

Q1. Vanadium is an important metal. Ferrovandium, an alloy of iron and vanadium, is used to make a strong type of vanadium-steel. Pure vanadium is used in nuclear reactors.

(a) The table shows some standard enthalpy of formation data.

	$V_2O_5(s)$	$CaO(s)$
$\Delta H_f^\ominus / \text{kJ mol}^{-1}$	-1560	-635

In the oldest method of extraction of vanadium, V_2O_5 is reacted with calcium at a high temperature.



Use data from the table and the equation to calculate the standard enthalpy change for this reaction.

State the type of reaction that V_2O_5 has undergone.

Suggest **one** major reason why this method of extracting vanadium is expensive, other than the cost of heating the reaction mixture.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(5)

(b) Ferrovandium is produced by the reaction of aluminium with a mixture of V_2O_5 and iron(III) oxide.

Write an equation for the reaction of aluminium with iron(III) oxide.

State the change in oxidation state of aluminium in this reaction.

.....
.....
.....
.....

(2)

- (c) Pure vanadium, for nuclear reactors, is formed by the reaction of hydrogen with purified VCl_2

Write an equation for this reaction in which the only other product is HCl gas.

Identify **two** hazards in this process, other than the fact that it operates at a high temperature.

Deduce why this process produces **pure** vanadium, other than the fact that purified VCl_2 is used.

.....
.....
.....
.....
.....
.....
.....
.....
.....

(4)

(Total 11 marks)

- Q2.(a)** Write an equation, including state symbols, for the reaction with enthalpy change equal to the standard enthalpy of formation for $\text{CF}_4(\text{g})$.

.....

(1)

- (b) Explain why CF_4 has a bond angle of 109.5° .

.....

.....

.....

.....

.....


(2)

(c) **Table 1** gives some values of standard enthalpies of formation ($\Delta_f H^\ominus$).

Table 1

Substance	F ₂ (g)	CF ₄ (g)	HF(g)
$\Delta_f H^\ominus / \text{kJ mol}^{-1}$	0	-680	-269

The enthalpy change for the following reaction is $-2889 \text{ kJ mol}^{-1}$.




Use this value and the standard enthalpies of formation in **Table 1** to calculate the standard enthalpy of formation of C₂H₆(g).

Standard enthalpy of formation of C₂H₆(g) = kJ mol⁻¹

(3)

(d) Methane reacts violently with fluorine according to the following equation.



Some mean bond enthalpies are given in **Table 2**.

Table 2

Bond	C-H	C-F	H-F
Mean bond enthalpy / kJ	412	484	562

mol ⁻¹			
-------------------	--	--	--

A student suggested that one reason for the high reactivity of fluorine is a weak F–F bond.

Is the student correct? Justify your answer with a calculation using these data.

.....

(4)
 (Total 10 marks)

Q3.Antimony is a solid element that is used in industry. The method used for the extraction of antimony depends on the grade of the ore.

(a) Antimony can be extracted by reacting scrap iron with low-grade ores that contain antimony sulfide (Sb₂S₃).

(i) Write an equation for the reaction of iron with antimony sulfide to form antimony and iron(II) sulfide.

.....

(1)

(ii) Write a half-equation to show what happens to the iron atoms in this reaction.

.....

(1)

(b) In the first stage of the extraction of antimony from a high-grade ore, antimony sulfide is roasted in air to convert it into antimony(III) oxide (Sb₂O₃) and sulfur dioxide.

(i) Write an equation for this reaction.

.....

(1)

- (ii) Identify **one** substance that is manufactured directly from the sulfur dioxide formed in this reaction.

.....

(1)

- (c) In the second stage of the extraction of antimony from a high-grade ore, antimony(III) oxide is reacted with carbon monoxide at high temperature.

- (i) Use the standard enthalpies of formation in the table and the equation given below the table to calculate a value for the standard enthalpy change for this reaction.

	Sb₂O₃(s)	CO(g)	Sb(l)	CO₂(g)
$\Delta H_{f\ominus} / \text{kJ mol}^{-1}$	-705	-111	+20	-394



.....

(3)

- (ii) Suggest why the value for the standard enthalpy of formation of liquid antimony, given in the table above, is **not** zero.

.....

(1)

- (iii) State the type of reaction that antimony(III) oxide has undergone in this reaction.

.....

(1)

- (d) Deduce **one** reason why the method of extraction of antimony from a low-grade ore, described in part (a), is a low-cost process. Do **not** include the cost of the ore.

.....
.....

(1)
(Total 10 marks)

Q4. The alcohol 2-methylpropan-2-ol, $(\text{CH}_3)_3\text{COH}$, reacts to form esters that are used as flavourings by the food industry. The alcohol can be oxidised to produce carbon dioxide and water.

A student carried out an experiment on a pure sample of 2-methylpropan-2-ol to determine its enthalpy of combustion. A sample of the alcohol was placed into a spirit burner and positioned under a beaker containing 50 cm^3 of water. The spirit burner was ignited and allowed to burn for several minutes before it was extinguished.

The results for the experiment are shown in **Table 1**.

Table 1

Initial temperature of the water / °C	18.1
Final temperature of the water / °C	45.4
Initial mass of spirit burner and alcohol / g	208.80
Final mass of spirit burner and alcohol / g	208.58

- (a) Use the results from **Table 1** to calculate a value for the heat energy released from the combustion of this sample of 2-methylpropan-2-ol. The specific heat capacity of water is $4.18 \text{ J K}^{-1} \text{ g}^{-1}$. Show your working.

.....
.....
.....
.....
.....

(2)

- (b) Calculate the amount, in moles, of 2-methylpropan-2-ol burned in the experiment. Hence calculate a value, in kJ mol^{-1} , for the enthalpy of combustion of 2-methylpropan-2-ol. Show your working.

(If you were unable to calculate an answer to part (a), you should assume that the heat energy released was 5580 J. This is **not** the correct value.)

.....

.....

.....

.....

.....

(3)

- (c) An equation for the combustion of 2-methylpropan-2-ol is



Table 2 contains some standard enthalpy of formation data.

Table 2

	$(\text{CH}_3)_3\text{COH}(\text{l})$	$\text{O}_2(\text{g})$	$\text{CO}_2(\text{g})$	$\text{H}_2\text{O}(\text{l})$
$\Delta H_f^\ominus / \text{kJ mol}^{-1}$	-360	0	-393	-286

Use the data from **Table 2** to calculate a value for the standard enthalpy of combustion of 2-methylpropan-2-ol. Show your working.

.....

.....

.....

.....

.....

.....

.....

.....

(3)

- (d) An accurate value for the enthalpy of combustion of 2-methylpropan-2-ol in which water is formed as a gas is $-2422 \text{ kJ mol}^{-1}$.

Use this value and your answer from part (b) to calculate the overall percentage error in the student's experimental value for the enthalpy of combustion of 2-methylpropan-2-ol.

.....
.....
.....

(1)

- (e) Suggest **one** improvement that would reduce errors due to heat loss in the student's experiment.

.....
.....
.....

(1)

- (f) Suggest **one** other source of error in the student's experiment. Do **not** include heat loss, apparatus error or student error.

.....
.....
.....

(1)


(Total 11 marks)

Q5. Group 2 metals and their compounds are used commercially in a variety of processes.

- (a) Strontium is extracted from strontium oxide (SrO) by heating a mixture of powdered strontium oxide and powdered aluminium.

Consider these standard enthalpies of formation.

	SrO(s)	Al ₂ O ₃ (s)
$\Delta H_f^\ominus / \text{kJ mol}^{-1}$	- 590	- 1669



Use these data and the equation to calculate the standard enthalpy change for this extraction of strontium.

The use of powdered strontium oxide and powdered aluminium increases the surface area of the reactants.

Suggest **one** reason why this increases the reaction rate.

Suggest **one** major reason why this method of extracting strontium is expensive.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(5)

(b) Explain why calcium has a higher melting point than strontium.

.....

.....

.....

.....

.....

.....

.....
.....

(2)

- (c) Magnesium is used in fireworks. It reacts rapidly with oxygen, burning with a bright white light. Magnesium reacts slowly with cold water.

Write an equation for the reaction of magnesium with oxygen.

Write an equation for the reaction of magnesium with cold water.

Give a medical use for the magnesium compound formed in the reaction of magnesium with cold water.

.....
.....
.....
.....
.....
.....
.....
.....

(3)

(Total 10 marks)