


M1.(a) (nucleophilic) addition-elimination
Not electrophilic addition-elimination

1



Allow C₆H₅ or benzene ring

Allow attack by :NH₂C₆H₅

M2 not allowed independent of M1, but allow M1 for correct attack on C⁺

M3 for correct structure with charges but lone pair on O is part of M4

M4 (for three arrows and lone pair) can be shown in more than one structure

4

(b) **The minimum quantity of hot water was used:**

To ensure the hot solution would be saturated / crystals would form on cooling

1

The flask was left to cool before crystals were filtered off:

Yield lower if warm / solubility higher if warm

1

The crystals were compressed in the funnel:

Air passes through the sample not just round it

Allow better drying but not water squeezed out

1

A little cold water was poured through the crystals:

To wash away soluble impurities

1

(c) Water

Do not allow unreacted reagents

1

Press the sample of crystals between filter papers

Allow give the sample time to dry in air

1

(d) M_r product = 135.0

1

$$\text{Expected mass} = 5.05 \times \frac{135.0}{93.0} = 7.33 \text{ g}$$


1

$$\text{Percentage yield} = \frac{4.82}{7.33} \times 100 = 65.75 = 65.8(\%)$$


Answer must be given to this precision

1

(e)



OR



1

(f) Electrophilic substitution

1

(g) Hydrolysis

1

(h) Sn / HCl

Ignore acid concentration; allow Fe / HCl

1

[18]

M2.(a) (Q = mcΔT)

$$= 50 \times 4.18 \times 27.3$$

If incorrect (eg mass = 0.22 or 50.22 g) CE = 0 / 2

1

$$= \mathbf{5706 \text{ J}}$$
 (accept 5700 and 5710)

Accept 5.7 kJ with correct unit. Ignore sign.

1

(b) M_r of 2-methylpropan-2-ol = 74(.0)

For incorrect M_r , lose M1 but mark on.

1

$$\begin{aligned} \text{Moles} &= \text{mass} / M_r \\ &= 0.22 / 74(.0) \\ &= \mathbf{0.00297 \text{ moles}} \end{aligned}$$

1

$$\Delta H = -5706 / (0.002970 \times 1000)$$

$$= \mathbf{-1921 \text{ (kJ mol}^{-1}\text{)}}$$

If 0.22 is used in part (a), answer = -8.45 kJ mol⁻¹ scores 3

(Allow -1920, -1919)

If uses the value given (5580 J), answer = -1879 kJ mol⁻¹ scores 3

Answer without working scores M3 only.

Do not penalise precision.

Lack of negative sign loses M3

1

- (c) $\Delta H = \Sigma \Delta H \text{ products} - \Sigma \Delta H \text{ reactants}$
OR a correct cycle

Correct answer with no working scores 1 mark only.

1

$$\Delta H = -(-360) + (4 \times -393) + (5 \times -286)$$

M2 also implies M1 scored.

1

$$\Delta H = \mathbf{-2642 \text{ (kJ mol}^{-1}\text{)}}$$
 This answer only.

Allow 1 mark out of 3 for correct value with incorrect sign.

1

- (d) $(-2422 - \text{part (b)}) \times 100 / -2422$
Ignore negative sign.

Expect answers in region of 20.7

If error carried forward, 0.22 allow 99.7

If 5580 J used earlier, then allow 22.4

1

- (e) Reduce the distance between the flame and the beaker / put a sleeve around the flame to protect from drafts / add a lid / use a copper calorimeter rather than a pyrex beaker / use a food calorimeter

Any reference to insulating material around the beaker must be on top.

Accept calibrate the equipment using an alcohol of known enthalpy of combustion.

1


- (f) Incomplete combustion

1

[11]

M3.(a) moles of $\text{Cr}_2\text{O}_7^{2-}$ per titration = $21.3 \times 0.0150 / 1000 = \underline{3.195 \times 10^{-4}}$

1



If 1:6 ratio incorrect cannot score M2 or M3

1

moles of $\text{Fe}^{2+} = 6 \times 3.195 \times 10^{-4} = 1.917 \times 10^{-3}$

Process mark for M1 $\times 6$ (also score M2)

1

original moles in $250 \text{ cm}^3 = 1.917 \times 10^{-3} \times 10 = 1.917 \times 10^{-2}$

Process mark for M3 $\times 10$

1

mass of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O} = 1.917 \times 10^{-2} \times 277.9 = 5.33 \text{ (g)}$

Mark for answer to M4 $\times 277.9$

(allow 5.30 to 5.40)

*Answer **must** be to at least 3 sig figs*

Note that an answer of 0.888 scores M1, M4 and M5 (ratio 1:1 used)

1

- (b) (Impurity is a) reducing agent / reacts with dichromate / impurity is a version of FeSO_4 with fewer than 7 waters (not fully hydrated)

Allow a reducing agent or compound that that converts Fe^{3+} into Fe^{2+}

1

Such that for a given mass, the impurity would react with more dichromate than a similar mass of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

OR for equal masses of the impurity and $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, the impurity would react with more dichromate.

Must compare mass of impurity with mass of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

1

[7]

- M4.(a)** This question is marked using levels of response. Refer to the Mark Scheme Instructions for Examiners for guidance on how to mark this question.

All stages are covered and the explanation of each stage is generally correct and virtually complete.

Answer is communicated coherently and shows a logical progression from stage 1 and stage 2 to stage 3. Steps in stage 3 must be complete, ordered and include a comparison.

Level 3
5 – 6 marks

All stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies OR two stages are covered and the explanations are generally correct and virtually complete.

Answer is mainly coherent and shows a progression from stage 1 and stage 2 to stage 3.

Level 2
3 – 4 marks

Two stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies, OR only one stage is covered but the explanation is generally correct and virtually complete.

Answer includes some isolated statements, but these are not presented in a logical order or show confused reasoning.

Level 1
1 – 2 marks

Insufficient correct Chemistry to warrant a mark.

Level 0

Indicative Chemistry content**Stage 1:** difference in structure of the two acids

- The acids are of the form RCOOH
- but in ethanoic acid R = CH₃
- whilst in ethanedioic acid R = COOH

Stage 2: the inductive effect

- The unionised COOH group contains two very electronegative oxygen atoms
- therefore has a negative inductive (electron withdrawing) effect
- The CH₃ group has a positive inductive (electron pushing) effect

Stage 3: how the polarity of OH affects acid strength

- The O–H bond in the ethanedioic acid is more polarised / H becomes more δ⁺
- More dissociation into H⁺ ions
- Ethanedioic acid is stronger than ethanoic acid

6

(b) Moles of NaOH = Moles of HOCCOO⁻ formed = 6.00×10^{-2} *Extended response*

1

Moles of HOCCOOH remaining = $1.00 \times 10^{-1} - 6.00 \times 10^{-2}$ = 4.00×10^{-2}

1

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$[\text{H}^+] = K_a \times \frac{[\text{HA}]}{[\text{A}^-]}$$

1



$$[\text{H}^+] = 5.89 \times 10^{-2} \times (4.00 \times 10^{-2} / V) / (6.00 \times 10^{-2} / V) = 3.927 \times 10^{-2}$$

1

$$\text{pH} = -\log_{10}(3.927 \times 10^{-2}) = 1.406 = 1.41$$

Answer must be given to this precision

1



1

Moles of $\text{KMnO}_4 = 20.2 \times 2.00 \times 10^{-2} / 1000 = 4.04 \times 10^{-4}$

1

Moles of $\text{H}_2\text{C}_2\text{O}_4 = 5 / 2 \times 4.04 \times 10^{-4} = 1.01 \times 10^{-3}$

1

Concentration = moles / volume (in dm^3)

$= 1.01 \times 10^{-3} \times 1000 / 25 = 4.04 \times 10^{-2} \text{ (mol dm}^{-3}\text{)}$

If 1:1 ratio or incorrect ratio used, M2 and M4 can be scored


1

[15]

M5.(a) Other product in equation is water

If product incorrect, CE = 0 / 2

1



Allow multiples, including fractions.

Ignore state symbols.

1

(b) Named indicator paper placed in gas / add named indicator to gas / collect gas and add named indicator

If indicator not named, CE = 0 / 2

Lose this mark if the indicator is added to the reaction mixture. Can still score the second mark.

1

Correct full colour change

If universal indicator is used, allow 'green to blue / purple' or

'yellow to blue / purple'.

If litmus is used, allow 'purple to blue' or 'red to blue'.

Allow one mark overall for 'add universal indicator' and 'turns purple / blue'.

Allow one mark overall for 'add litmus' and 'turns blue'.

1

[4]