




Q1. Ammonia is used to make nitric acid (HNO₃) by the Ostwald Process.
Three reactions occur in this process.



- (a) In one production run, the gases formed in Reaction 1 occupied a total volume of 4.31 m³ at 25 °C and 100 kPa.

Calculate the amount, in moles, of NO produced.

Give your answer to 3 significant figures.

(The gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$)

.....
.....
.....
.....
.....
.....

(Extra space)
.....
.....

(4)

- (b) In another production run, 3.00 kg of ammonia gas were used in Reaction 1 and all of the NO gas produced was used to make NO₂ gas in Reaction 2.

- (i) Calculate the amount, in moles, of ammonia in 3.00 kg.

.....
.....
.....
.....

(2)


- (ii) Calculate the mass of NO₂ formed from 3.00 kg of ammonia in Reaction 2 assuming an 80.0% yield.
Give your answer in kilograms.
(If you have been unable to calculate an answer for part (b)(i), you may assume a value of 163 mol. This is **not** the correct answer.)

.....
.....
.....
.....
.....
.....

(Extra space)
.....
.....

(3)

- (c) Consider Reaction 3 in this process.



Calculate the concentration of nitric acid produced when 0.543 mol of NO₂ is reacted with water and the solution is made up to 250 cm³.

.....
.....
.....
.....
.....

(Extra space)
.....

(2)

- (d) Suggest why a leak of NO₂ gas from the Ostwald Process will cause atmospheric pollution.

.....
.....


(1)

(e) Give **one** reason why excess air is used in the Ostwald Process.

.....
.....

(1)

(f) Ammonia reacts with nitric acid as shown in this equation.




Deduce the type of reaction occurring.

.....

(1)

(Total 14 marks)

Q2. (a) Boron trichloride (BCl_3) can be prepared as shown by the following equation.



A sample of boron oxide (B_2O_3) was reacted completely with carbon and chlorine. The two gases produced occupied a total volume of 5000 cm^3 at a pressure of 100 kPa and a temperature of 298 K.

Calculate the mass of boron oxide that reacted.
Give your answer to 3 significant figures.

(The gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$)

.....
.....
.....

.....
.....
.....
.....
.....
.....

(Extra space)
.....
.....
.....

(5)

(b) Boron trichloride can also be prepared from its elements.

Write an equation for this reaction.
Explain why boron trichloride has a trigonal planar shape with equal bond angles.

.....
.....
.....
.....

(Extra space)
.....

(3)

(c) (i) Boron trichloride is easily hydrolysed to form two different acids as shown in the following equation.



Calculate the concentration, in mol dm⁻³, of hydrochloric acid produced when 43.2 g of boron trichloride are added to water to form 500 cm³ of solution. Give your answer to 3 significant figures.

.....

.....
.....
.....
.....
.....
.....
.....
.....
(Extra space)


(4)

(ii) Boric acid (H₃BO₃) can react with sodium hydroxide to form sodium borate and water.
Write an equation for this reaction.

.....

(1)

(d) Boron trichloride can be reduced by using hydrogen to form pure boron.



Calculate the percentage atom economy for the formation of boron in this reaction.

Apart from changing the reaction conditions, suggest **one** way a company producing pure boron could increase its profits from this reaction.

.....
.....
.....
.....
.....
.....

(Extra space)

(3)

- (e) A different compound of boron and chlorine has a relative molecular mass of 163.6 and contains 13.2% of boron by mass.

Calculate the molecular formula of this compound.
Show your working.

.....

.....

.....

.....

.....

.....

.....


.....

(Extra space)

.....

(4)
(Total 20 marks)

Q3. Solid iron(II) ethanedioate dihydrate ($\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$) has a polymeric structure.
Two repeating units in the polymer chain are shown.



- (a) Name the type of bond that is represented by the arrows.

.....

(1)

- (b) In terms of electrons explain how the water molecules, **not** shown in the diagram, form bonds to the iron.

.....
.....
.....
.....

(2)

- (c) Predict the value of the bond angle between the two bonds to iron that are formed by these two water molecules.



.....

(1)

- (d) Iron(II) ethanedioate dihydrate can be analysed by titration using potassium manganate(VII) in acidic solution. In this reaction, manganate(VII) ions oxidise iron(II) ions and ethanedioate ions.

A 1.381 g sample of impure $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ was dissolved in an excess of dilute sulfuric acid and made up to 250 cm^3 of solution.
 25.0 cm^3 of this solution decolourised 22.35 cm^3 of a $0.0193 \text{ mol dm}^{-3}$ solution of potassium manganate(VII).

- (i) Use the half-equations given below to calculate the reacting ratio of moles of manganate(VII) ions to moles of iron(II) ethanedioate.



.....
.....
.....

(1)

- (ii) Calculate the percentage by mass of $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ in the original sample.

(If you have been unable to answer part (d)(i) you may assume that three moles of manganate(VII) ions react with seven moles of iron(II) ethanedioate. This is **not** the correct ratio.)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(5)
(Total 10 marks)

Q4.A student calculated that a value for the enthalpy change of neutralisation is $-51.2 \text{ kJ mol}^{-1}$.

The design of a possible hand-warmer using hydrochloric acid and sodium hydroxide was discussed. It was proposed that 500 cm^3 of hydrochloric acid should be used in a flexible, sealed plastic container with a breakable tube of solid sodium hydroxide also in the container. On breaking the tube, the sodium hydroxide would be released, react with the acid and produce heat.

A $40 \text{ }^\circ\text{C}$ temperature rise was thought to be suitable.

- (a) Calculate the heat energy, in J, required to raise the temperature of the reaction mixture by $40 \text{ }^\circ\text{C}$. Assume that the reaction mixture has a density of 1.00 g cm^{-3} and a specific heat capacity of $4.18 \text{ J K}^{-1} \text{ g}^{-1}$. Assume that all of the heat energy given out is used to heat the reaction mixture.

.....

.....

.....
.....

(2)

- (b) Use your answer from part (a) and the value for the enthalpy change of neutralisation of $-51.2 \text{ kJ mol}^{-1}$ to calculate the minimum amount, in moles, and hence the minimum mass of sodium hydroxide required in the breakable tube. (If you could not complete the calculation in part (a) assume that the heat energy required was 77 400 J. This is **not** the correct answer).

Show your working.

Moles of NaOH

.....

Mass of NaOH

.....

(3)

- (c) Use the amount, in moles, of sodium hydroxide from part (b) to calculate the minimum concentration, in mol dm^{-3} , of hydrochloric acid required in the 500 cm^3 of solution used in the sealed container.

.....

.....

.....

(1)

- (d) Suggest **one** possible risk to a person who uses a hand-warmer containing sodium hydroxide and hydrochloric acid.

.....

.....

(1)

- (e) A commercial hand-warmer uses powdered iron sealed in a plastic container. A valve allows air to enter the container, and oxygen in the air reacts slowly with the iron to form solid iron(III) oxide. The heat released warms the container.

- (i) Write an equation for this reaction between iron and oxygen to form iron(III)

oxide.

.....

(1)

- (ii) One version of an iron-oxygen hand-warmer advertises that it is designed to stay warm for up to four hours. Other than by increasing the amount of iron in the container, state **one** change to the iron in the hand-warmer that would increase this time. Explain why this change to the iron might **not** be an advantage.

Change to the iron

.....

Explanation

.....

.....

(3)

- (f) Another type of hand-warmer uses sodium thiosulfate. Sodium thiosulfate is very soluble in water at 80 °C but is much less soluble at room temperature. When a hot, concentrated solution of sodium thiosulfate is cooled it does not immediately crystallise. The sodium thiosulfate stays dissolved as a stable 'super-saturated' solution until crystallisation is triggered. Heat energy is then released when the sodium thiosulfate crystallises.

- (i) This type of hand-warmer is re-usable. Suggest **one** environmental advantage that a sodium thiosulfate hand-warmer has over the other two types.

.....

(1)

- (ii) Describe the **two** steps that you would take to make the sodium thiosulfate hand-warmer ready for re-use.

Step 1

.....

Step 2

.....

(2)

(Total 14 marks)

