M1. (a) enthalpy change/ heat energy change when 1 mol of a substance

1

is completely burned in oxygen

1

at 298K and 100 kPa or standard conditions

1

(not 1atm)

(b)  $\Delta H = \sum \text{bonds broken} - \sum \text{bonds formed}$ 

1

$$= (6 \times 412) + 612 + 348 + (4.5 \times 496) - ((6 \times 743) + (6 \times 463))$$

1

1

(c) by definition  $\Delta H_i$  is formation from an element

1

(d)  $\Delta H_c = \sum \Delta H_t$  products  $-\sum \Delta H_t$  reactants or cycle

1

$$= (3 \times -394) + (3 \times -242) - (+20)$$

1

1

(e) bond enthalpies are mean/average values

1

from a range of compounds

[12]

M2. (a) enthalpy (or energy) to break (or dissociate) a bond; 1 averaged over different molecules (environments); 1 enthalpy (or heat energy) change when one mole of a compound; 1 is formed from its elements; 1 in their standard states; 1 enthalpy change =  $\Sigma$ (bonds broken) –  $\Sigma$ (bonds formed) or cycle; (b) 1  $= 4 \times 388 + 163 + 2 \times 146 + 4 \times 463 - (944 + 8 \times 463);$ (or similar) 1 =-789; (+ 789 scores 1 only) 1 (c) (i) zero; 1 (ii)  $AH = \Sigma$  (enthalpies of formation of products)  $-\Sigma$  (enthalpies of formation of reactants) 1  $= 4 \times -242 - (75 + 2 \times -133);$ 1 =-777;(+ 777 scores one only) (d) mean bond enthalpies are not exact (or indication that actual values are different from real values) 1

[13]

M3.  $\Delta H = \Sigma$ (bonds broken) –  $\Sigma$ (bonds formed) (or cycle) (a) 1 = +146 - 496/2 (or  $2 \times 463 + 146 - (2 \times 463 + 496/2)$ 1  $= -102 \text{ (kJ mol}^{-1}) (1)$ (accept no units, wrong units loses a mark; +102 scores (1) 1  $C(s) + 2H_2(g) \rightarrow CH_4(g)$  equation (1) Correct state symbols (1) (b) 2 Macromolecular (c) (i) (accept giant molecule or carbon has many (4) bonds) 1 (ii)  $\Delta H = \Sigma \Delta H_t$ (products) –  $\Sigma \Delta H_t$ (reactants) (or cycle) 1  $= 715 + 4 \times 218 - (-74.9)$ 1 = 1662 (kJ mol<sup>-1</sup>) (accept no units, wrong units loses one mark, allow 1660 to 1663, -1662 scores one mark only) 1 (iii) 1662/4 = 415.5(mark is for divide by four, allow if answer to (c)(ii) is wrong) 1 [10]

**M4.**D

[1]

## **M5.** (a) (Energy required) to break a given <u>covalent</u> bond **(1)** averaged over a range of compounds **(1)**

Penalise first mark if 'energy' / 'enthalpy' evolved

2

(b) (i) 
$$4 \times C - H = 4 \times 413 = +1652$$
  
 $1 \times C - C = 1 \times 347 = 347$   
 $1 \times C = O = 1 \times 736 = 736$   
 $2\frac{1}{2} \times O = O = 2.5 \times 498 = 1245$  (1)  
 $= 2735 + 1245 = +3980$  (1)

first mark for 4:1:1 or 2735 ignore sign

(ii) 
$$4 \times H-O = -4 \times 464 = -1856$$
  
 $4 \times C-O = -4 \times 736 = -2944$  (1)  
 $= -4800$  (1)

First mark for 4:4

(iii) 
$$\Delta H_R = \Sigma Bonds \ broken - \Sigma Bonds \ made$$
  
= +3980 - 4800 = -820 **(1)**  
Conseq Mark for incorrect answers in (i) and (ii) as  
(i) Answer + (ii) Answer =

[7]

5

M6.(a) Enthalpy (Energy) to break a (covalent) bond (1) OR dissociation energy Varies between compounds so average value used (1) QL mark

OR average of dissociation energies in a single molecule / e.g.  $CH_4$ 

Do not allow mention of energy to form bonds but with this case can allow second mark otherwise 2<sup>nd</sup> mark consequential on first

2

(b) (i) 
$$1/2 N_2 + 3/2 H_2 \rightarrow NH_3$$
 (1)

## Ignore s s

```
(ii) \Delta H = (\Sigma)bonds broken -(\Sigma)bonds formed (1)
= 1/2 \times 944 + 3/2 \times 436 - 3 \times 388 (1)
= -38 kJ mol<sup>-1</sup> (1)
Ignore no units, penalise wrong units
Score 2/3 for -76
1/3 for +38
Allow 1/3 for +76
```

4

3

(c) 
$$4 (C-H) + (C=C) + (H-H) - (6 (C-H) + (C-C)) = -136$$
 (1) OR  $(C=C) + (H-H) - ((C-C) + 2 (C-H)) = -136$  2  $(C-H) = 836$  (1)  $(C-H) = 418$  (kJ mol<sup>-1</sup>) (1) Note: allow (1) for  $-836$  another (1) for  $-418$ 

[9]