

**Q1.** This question is about the pH of several solutions.

Give all values of pH to 2 decimal places.

(a) (i) Write an expression for pH.

.....

(1)

(ii) Calculate the pH of 0.154 mol dm<sup>-3</sup> hydrochloric acid.

.....

.....

(1)

(iii) Calculate the pH of the solution formed when 10.0 cm<sup>3</sup> of 0.154 mol dm<sup>-3</sup> hydrochloric acid are added to 990 cm<sup>3</sup> of water.

.....

.....

.....

.....

(2)

(b) The acid dissociation constant,  $K_a$ , for the weak acid HX has the value  $4.83 \times 10^{-5}$  mol dm<sup>-3</sup> at 25 °C.  
A solution of HX has a pH of 2.48

Calculate the concentration of HX in the solution.

.....

.....

.....

.....

.....

.....

(4)

- (c) Explain why the pH of an acidic buffer solution remains almost constant despite the addition of a small amount of sodium hydroxide.

.....

.....

.....

.....

.....

.....

(2)

- (d) The acid dissociation constant,  $K_a$ , for the weak acid HY has the value  $1.35 \times 10^{-5} \text{ mol dm}^{-3}$  at  $25^\circ\text{C}$ .

A buffer solution was prepared by dissolving 0.0236 mol of the salt NaY in  $50.0 \text{ cm}^3$  of a  $0.428 \text{ mol dm}^{-3}$  solution of the weak acid HY

- (i) Calculate the pH of this buffer solution.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(4)

- (ii) A  $5.00 \times 10^{-4} \text{ mol}$  sample of sodium hydroxide was added to this buffer solution.

Calculate the pH of the buffer solution after the sodium hydroxide was added.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

(4)  
(Total 18 marks)

**Q2.** In this question, give all values of pH to 2 decimal places.

(a) (i) Write an expression for the term pH.

.....

(1)

(ii) Calculate the concentration, in mol dm<sup>-3</sup>, of an aqueous solution of sulfuric acid that has a pH of 0.25

.....  
.....  
.....  
.....

(2)

(b) A student carried out a titration by adding an aqueous solution of sodium hydroxide from a burette to an aqueous solution of ethanoic acid. The end-point was reached

when 22.60 cm<sup>3</sup> of the sodium hydroxide solution had been added to 25.00 cm<sup>3</sup> of 0.410 mol dm<sup>-3</sup> ethanoic acid.

- (i) Write an equation for the reaction between sodium hydroxide and ethanoic acid.

.....

(1)

- (ii) Calculate the concentration, in mol dm<sup>-3</sup>, of the sodium hydroxide solution used.

.....

.....

.....

.....

(2)

- (iii) A list of indicators is shown below.

Indicator	pH range
thymol blue	1.2–2.8
bromophenol blue	3.0–4.6
litmus	5.0–8.0
cresol purple	7.6–9.2

Select from the list the most suitable indicator for the end-point of this titration.

.....

(1)

- (iv) Suggest why the concentration of sodium hydroxide in a solution slowly decreases when left open to air.

.....

.....

(1)

(c) At 298 K, the value of the acid dissociation constant,  $K_a$ , for ethanoic acid in aqueous solution is  $1.74 \times 10^{-5} \text{ mol dm}^{-3}$

(i) Write an expression for the acid dissociation constant,  $K_a$ , for ethanoic acid.

.....  
.....

(1)

(ii) Calculate the pH of  $0.410 \text{ mol dm}^{-3}$  ethanoic acid at this temperature.

.....  
.....  
.....  
.....

(3)

(iii) Calculate the pH of the buffer solution formed when  $10.00 \text{ cm}^3$  of  $0.100 \text{ mol dm}^{-3}$  potassium hydroxide are added to  $25.00 \text{ cm}^3$  of  $0.410 \text{ mol dm}^{-3}$  ethanoic acid.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

(6)

(Total 18 marks)

**Q3.**In this question, give all pH values to 2 decimal places.

- (a) (i) Write expressions for the ionic product of water,  $K_w$ , and for pH.

$K_w =$  .....

pH = .....

- (ii) At 318 K, the value of  $K_w$  is  $4.02 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$  and hence the pH of pure water is 6.70  
State why pure water is not acidic at 318 K.

.....  
.....

- (iii) Calculate the number of moles of sodium hydroxide in  $2.00 \text{ cm}^3$  of  $0.500 \text{ mol dm}^{-3}$  aqueous sodium hydroxide.

.....  
.....

- (iv) Use the value of  $K_w$  given above and your answer to part (a)(iii) to calculate the pH of the solution formed when  $2.00 \text{ cm}^3$  of  $0.500 \text{ mol dm}^{-3}$  aqueous sodium hydroxide are added to  $998 \text{ cm}^3$  of pure water at 318 K.

.....  
.....  
.....  
.....  
.....

(6)

(b) At 298 K, the acid dissociation constant,  $K_a$ , for propanoic acid,  $\text{CH}_3\text{CH}_2\text{COOH}$ , has the value  $1.35 \times 10^{-5} \text{ mol dm}^{-3}$ .

(i) Write an expression for  $K_a$  for propanoic acid.


.....  
.....

(ii) Calculate the pH of  $0.125 \text{ mol dm}^{-3}$  aqueous propanoic acid at 298 K.

.....  
.....  
.....  
.....  
.....  
.....

(4)

(c) Sodium hydroxide reacts with propanoic acid as shown in the following equation.



A buffer solution is formed when sodium hydroxide is added to an excess of aqueous propanoic acid.

(i) Calculate the number of moles of propanoic acid in  $50.0 \text{ cm}^3$  of  $0.125 \text{ mol dm}^{-3}$  aqueous propanoic acid.

.....  
.....

(ii) Use your answers to part (a)(iii) and part (c)(i) to calculate the number of moles of propanoic acid in the buffer solution formed when  $2.00 \text{ cm}^3$  of  $0.500 \text{ mol dm}^{-3}$  aqueous sodium hydroxide are added to  $50.0 \text{ cm}^3$  of  $0.125 \text{ mol dm}^{-3}$  aqueous propanoic acid.

.....  
.....  
.....

(iii) Hence calculate the pH of this buffer solution at 298 K.

.....  
.....  
.....  
.....  
.....  
.....

(6)  
(Total 16 marks)