

1

(ii) to speed up the reaction

OR

(provide a) catalyst or catalyses the reaction or biological catalyst

OR


release / contain / provides an enzyme

Ignore "fermentation"

Ignore "to break down the glucose"

Not simply "enzyme" on its own

1



1

(ii) Secondary (alcohol) OR 2° (alcohol)

1

(c) **M1** $q = m c \Delta T$

OR $q = 150 \times 4.18 \times 8.0$

Award full marks for correct answer

*In **M1**, do not penalise incorrect cases in the formula*

M2 = (±) 5016 (J) **OR** 5.016 (kJ) **OR** 5.02 (kJ)
 (also scores M1)

M3 This mark is for dividing correctly the number of kJ by the number of moles and arriving at a final answer in the range shown.
 Using 0.00450 mol

therefore $\Delta H = - \underline{1115}$ (kJ mol⁻¹)

OR $- \underline{1114.6}$ to $- \underline{1120}$ (kJ mol⁻¹)

Range (+)1114.6 to (+)1120 gains 2 marks

BUT - 1110 gains 3 marks and +1110 gains 2 marks

AND – 1100 gains 3 marks and +1100 gains 2 marks

Award full marks for correct answer

In M1, do not penalise incorrect cases in the formula

Penalise M3 ONLY if correct numerical answer but sign is incorrect; (+)1114.6 to (+)1120 gains 2 marks

Penalise M2 for arithmetic error and mark on

If $\Delta T = 281$; score $q = m c \Delta T$ only

If $c = 4.81$ (leads to 5772) penalise M2 ONLY and mark on for M3 = - 1283

Ignore incorrect units in M2

If units are given in M3 they must be either kJ or kJ mol⁻¹ in this case

3

(d) **M1** The enthalpy change / heat change at constant pressure when 1 mol of a compound / substance / element

M2 is burned / combusts / reacts completely in oxygen
OR
burned / combusted / reacted in excess oxygen

M3 with (all) reactants and products / (all) substances in standard / specified states

OR

(all) reactants and products / (all) substances in normal states under standard conditions / 100 kPa / 1 bar and specified T / 298 K

For M3

Ignore reference to 1 atmosphere

3

(e) **M1**

$\sum B(\text{reactants}) - \sum B(\text{products}) = \Delta H$

OR

Sum of bonds broken – Sum of bonds formed = ΔH

OR

2B(C–C) + B(C=O) + 6B(C–H) + 4B(O=O) (LHS)

– 6B(C=O) – 6B(O–H) (RHS) = ΔH

M2 (also scores **M1**)

2(348)+805+6(412)+4(496) [LHS = **5957**]

(696) (2472) (1984)

– 6(805) – 6(463) [RHS = (–) **7608**] = ΔH

(4830) (2778)

OR using only bonds broken and formed (5152 – 6803)

M3

$\Delta H = -1651$ (kJ mol⁻¹)

Candidates may use a cycle and gain full marks.

Correct answer gains full marks

Credit 1 mark for (+) 1651 (kJ mol⁻¹)

For other incorrect or incomplete answers, proceed as follows

- *check for an arithmetic error (AE), which is either a transposition error or an incorrect multiplication / addition error; this would score 2 marks (M1 and M2)*

- *If no AE, check for a correct method; this requires either a correct cycle with 4O₂, 3CO₂ and 3H₂O OR a clear statement of M1 which could be in words and scores only M1*

Allow a maximum of one mark if the only scoring point is LHS = 5957 (or 5152) OR RHS = 7608 (or 6803)

Award 1 mark for + 1651

3

(f) **For the two marks M1 and M2, any two from**

- heat loss or not all heat transferred to the apparatus or heat absorbed by the apparatus or (specific) heat capacity of the apparatus not considered
- incomplete combustion / not completely burned / reaction is not complete
- The idea that the water may end up in the gaseous state (rather than liquid)
- reactants and / or products may not be in standard states.
- MBE data refers to gaseous species but the enthalpy of combustion refers to liquids in their standard states / liquid propanone and liquid water in standard states
- MBE do not refer to specific compounds OR MBE values vary with different compounds / molecules OR are average / mean values taken from a range of compounds / molecules

Apply the list principle but ignore incomplete reasons that contain correct chemistry

Ignore "evaporation"

Ignore "faulty equipment"

Ignore "human error"

Not enough simply to state that "MBE are mean / average values"

2

[15]

M2.Increase in volume


If a volume is quoted it must be less than 300

1

Smaller increase in T above room temperature
Or increased contact between calorimeter and water
Or smaller heat loss by evaporation / from the surface

1

[2]



1

(b) Temperature rise = 20.1

$$q = 50.0 \times 4.18 \times 20.1 = 4201 \text{ (J)}$$

1

Mass of alcohol burned = 0.54 g and M_r alcohol = 100.0

$$\therefore \text{mol of alcohol} = n = 0.54 / 100 = 0.0054$$

1

Heat change per mole = $q / 1000n$ **OR** q / n

$$= 778 \text{ kJ mol}^{-1} \text{ **OR** } 778\,000 \text{ J mol}^{-1}$$

1

$$\Delta H = -778 \text{ kJ mol}^{-1} \text{ **OR** } -778\,000 \text{ J mol}^{-1}$$

M4 is for answer with negative sign for exothermic reaction

Units are tied to the final answer and must match

1

(c) Less negative than the reference

1

Heat loss **OR** incomplete combustion **OR** evaporation of alcohol **OR** heat transferred to beaker not taken into account

(d) Water has a known density (of 1.0 g cm^{-3})


1

Therefore, a volume of 50.0 cm^3 could be measured out

1

[9]

M4.(a)



Penalise C₂H₆O for ethanol in M1.

M2 and M3

Mark M2 and M3 independently.

Any **two** conditions in any order for **M2** and **M3** from

- (enzymes from) yeast or zymase
- $25 \text{ }^\circ\text{C} \leq T \leq 42 \text{ }^\circ\text{C}$ OR $298 \text{ K} \leq T \leq 315 \text{ K}$
- anaerobic / no oxygen / no air OR neutral pH

A lack of oxygen can mean either without oxygen or not having enough oxygen and does not ensure no oxygen, therefore only credit "lack of oxygen" if it is qualified.

Penalise 'bacteria', 'phosphoric acid', 'high pressure' using the list principle.

M4 (fractional) distillation or GLC

Ignore reference to 'aqueous' or 'water' (ie not part of the list principle).

M5 Carbon-neutral **in this context** means

There is no net / overall (annual) carbon dioxide / CO₂ emission to the atmosphere

OR

There is no change in the total amount / level of carbon dioxide / CO₂ present, in the atmosphere

For **M5** – must be about CO₂ and the atmosphere.
The idea that the carbon dioxide / CO₂ given out equals the carbon dioxide / CO₂ that was taken in from the atmosphere.

5

- (b) **M1** $q = m c \Delta T$ (this mark for correct mathematical formula)
*Full marks for **M1**, **M2** and **M3** for the correct answer.*
*In **M1**, do not penalise incorrect cases in the formula.*

$$\mathbf{M2} = (75 \times 4.18 \times 5.5)$$

$$1724 \text{ (J) OR } 1.724 \text{ (kJ) OR } 1.72 \text{ (kJ) OR } 1.7 \text{ (kJ)}$$

(also scores **M1**)

*Ignore incorrect units in **M2**.*

M3 Using 0.0024 mol

therefore $\Delta H = \underline{-718}$ (kJ mol⁻¹)

(Accept a range from -708 to -719 but do not penalise more than 3 significant figures)

*Penalise **M3** ONLY if correct numerical answer but sign is incorrect. Therefore **+718** gains two marks.*

*If units are quoted in **M3** they must be correct.*

*If $\Delta T = 278.5$, CE for the calculation and penalise **M2** and **M3**.*

M4 and **M5** in any order

Any **two** from

- incomplete combustion
- heat loss
- heat capacity of Cu not included
- some ethanol lost by evaporation
- not all of the (2.40×10^{-3} mol) ethanol is burned / reaction is incomplete
*If $c = 4.81$ (leads to 1984) penalise **M2** ONLY and mark on for **M3** = - 827*

5

- (c) (i) **M1** enthalpy / heat / energy change (at constant pressure) or enthalpy / heat / energy needed in breaking / dissociating (a) covalent bond(s)
Ignore bond making.

M2 averaged for that type of bond over different / a range of molecules /

compounds

Ignore reference to moles.

2

(ii) **M1**

$$\underline{\sum B(\text{reactants}) - \sum B(\text{products}) = \Delta H}$$

OR

$$\underline{\text{Sum of bonds broken} - \text{Sum of bonds formed} = \Delta H}$$

OR

$$\begin{aligned} & B(\text{C-C}) + B(\text{C-O}) + B(\text{O-H}) + 5B(\text{C-H}) + 3B(\text{O=O}) \\ & - 4B(\text{C=O}) - 6B(\text{O-H}) = \Delta H = -1279 \end{aligned}$$

Correct answer gains full marks.

Credit 1 mark for - 496 (kJ mol⁻¹)

For other incorrect or incomplete answers, proceed as follows

- *check for an arithmetic error (AE), which is either a transposition error or an incorrect multiplication; this would score 2 marks (M1 and M2).*

*If no AE, check for a correct method; this requires either a correct cycle with 2CO₂ and 3H₂O OR a clear statement of M1 which could be in words and scores **only M1**.*

M2 (also scores **M1**)

$$348+360+463+5(412)+ 3B(\text{O=O})$$

$$\begin{aligned} & \quad \quad \quad (3231) \quad \quad \quad (\text{or } 2768 \text{ if O-H cancelled}) \\ & - 4(805) - 6(463) = \Delta H = -1279 \end{aligned}$$

$$\quad \quad \quad (5998) \quad \quad \quad (\text{or } 5535 \text{ if O-H cancelled})$$

$$3B(\text{O=O}) = \underline{1488} \text{ (kJ mol}^{-1}\text{)}$$

Credit a maximum of one mark if the only scoring point is bonds formed adds up to 5998 (or 5535) OR bonds broken includes the calculated value of 3231 (or 2768).

M3

$$B(\text{O=O}) = \underline{496} \text{ (kJ mol}^{-1}\text{)}$$

Award 1 mark for -496

Students may use a cycle and gain full marks

3

[15]

M5.(a) Start a clock when KCl is added to water 1

Record the temperature every subsequent minute for about 5 minutes

*Allow record the temperature at regular time intervals
until some time after all the solid has dissolved for M2*

1

Plot a graph of temperature vs time

1

Extrapolate back to time of mixing = 0 and determine the temperature

1

(b) Heat taken in = $m \times c \times \Delta T = 50 \times 4.18 \times 5.4 = 1128.6 \text{ J}$

Max 2 if 14.6 °C used as ΔT

1

Moles of KCl = $5.00 / 74.6 = 0.0670$

1

Enthalpy change per mole = $+1128.6 / 0.0670 = 16\,839 \text{ J mol}^{-1}$

1

= $+16.8 \text{ (kJ mol}^{-1}\text{)}$

Answer must be given to this precision

1

(c) $\Delta H_{\text{solution}} = \Delta H_{\text{lattice}} + \Delta H(\text{hydration of calcium ions}) + 2 \times \Delta H(\text{hydration of chloride ions})$

$\Delta H_{\text{lattice}} = \Delta H_{\text{solution}} - \Delta H(\text{hydration of calcium ions}) - 2 \times \Delta H(\text{hydration of chloride ions})$

1

$$\Delta H_{\text{lattice}} = -82-9 - (-1650 + 2 \times -364) = +2295 \text{ (kJ mol}^{-1}\text{)}$$

1

(d) Magnesium ion is smaller than the calcium ion

1

Therefore, it attracts the chloride ion more strongly / stronger ionic bonding

1

[12]