



**Mark Scheme**

Q1.

Question Number	Answer	Mark	
(i)	<p>The only correct answer is <b>B</b> (elimination)</p> <p><i>A is not correct because this is a typical reaction of alkenes, not a reaction to form alkenes</i></p> <p><i>C is not correct because alcohols are typically oxidised to aldehydes, ketones or carboxylic acids</i></p> <p><i>D is not correct because substitution removes just the -OH not an -H as well</i></p>	<b>(1)</b>	
Question Number	Answer	Additional Guidance	Mark
(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• compounds with the same structural formula <b>(1)</b></li> <li>• where the atoms have a different arrangement in space <b>(1)</b></li> </ul>	<p>Allow the bonds/groups have different spatial arrangements or orientation or configuration or 3D arrangement</p> <p>Allow have a different displayed formula</p> <p>Do not award where the molecules have a different arrangement in space</p> <p>Do not award a discussion of optical isomerism</p> <p>Do not award just 'cis/trans isomerism / E/Z isomerism'</p>	<b>(2)</b>

## Edexcel Chemistry A-level - Alcohols

Question Number	Answer	Additional Guidance	Mark
(iii)	<ul style="list-style-type: none"> <li>any two of structures and/or names correct (1)</li> <li>both structures and names correct. (1)</li> </ul>	  <p>Z/cis-but-2-ene                      E/trans-but-2-ene</p> <p>Can be in either order.</p> <p>If the isomerism described in (b)(ii) is the position of the double bond allow but-1-ene and either Z/cis- or E/trans-but-2-ene here.</p> <p>Allow skeletal/displayed formulae</p>	(2)
Question Number	Answer	Additional Guidance	Mark
(iv)	<ul style="list-style-type: none"> <li>geometric (isomerism)</li> </ul>	Accept <i>cis-trans</i> / <i>E-Z</i>	(1)

Q2.

Question Number	Answer	Additional Guidance	Mark
(i)	<p>EITHER</p> <ul style="list-style-type: none"> <li>correct equation (1)</li> <li>butanal (1)</li> <li>distil (off immediately) / distillation (1)</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>correct equation (1)</li> <li>butanoic acid (1)</li> <li>heat under reflux (1)</li> </ul>	<p>The condition mark is dependent on one of the other two marks being scored</p> <p>Allow 2 marks for correct use of propan-1-ol</p> $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH} + [\text{O}] \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{CHO} + \text{H}_2\text{O}$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH} + 2[\text{O}] \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{COOH} + \text{H}_2\text{O}$ <p>Allow just 'reflux'</p> <p>Award other correct formulae for butan-1-ol, butanal and butanoic acid, e.g. <math>\text{C}_3\text{H}_7\text{CH}_2\text{OH}</math>, <math>\text{C}_3\text{H}_7\text{CHO}</math> and <math>\text{C}_3\text{H}_7\text{COOH}</math></p> <p>Do not award molecular formulae for butanal and butanoic acid</p>	(3)

## Edexcel Chemistry A-level - Alcohols

Question Number	Answer	Mark
(ii)	<p>The only correct answer is <b>B</b> (green)</p> <p><i>A is not correct because brown is not a colour which is associated with this reaction</i></p> <p><i>C is not correct because this is the colour of potassium dichromate(VI) before the reaction</i></p> <p><i>D is not correct because this is the colour of potassium chromate(VI)</i></p>	(1)

Q3.

Question Number	Acceptable Answer	Additional guidance	Mark
	<ul style="list-style-type: none"> <li>Balanced equation (1)</li> <li>Appropriate colours (1)</li> </ul>	<p><u>Example of equation:</u>  <math>\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}</math>                      Allow multiples</p> <p>Orange colourless green colourless</p> <p>Allow 'No colour' for colourless</p> <p>Do not award 'blue' for 'green'                      Do not award 'clear' for colourless                      Do not award if any spaces left blank</p> <p>Ignore any colour given for electrons                      Ignore any shades of colour</p>	(2)

Q4.

Question Number	Acceptable Answer	Additional Guidance	Mark
(i)	<p>moles of NaOH  <math>18.45 \times 0.400/1000 = 7.38 \times 10^{-3}/0.00738</math>                      (1)</p> <p>moles of propanedioic acid  <math>7.38 \times 10^{-3}/2 = 3.69 \times 10^{-3}/0.00369</math>                      (1)</p>	TE: moles of NaOH/2	(2)

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	<p>moles of propanedioic acid in 250 cm<sup>3</sup> (1)</p> <p>mass of propanedioic acid in 250 cm<sup>3</sup> (1)</p>	<p><u>example of calculation</u></p> <p>moles of propanedioic acid 25 x answer to (c)(i) = 25 x 3.69 x 10<sup>-3</sup> = 0.09225</p> <p>0.09225 x 104 = 9.6/9.59/9.594 (g)</p> <p>Allow calculation in either order e.g. calculate mass propanedioic acid in 10.0 cm<sup>3</sup> first then x 25</p> <p>Allow TE from c(i) eg 0.00738 gives 19.188 (g)</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
(iii)	<p>theoretical yield (1)</p> <p>% yield (1)</p>	<p><u>example of calculation</u></p> <p>theoretical yield 15.2 x 104/76 = 20.8 g</p> <p>% yield answer to c(ii) x 100/20.8 9.594 x 100/20.8 = 46/46.1/46.12/46.13/46.125 (%)</p> <p>use of 9.6 gives 46.15385</p> <p>allow any number of sig figs except one</p> <p>Correct answer with or without working scores 2 marks TE on incorrect theoretical yield and answer to c(ii)</p> <p>Both marks will be lost for use of 15.2 as theoretical yield (gives 63.1%)</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
(iv)	<p>an answer that makes reference to one of the following points:</p> <ul style="list-style-type: none"> <li>transfer losses</li> <li>incomplete reaction/oxidation/ formation of aldehyde</li> <li>side reaction(s)</li> </ul>	<p>Ignore spillage/impure reactants/incompetence/references to uncertainties</p> <p>Ignore other products formed/loss by evaporation</p> <p>Penalise additional incorrect reasons ie +1 -1 = zero</p>	(1)

Q5.

Question Number	Answer	Additional Guidance	Mark														
	<p>A description that makes reference to two of the following points:</p> <ul style="list-style-type: none"> <li>reagent (1)</li> <li>corresponding observation (1)</li> </ul>	<p><u>Examples of reagents and observations</u></p> <table border="1"> <thead> <tr> <th>Reagent</th> <th>Observation</th> </tr> </thead> <tbody> <tr> <td>bromine water Allow bromine (in an organic solvent)</td> <td>orange / yellow / brown solution goes colourless Allow bromine water is decolourised</td> </tr> <tr> <td>carboxylic acid and (concentrated) <math>\text{H}_2\text{SO}_4 / \text{HCl} / \text{H}^+</math></td> <td>characteristic smell (of an ester)</td> </tr> <tr> <td>acidified potassium manganate(VII) / permanganate</td> <td>purple to colourless / decolourised</td> </tr> <tr> <td>alkaline potassium manganate(VII)</td> <td>purple to green</td> </tr> <tr> <td>(neutral) potassium manganate(VII)</td> <td>purple to brown ppt</td> </tr> <tr> <td>acidified (potassium) dichromate(VI) (ions)</td> <td>orange to green</td> </tr> </tbody> </table> <p>Allow names or formulae for reagents but if both are given, both must be correct</p> <p>Ignore conditions e.g. heat</p> <p>Do not award <math>\text{PCl}_5 / \text{Na}</math></p> <p>If more than one test is given, penalise any incorrect tests</p>	Reagent	Observation	bromine water Allow bromine (in an organic solvent)	orange / yellow / brown solution goes colourless Allow bromine water is decolourised	carboxylic acid and (concentrated) $\text{H}_2\text{SO}_4 / \text{HCl} / \text{H}^+$	characteristic smell (of an ester)	acidified potassium manganate(VII) / permanganate	purple to colourless / decolourised	alkaline potassium manganate(VII)	purple to green	(neutral) potassium manganate(VII)	purple to brown ppt	acidified (potassium) dichromate(VI) (ions)	orange to green	(2)
Reagent	Observation																
bromine water Allow bromine (in an organic solvent)	orange / yellow / brown solution goes colourless Allow bromine water is decolourised																
carboxylic acid and (concentrated) $\text{H}_2\text{SO}_4 / \text{HCl} / \text{H}^+$	characteristic smell (of an ester)																
acidified potassium manganate(VII) / permanganate	purple to colourless / decolourised																
alkaline potassium manganate(VII)	purple to green																
(neutral) potassium manganate(VII)	purple to brown ppt																
acidified (potassium) dichromate(VI) (ions)	orange to green																

## Edexcel Chemistry A-level - Alcohols

Q6.


Question Number	Answer	Mark
(i)	<p>The only correct answer is B (elimination)</p> <p><i>A is not correct because the reaction involves only one reacting molecule (cyclohexanol)</i></p> <p><i>C is not correct because there is no change in the oxidation numbers of any of the elements involved</i></p> <p><i>D is not correct because nothing has been substituted.</i></p>	(1)

Question Number	Answer	Additional Guidance	Mark
(ii)	<ul style="list-style-type: none"> <li>calculation of mass and mol of cyclohexanol (1)</li> <li>calculation of mol of cyclohexene (1)</li> <li>calculation of mass of cyclohexene (1)</li> <li>calculation of volume of cyclohexene (1)</li> </ul>	<p><u>Example of calculation</u></p> <p>Mass cyclohexanol = <math>10.0 \times 0.962 = 9.62</math> (g) and Mol of cyclohexanol = <math>9.62 \div 100 = 0.0962</math> (mol)</p> <p>Mol of cyclohexene = <math>0.63 \times 0.0962 = 0.060606</math> (mol)</p> <p>Mass of cyclohexene = <math>0.060606 \times 82.0 = 4.9697</math> (g)</p> <p>Volume of cyclohexene = <math>4.9697 \div 0.811 = 6.1279</math> = <math>6 / 6.1 / 6.13</math> (cm<sup>3</sup>) / <math>6.1 \times 10^{-3}</math> dm<sup>3</sup></p> <p>Ignore SF in final answer Allow TE at each stage Correct answer with no working scores 4</p>	(4)

## Edexcel Chemistry A-level - Alcohols

Question Number	Answer	Additional Guidance	Mark																				
* (iii)	<p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table border="1"> <thead> <tr> <th></th> <th>Number of marks awarded for structure and sustained lines of reasoning</th> </tr> </thead> <tbody> <tr> <td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.</td> <td>2</td> </tr> <tr> <td>Answer is partially structured with some linkages and lines of reasoning.</td> <td>1</td> </tr> <tr> <td>Answer has no linkages between points and is unstructured.</td> <td>0</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure and sustained lines of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2	Answer is partially structured with some linkages and lines of reasoning.	1	Answer has no linkages between points and is unstructured.	0	<p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning, scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> <p>In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.</p> <p>If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded do not deduct mark(s).</p> <p>Comment: Look for the indicative marking points first, then consider the mark for the structure of the answer and sustained line of reasoning.</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																						
6	4																						
5-4	3																						
3-2	2																						
1	1																						
0	0																						
	Number of marks awarded for structure and sustained lines of reasoning																						
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2																						
Answer is partially structured with some linkages and lines of reasoning.	1																						
Answer has no linkages between points and is unstructured.	0																						
	<p><b>Indicative content:</b></p> <ul style="list-style-type: none"> <li>IP1 separate cyclohexene from water using separating funnel</li> <li>IP2 remove the lower aqueous layer because it has a higher density</li> <li>IP3 suitable drying agent</li> <li>IP4 separate the cyclohexene from the drying agent</li> </ul>	<p>Mark independently</p> <p>Suitable specified drying agents: (anhydrous) CaCl<sub>2</sub>, Na<sub>2</sub>SO<sub>4</sub>, CaSO<sub>4</sub>, MgSO<sub>4</sub></p>																					
	<ul style="list-style-type: none"> <li>IP5 redistil product</li> <li>IP6 collect the distillate boiling over between 80 and 86 °C to collect the pure cyclohexene</li> </ul>	By filtering or decanting																					

Q7.


Question Number	Answer	Additional Guidance	Mark
(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>name (1)</li> <li>displayed formula (1)</li> </ul>	<p><u>Example of displayed formula</u></p>  <p>Allow CH<sub>2</sub> groups Allow skeletal formula Do not award molecular formula</p>	(2)

Question Number	Answer	Additional Guidance	Mark
(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>O-H bond (stretching) 3750 – 3200 cm<sup>-1</sup> in cyclohexanol is not present in cyclohexanone /disappears (when cyclohexanol reacts). (1)</li> <li>C=O bond (stretching) 1720 – 1700 cm<sup>-1</sup> appears in cyclohexanone (1)</li> </ul>	<p>Allow a range within the specified range</p> <p>Allow 1725 – 1700 cm<sup>-1</sup> Do not allow 1740 – 1720 cm<sup>-1</sup> (aldehyde)</p>	(2)

Question Number	Answer	Additional Guidance	Mark
(iii)	<ul style="list-style-type: none"> <li>highest <math>m/z = M_r = 98</math></li> </ul>	<p>Check, answer may be shown on mass spectrum Do not accept just '98' with no supporting evidence</p> <p>Allow peak furthest to the right / molecular ion peak is 98</p>	(1)



## Edexcel Chemistry A-level - Alcohols


Question Number	Answer	Additional Guidance	Mark
(iv)	<ul style="list-style-type: none"> <li>fragment (1)</li> <li>charge (1)</li> </ul>	<p><u>Examples of fragment structure</u></p>  <p>Allow charge anywhere on fragment, including outside brackets around the fragment</p> <p>Allow straight chain fragment provided it has the correct number of C and H atoms</p>	(2)

Q8.

Question Number	Acceptable Answer	Additional guidance	Mark
	<p>An explanation that makes reference to the following</p> <ul style="list-style-type: none"> <li>ethanol would be oxidised to ethanal (1)</li> <li>because ethanal has a low boiling temperature or ethanal will distil before ethanoic acid can be formed (1)</li> </ul>	<p>Allow aldehyde for ethanal</p> <p>Allow ethanal will be formed</p> <p>Allow ethanal is (more) volatile</p> <p>Accept reverse argument in terms of reflux condensing ethanal for oxidation to ethanoic acid</p>	(2)

## Edexcel Chemistry A-level - Alcohols

Q9.

Question Number	Answer	Additional Guidance	Mark
	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>a separating funnel (with or without a stopper or bung) <b>(1)</b></li> <li>a two layer system with the top layer labelled as the organic layer / 2-chloro-2- methylpropane <b>(1)</b></li> </ul>	<p>Allow any shape separating funnel with a tap at the bottom (no label required) with a bung, stopper or appropriate joint / gap at the top. Allow anything labelled as a tap</p> <p>Do not award a sealed apparatus if stopper / bung is unclear</p>  <p>Do not award a three layer system</p>	<b>(2)</b>

Q10.

Question Number	Answer	Additional Guidance	Mark
(i)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>carbon dioxide / CO<sub>2</sub></li> </ul>	Ignore references to limewater turning cloudy	<b>(1)</b>

Question Number	Answer	Additional Guidance	Mark
(ii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>H<sup>+</sup> / H<sub>3</sub>O<sup>+</sup></li> </ul>	Ignore 'hydrogen ion' Ignore numbers before e.g. 2H <sup>+</sup>	<b>(1)</b>

## Edexcel Chemistry A-level - Alcohols

Question Number	Answer	Additional Guidance	Mark
(iii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• mixed with an appropriate named drying agent, e.g. (anhydrous) calcium chloride / <math>\text{CaCl}_2</math> / (anhydrous) magnesium sulfate / <math>\text{MgSO}_4</math> / (anhydrous) sodium sulfate / <math>\text{Na}_2\text{SO}_4</math> / silica gel</li> <li>• leave until the solution becomes clear / left until added drying agent remains powdered / left until added drying agent does not clump together</li> </ul> <p><b>or</b></p> <p>decant the liquid / filter the solid (to separate from the drying agent)</p>	<p>M2 is dependent on a drying agent being added in M1</p> <p>Do not award sodium hydroxide, potassium hydroxide, anhydrous copper sulfate, anhydrous cobalt chloride, calcium sulfate, calcium carbonate, potassium sulfate</p>	(2)

Q11.

Question Number	Answer	Additional Guidance	Mark
(i)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>• 50-52 (°C)</li> </ul>	<p>Allow 48-54 (°C) Allow a range within these limits to include 51 (°C) Do not award just 51 (°C)</p>	(1)

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answer	Additional Guidance	Mark												
* (ii)	<p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="389 786 836 1106"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0	<p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning.</p> <p>For example, an answer with five indicative marking points, which is partially structured with some linkages and lines of reasoning, scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points														
6	4														
5-4	3														
3-2	2														
1	1														
0	0														

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answer	Additional Guidance	Mark								
* (ii) contd	<p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table border="1" data-bbox="355 344 1018 875"> <thead> <tr> <th data-bbox="355 344 724 488"></th> <th data-bbox="724 344 1018 488">Number of marks awarded for structure of answer and sustained line of reasoning</th> </tr> </thead> <tbody> <tr> <td data-bbox="355 488 724 663">Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.</td> <td data-bbox="724 488 1018 663">2</td> </tr> <tr> <td data-bbox="355 663 724 770">Answer is partially structured with some linkages and lines of reasoning.</td> <td data-bbox="724 663 1018 770">1</td> </tr> <tr> <td data-bbox="355 770 724 875">Answer has no linkages between points and is unstructured.</td> <td data-bbox="724 770 1018 875">0</td> </tr> </tbody> </table>		Number of marks awarded for structure of answer and sustained line of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2	Answer is partially structured with some linkages and lines of reasoning.	1	Answer has no linkages between points and is unstructured.	0	<p>In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.</p> <p>Reasoning marks may be reduced for extra incorrect chemistry</p> <p>Ignore stated errors which are not present</p> <p>Allow to prevent uneven boiling / ensure smooth boiling Ignore prevents bumping Do not award so reaction does not explode / shatter glassware / damage apparatus</p>	(6)
	Number of marks awarded for structure of answer and sustained line of reasoning										
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2										
Answer is partially structured with some linkages and lines of reasoning.	1										
Answer has no linkages between points and is unstructured.	0										

## Edexcel Chemistry A-level - Alcohols

	<p><b>Indicative content:</b></p> <ul style="list-style-type: none"> <li>• IP1 add anti-bumping granules</li> <li>• IP2 to prevent the formation of large bubbles / rapid heating / transfer of reaction mixture to collecting vessel (leading to impure product)</li> <li>• IP3 the thermometer should be opposite the entrance of the condenser</li> <li>• IP4 collecting over the wrong temperature range (therefore impure or the wrong product)</li> <li>• IP5 add more ice-water mixture</li> <li>• IP6 ensure you collect as much product as possible</li> </ul>	<p>Allow thermometer should be measuring the vapour temperature not the liquid temperature</p> <p>Allow collecting impure product but must be linked to wrong position of thermometer Do not award just the temperature is inaccurate without mention of vapour</p> <p>Allow collection flask should be further in the ice-water mixture</p> <p>Allow to ensure greater / quicker condensation</p>	
--	---	--	--

Q12.

Question Number	Acceptable Answer	Additional Guidance	Mark
(a)	$  \begin{array}{c}  \text{H} \\    \\  \text{H}-\text{C}-\text{H} \\    \\  \text{H}-\text{C}-\text{C}-\text{O}-\text{H} \\    \quad   \\  \text{H} \quad \text{H}-\text{C}-\text{H} \\    \\  \text{H}  \end{array}  $	<p>display all three methyl groups allow -OH do not award C-H-O</p>	(1)

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answer	Additional Guidance	Mark
(b)(i)	An answer that makes reference to one of the following:  molecular ion/molecule fragments/is unstable		(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	$\begin{array}{c} \text{CH}_3-\overset{+}{\text{C}}-\text{CH}_3 \\   \\ \text{O}-\text{H} \end{array}$	allow + charge on any part of the ion/outside the structure but + must be shown  allow displayed/structural/skeletal/molecular formulae or any combination of these.	(1)


Question Number	Acceptable Answer	Additional Guidance	Mark
(c)(i)	<ul style="list-style-type: none"> <li>calculation for bonds broken in the alcohol (*) <b>(1)</b></li> <li>calculation for bonds broken in oxygen <b>and</b> total energy for bonds broken(**) <b>(1)</b></li> <li>calculation for bonds made(***) <b>(1)</b></li> <li>calculation of <math>\Delta_c H</math> (2-methylpropan-2-ol) with sign <b>(1)</b></li> </ul>	<p><u>Example of calculation</u></p> $3(\text{C}-\text{C}) + 9(\text{C}-\text{H}) + (\text{C}-\text{O}) + (\text{O}-\text{H})$ $= (3 \times 347) + (9 \times 413) + 358 + 464 = (+)5580 \text{ (kJ mol}^{-1}\text{)}$ $6(\text{O}=\text{O}) = (6 \times 498) = (+)2988 \text{ (kJ mol}^{-1}\text{)}$ <p>total = + 5580 + 2988 = (+)8568 (kJ mol<sup>-1</sup>) TE from ans * M1 + 2988</p> $= 8(\text{C}=\text{O}) + 10(\text{O}-\text{H})$ $= (8 \times 805) + (10 \times 464) = -11080 \text{ (kJ mol}^{-1}\text{)}$ $= +8568 - 11080 = -2512 \text{ (kJ mol}^{-1}\text{)}$ <p>allow TE for answer(**) + answer(***) units not required but if given they must be correct correct final answer with no working scores 4 marks</p>	(4)

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>incomplete combustion (1)</li> <li><math>\Delta_c H</math> (2-methylpropan-2-ol) will be less negative /less exothermic than data book value (1)</li> </ul>	<p>mark independently</p> <p>do not award just lower/smaller/decreases/ more positive allow reduce the magnitude (of the value)</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
(iii)	<p>An answer that makes reference to the following points:</p> <p><math>\Delta_c H</math> figures are at 298 K /data book bond energies refer to gaseous state <u>and</u> water and/or 2-methylpropan-2-ol are/is (both) liquid(s) (at 298 K)</p>	<p>allow just liquid involved</p> <p>do not award data book bond energies are mean (values)/not specific to 2-methylpropan-2-ol</p>	(1)

Q13.

Question Number	Answer	Additional Guidance	Mark
	<ul style="list-style-type: none"> <li>skeletal formula</li> </ul>	<p>Example of skeletal formula</p>  <p>Bond from ring to Br must be shown but the length is not important.</p>	(1)




Edexcel Chemistry A-level - Alcohols


Q14.

Question Number	Acceptable Answers	Additional Guidance	Mark
(a)	<ul style="list-style-type: none"> <li>calculation of empirical formula (1)</li> <li>uses molecular ion to prove molecular formula (1)</li> </ul> <p>or</p> <ul style="list-style-type: none"> <li>calculation of percentage of each element in compound all 3 correct scores (2) any 2 correct scores (1)</li> </ul> <p>or</p> <ul style="list-style-type: none"> <li>calculation of the number of atoms of each element directly all 3 correct scores (2) any 2 correct scores (1)</li> </ul>	<p>Example of calculation</p> $\begin{array}{r} \text{C} : \text{H} : \text{O} \\ \hline 68.2 \quad 13.6 \quad 18.2 \\ 12 \quad 1 \quad 16 \\ = \quad 5.68 \quad 13.6 \quad 1.14 \\ = \quad 5 \quad 12 \quad 1 \end{array}$ <p>Use of 88 to show molecular formula is <math>\text{C}_5\text{H}_{12}\text{O}</math> e.g. <math>M_r</math> is <math>(5 \times 12) + (12 \times 1) + 16 = 88</math> or states that <math>M_r</math> of empirical formula is 88</p> <p>or</p> <p>% C = <math>\frac{5 \times 12 \times 100}{88} = 68.2</math> % H = <math>\frac{12 \times 1 \times 100}{88} = 13.6</math> % O = <math>\frac{1 \times 16 \times 100}{88} = 18.2</math></p> <p>or</p> <p>C atoms = <math>\frac{68.2 \times 88}{100 \times 12} = 5</math> H atoms = <math>\frac{13.6 \times 88}{100 \times 1} = 12</math> O atoms = <math>\frac{18.2 \times 88}{100 \times 16} = 1</math></p>	(2)
Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(i)	<ul style="list-style-type: none"> <li>(X is a) primary/ 1° (alcohol)</li> </ul>		(1)


# Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(ii)		<p>Allow alcohols in any order</p> <p>Allow CH<sub>3</sub> / OH</p> <p>Allow slip of 1 H missing from 1 alcohol / 1 C-C bond missing</p> <p>Ignore names, even if incorrect</p> <p>Penalise O-H-C- / -C-H-O at end of molecule once only</p> <p>If no other mark is given, allow (2) for 4 correct skeletal / structural formulae or any combination of these or (1) for 3 correct</p> <p>Allow (2) for displayed formulae of pentan-2-ol, pentan-3-ol and 3-methylbutan-2-ol if secondary alcohol in (b)(i), or (1) for any two of those</p>	(3)

<ul style="list-style-type: none"> <li>• 4 correct</li> <li>• 3 correct</li> <li>• 2 correct</li> </ul>	<p>(3)</p> <p>(2)</p> <p>(1)</p>	<p>If no other mark awarded and if (b)(i) is blank or incorrect, allow (2) for any 4 different alcohols with formula C<sub>5</sub>H<sub>12</sub>O, (1) for 3 alcohols</p>	
---	----------------------------------	---	--

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(iii)	<ul style="list-style-type: none"> <li>• </li> </ul>	<p>Allow structural formula or any combination of displayed and structural formula</p> <p>Allow + anywhere on structure or outside of a formula in a bracket</p> <p>Do not allow C<sub>2</sub>H<sub>5</sub>O<sup>+</sup>/C<sub>2</sub>H<sub>4</sub>OH<sup>+</sup></p> <p>Do not allow missing charge</p> <p>Allow CH<sub>3</sub>C<sup>+</sup>HOH if secondary alcohol identified in (b)(i)</p>	(1)

Edexcel Chemistry A-level - Alcohols



Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(iv)	<ul style="list-style-type: none"> <li>•           <div style="text-align: center;">  </div> </li> </ul> <p style="text-align: right;"><b>(1)</b></p> <ul style="list-style-type: none"> <li>• because this is the <b>only</b> alcohol with a branched chain <u>and</u> forms <math>\text{CH}_2\text{OHCH}_2^+</math> / <math>\text{C}_2\text{H}_4\text{OH}^+</math> / peak at 45 / fragment identified in (b)(iii)</li> </ul> <p style="text-align: right;"><b>(1)</b></p>	<p>Allow any type of identification, including name 3-methylbutan-1-ol</p> <p>Ignore incorrect name with correct structure</p> <p>Conditional on correct identification Ignore missing charge on fragment</p> <p>Allow reasons why the others are not correct e.g. not pentan-1-ol as it is not branched <u>and</u> not 2-methylbutan-1-ol or 2,2-dimethylpropan-1-ol as they do not form <math>\text{CH}_2\text{OHCH}_2^+</math></p> <p>If secondary alcohol identified in (b)(i): Allow 3-methylbutan-2-ol (1) as it is the only alcohol with a branched chain that forms <math>\text{CH}_3\text{C}^+\text{HOH}</math> (1)</p>	<b>(2)</b>

# Edexcel Chemistry A-level - Alcohols

Q15.

Question Number	Answer	Additional Guidance	Mark																				
*	<p>This question assesses the student's ability to show a coherent and logically structured answer with linkages and fully sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning</p> <table border="1"> <thead> <tr> <th></th> <th>Number of marks awarded for structure of answer and sustained lines of reasoning</th> </tr> </thead> <tbody> <tr> <td>Answer shows a coherent logical structure with linkages and fully sustained lines of reasoning demonstrated throughout</td> <td>2</td> </tr> <tr> <td>Answer is partially structured with some linkages and lines of reasoning</td> <td>1</td> </tr> <tr> <td>Answer has no linkages between points and is unstructured</td> <td>0</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure of answer and sustained lines of reasoning	Answer shows a coherent logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2	Answer is partially structured with some linkages and lines of reasoning	1	Answer has no linkages between points and is unstructured	0	<p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, a response with four indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there were no linkages between the points, then the same indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and zero marks for linkages).</p> <p><b>Typically</b></p> <p><b>Number of IPs Reasoning mark</b></p> <p>6 or 5 scores 2</p> <p>4 or 3 scores 1</p> <p>2 or 1 or 0 scores 0</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																						
6	4																						
5-4	3																						
3-2	2																						
1	1																						
0	0																						
	Number of marks awarded for structure of answer and sustained lines of reasoning																						
Answer shows a coherent logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2																						
Answer is partially structured with some linkages and lines of reasoning	1																						
Answer has no linkages between points and is unstructured	0																						

## Edexcel Chemistry A-level - Alcohols

	<p><b>Indicative content</b></p> <ul style="list-style-type: none"> <li>• <b>IP1</b> any mention of <b>oxidation</b> of ethanol or <b>oxidation</b> of flavan-3-ol (by oxygen in the air)</li> <li>• <b>IP2</b> for formation of <b>either</b> ethanoic acid or ethanal (from ethanol)</li> <li>• <b>IP3</b> for formation of ethyl ethanoate (from the reaction between ethanol and ethanoic acid)</li> <li>• <b>IP4</b> for structure / name of flavan-3-one</li> <li>• <b>IP5</b> for (-OH group on) flavan-3-ol forms an ester with ethanoic acid</li> <li>• <b>IP6</b> correct structure of the ester formed between flavan-3-ol and ethanoic acid</li> </ul>  <p>This is the structure of the ester formed between flavan-3-ol and ethanoic acid</p>	<p>Allow names or formulae <b>but</b> if both are given <b>both</b> must be correct</p>  <p><b>Comment</b> For correct structure of the ester formed between flavan-3-ol and ethanoic acid award both <b>IP5</b> and <b>IP6</b></p> <p>Do not award <b>IP4</b> if the product is described as an aldehyde</p>	
--	---	---	--


### Q16.

Question Number	Acceptable Answers	Additional Guidance	Mark
(i)	$\text{C}_5\text{H}_{12}\text{O} + [\text{O}] \rightarrow \text{C}_5\text{H}_{10}\text{O} + \text{H}_2\text{O}$ <ul style="list-style-type: none"> <li>• left-hand side of equation correct <b>(1)</b></li> <li>• right-hand side of equation correct <b>(1)</b></li> </ul>	<p><b>Molecular</b> formulae must be used throughout</p> <p>Allow [O] above the arrow</p> <p>Do not award for <math>\text{C}_5\text{H}_{11}\text{OH}</math> as the alcohol</p> <p>Ignore state symbols if incorrect or conditions mentioned</p>	<b>(2)</b>

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(ii)	<ul style="list-style-type: none"> <li>calculation of moles of both of C<sub>5</sub>H<sub>10</sub>O and C<sub>5</sub>H<sub>12</sub>O (1)</li> <li>calculation of mass of C<sub>5</sub>H<sub>12</sub>O (1)</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>calculation of theoretical mass of C<sub>5</sub>H<sub>10</sub>O <b>and</b> moles of C<sub>5</sub>H<sub>10</sub>O (1)</li> <li>calculation of mass of C<sub>5</sub>H<sub>12</sub>O (1)</li> </ul>	<p>Example of calculation</p> <p>Moles C<sub>5</sub>H<sub>10</sub>O = <math>\frac{2.15}{86.0} = 0.025(0)</math> (mol)</p> <p><b>and</b></p> <p>moles C<sub>5</sub>H<sub>12</sub>O = <math>\frac{0.025(0)}{62.5} \times 100 = 0.04(00)</math></p> <p>(So) mass of C<sub>5</sub>H<sub>12</sub>O = 0.04(00) x 88 = 3.52 g</p> <p>Theoretical mass C<sub>5</sub>H<sub>10</sub>O = <math>\frac{2.15}{62.5} \times 100 = 3.44</math> g</p> <p><b>and</b></p> <p>moles C<sub>5</sub>H<sub>10</sub>O = <math>\frac{3.44}{86.0} = 0.04(00) = \text{mol C}_5\text{H}_{12}\text{O}</math></p> <p>(So) mass of C<sub>5</sub>H<sub>12</sub>O = 0.04(00) x 88 = 3.52 g</p> <p>Correct answer with no working scores (2)</p> <p>Allow TE from MP1</p> <p>Award 1 mark for 3.36 g, 1.375 g or 2.2 g</p>	(2)

Q17.

Question Number	Acceptable Answers	Additional Guidance	Mark
(a)	<ul style="list-style-type: none"> <li>diagram of separating funnel (1)</li> <li>aqueous and organic layers labelled as shown (1)</li> </ul>	<p>Mark independently</p> <p>Allow any shape separating funnel with tap at the bottom (does not need to be labelled), with a narrowing top or vertical sides but do not allow a burette</p> <p>Allow stopper/bung in separating funnel</p>  <p>Allow two layers shown and just one labelled correctly</p> <p>Allow organic layer/ product for top layer / hydrochloric acid for aqueous layer</p> <p>Do not allow 'reactant' for top layer</p>	(2)

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)	<ul style="list-style-type: none"> <li>to react with/ neutralise any (unreacted/ excess hydrochloric) acid (1)</li> <li>to release the carbon dioxide produced or to relieve the build-up of pressure (1)</li> </ul>	Mark independently Allow to remove the (hydrochloric) acid Allow to neutralise the organic layer/ solution Allow to release gases Ignore just 'pressure builds up' Do not allow incorrect gases e.g. hydrogen	(2)

Question Number	Answer	Mark
(c)	D (sodium sulfate)	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
(d)(i)	A description that makes reference to the following points: <ul style="list-style-type: none"> <li>the (bulb of the) thermometer should be opposite the opening to the condenser (1)</li> <li>the water in and out of the condenser should be reversed (1)</li> <li>put a vent after the condenser or leave a gap between the condenser and the receiver or conical flask must be open (1)</li> </ul>	Allow these changes if shown on the diagram Allow thermometer should be higher up / above the liquid / should measure the temperature of the vapour / out of the mixture/liquid Allow water should enter the bottom (of the condenser) Ignore just 'vent' / the apparatus should not be completely sealed Ignore references to using a fume cupboard	(3)

Question Number	Acceptable Answers	Additional Guidance	Mark
(d)(ii)	<ul style="list-style-type: none"> <li>50-52°C</li> </ul>	Allow any range between 49 and 53°C, <u>provided</u> it includes 51°C Do not allow just 51°C	(1)

# Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(e)	<ul style="list-style-type: none"> <li>calculation of moles of alcohol used (1)</li> <li>calculation of theoretical volume of 2-chloro-2-methylpropane made or calculation of actual moles of 2-chloro-2-methylpropane or calculation of actual mass of 2-chloro-2-methylpropane (1)</li> <li>calculation of percentage yield (1)</li> </ul>	<p><u>Example of calculation</u>  mass of alcohol used = <math>15.0 \times 0.79 = 11.85</math> (g)  moles of alcohol used = <math>11.85/74.0 = 0.16014</math></p> <p>theoretical mass of chloro compound = <math>0.16014 \times 92.5 = 14.8125</math> (g)  theoretical volume = <math>14.8125/0.84 = 17.634</math> (cm<sup>3</sup>)  or  actual moles of chloro compound = <math>6.9 \times 0.84 / 92.5 = 0.062659</math>  or  actual mass of chloro compound = <math>0.062659 \times 92.5 = 5.796</math> (g)</p> <p>% yield = <math>(6.9/17.634) \times 100 = 39.1\%</math>  or  = <math>(0.062659/0.16014) \times 100 = 39.1\%</math>  or  = <math>(5.796/14.8125) \times 100 = 39.1\%</math></p> <p>TE on M1 and M2  Ignore SF except 1 SF</p>	(3)

		Correct answer without working scores 3	
--	--	---	--

Question Number	Acceptable Answers	Additional Guidance	Mark
(f)	<ul style="list-style-type: none"> <li>curly arrow from C-O bond to O (1)</li> <li>curly arrow from lone pair on Cl<sup>-</sup> to C<sup>+</sup> (1)</li> </ul>	<p>Do not allow single-headed arrows  Do not allow additional, incorrect arrows</p>	(2)



## Edexcel Chemistry A-level - Alcohols

Q18.

Question Number	Acceptable Answers	Additional Guidance	Mark
<b>(a)(i)</b>	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• cool the mixture because the reaction (with concentrated sulfuric acid / <math>\text{H}_2\text{SO}_4</math>) is (very) exothermic / releases (a lot of) heat <b>(1)</b></li>   <li>• anti-bumping granules are added to prevent violent / sudden / localised boiling  <b>or</b>                      to prevent superheating / large bubbles forming  <b>or</b>                      to promote smooth / even / controlled boiling  <b>or</b>                      to promote the formation of small bubbles  <b>or</b>                      to provide nucleation centres / (rough) surface for bubble formation <b>(1)</b></li>   <li>• heating under reflux is used to prevent the loss of any volatile substances / volatile reactants / volatile products / organic compound / named organic compound  <b>or</b>                      to make sure the vapour / gas condenses  <b>or</b>                      to prevent vapour escaping <b>(1)</b></li> </ul>	<p>Ignore reaction is violent / to prevent splashing / to slow down the reaction / to stop reactants evaporating</p> <p>Do not award to quench the reaction / reference to explosion</p> <p>Ignore to stop bumping / spitting / explosion / liquid splashing out / vigorous reaction / loss of reactants / to distribute heat more evenly / any reference to rate / to promote smooth heating</p> <p>Allow so that the reaction goes to completion                      Ignore just 'to prevent gas escaping' / just 'to prevent loss of reactants / products' / just 'reactants / products are volatile' / 'because 1-bromobutane / butan-1-ol is flammable' / to increase yield / reference to safety</p> <p>Do not award for reference to oxidation or reduction</p>	<b>(3)</b>

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(a)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>there is a gap between the condenser and the flask / seal the joint between the flask and condenser</li> </ul> <p><b>and</b> so vapour / gas / reactants / products will escape (1)</p> <ul style="list-style-type: none"> <li>the water is flowing the wrong way through the condenser / the water should go in at the bottom (and out at the top)</li> </ul> <p><b>and</b> so it doesn't fill with water / is only part filled / there is an airlock (1)</p> <ul style="list-style-type: none"> <li>there is a stopper on the condenser / there should not be a stopper on the condenser</li> </ul> <p><b>and</b> so there will be a build-up of pressure (if the gap between condenser and flask is closed) (1)  </p>	<p>Allow answers shown on annotated diagram e.g. gap circled Ignore any additional errors Ignore additional suggested modifications even if incorrect</p> <p>Do not award just 'the apparatus is not sealed' unless it is clear it means between the condenser and flask</p> <p>Allow so this will lower the yield of product / 1-bromobutane</p> <p>Ignore condenser is fitted the wrong way up</p> <p>Allow so there will be inefficient condensation / cooling</p> <p>Allow so the stopper will blow off / there will be an explosion / it will be dangerous</p>	(3)

Question Number	Acceptable Answers	Additional Guidance	Mark
(a)(iii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(brown vapour / it) is bromine / Br<sub>2</sub> (1)</li> <li>bromide ions / Br<sup>-</sup> / HBr oxidised (by concentrated sulfuric acid) (1)</li> </ul>	<p>Stand alone mark Do not award just 'Br' Do not award any other brown gas in addition to bromine</p> <p>Allow bromide ions / Br<sup>-</sup> / HBr reduce sulfuric acid / act as a reducing agent Ignore sodium bromide / NaBr is oxidised Ignore just 'redox reaction' Do not award bromine is oxidised Do not award oxidation by anything other than sulfuric acid</p>	(2)

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(i)	An answer that makes reference to the following: <ul style="list-style-type: none"> <li>aqueous layer is on the top  <b>and</b>  because water / it has a lower density than 1-bromobutane</li> </ul>	Allow 'it' for aqueous layer Allow because 1-bromobutane has a higher density than water  Ignore reference to butanol unless in a third layer  Do not award water is 'lighter' Do not award reference to three layers	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(ii)	An answer that makes reference to the following points:  Step 8 <ul style="list-style-type: none"> <li>(aqueous sodium hydrogencarbonate / <math>\text{NaHCO}_3</math>) reacts with / neutralises / removes the (hydrochloric) acid / <math>\text{H}^+</math> (ions) in the mixture (1)</li> </ul> Step 9 <ul style="list-style-type: none"> <li>(the tap is opened) to release the carbon dioxide / gas (formed)</li> </ul> <b>or</b> to allow the carbon dioxide / gas to escape <b>or</b> to prevent the build-up of pressure (1)  Step 10 <ul style="list-style-type: none"> <li>(anhydrous sodium sulfate is added) to remove / absorb water (1)</li> </ul>	Do not award reacts with incorrect acid e.g. $\text{H}_2\text{SO}_4$ / $\text{HBr}$ / ethanoic acid Ignore removes water  Do not award an incorrect gas e.g. hydrogen  Allow (anhydrous) sodium sulfate is a drying agent / added to dry the product  Do not award dehydration / reacts with water	(3)

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(iii)	<ul style="list-style-type: none"> <li>starting temperature 99 or 100 or 101 (<math>^{\circ}\text{C}</math>)  <b>and</b>  final temperature 103 or 104 or 105 (<math>^{\circ}\text{C}</math>)</li> </ul>	Do not award just one value / 102 ( $^{\circ}\text{C}$ ) Do not award 102 ( $^{\circ}\text{C}$ ) with another temperature	(1)

## Edexcel Chemistry A-level - Alcohols


Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(iv)	<ul style="list-style-type: none"> <li>calculation of mol of 1- bromobutane <b>(1)</b></li> <li>calculation of number of molecules <b>and</b> answer to 2/3 SF <b>(1)</b></li> </ul>	<p><u>Example of calculation</u></p> <p>mol of 1-bromobutane = <math>\frac{12.0 \times 1.27}{136.9}</math></p> <p>= 0.11132 (mol)</p> <p>Do not award 0.1</p> <p>number of molecules = <math>0.11132 \times 6.02 \times 10^{23}</math></p> <p>= <math>6.7 \times 10^{22} / 6.70 \times 10^{22}</math></p> <p>TE on a calculated mol 1-bromobutane using <math>M_r</math></p> <p>Correct answer to 2 or 3 SF with no working scores (2)</p>	<b>(2)</b>

Q19.

Question Number	Answer	Additional Guidance	Mark
	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li><math>\text{PCl}_5</math> / phosphorus(V) chloride / phosphorus pentachloride</li> </ul>	<p>Allow thionyl chloride / <math>\text{SOCl}_2</math></p> <p>Allow phosphorus(III) chloride / <math>\text{PCl}_3</math> / phosphorus trichloride</p> <p>Ignore phosphorus chloride</p> <p>If name and formula are given both must be correct</p>	<b>(1)</b>

# Edexcel Chemistry A-level - Alcohols

Q20.

Question Number	Acceptable Answer	Additional guidance	Mark
(i)	A diagram with any shading that is not 100%	<p>An example of a suitable diagram:</p>  <p>Allow shaded area to show 'air pockets'</p>	(1)

Question Number	Acceptable Answer	Additional guidance	Mark
(ii)	<p>An answer that makes reference to the following</p> <p>prevention of uncontrolled boiling by:</p> <ul style="list-style-type: none"> <li>distributing the heat more evenly</li> </ul> <p>or</p> <ul style="list-style-type: none"> <li>providing a surface for bubbles to form/allow smaller bubbles to form/provides nucleation sites for bubbles</li> </ul>	<p>Do not award provide surface area for reaction</p> <p>Ignore reference to mixing the reagents/provide smooth boiling</p>	(1)

Edexcel Chemistry A-level - Alcohols

Q21.

Question Number	Acceptable Answers	Additional Guidance	Mark
	<p><b>Identifications</b></p> <ul style="list-style-type: none"> <li>structure of alcohol <b>B</b> (1)</li> <li>structure of ester <b>A</b> (1)</li> </ul> <p><b>Justification</b></p> <ul style="list-style-type: none"> <li>butan-2-ol / <math>\text{CH}_3\text{CH}_2\text{CHOHCH}_3</math> is the <b>only</b> alcohol (with formula <math>\text{C}_4\text{H}_{10}\text{O}</math>) that (undergoes elimination and) produces (but-1-ene and) but-2-ene (1)</li> </ul>	<p>Allow any combination of structural, displayed or skeletal formulae / correct species in unbalanced equations</p> <p>Allow structures not labelled <b>A</b> and <b>B</b></p> <p>Penalise missing H once only <u>Examples of identification</u></p> <p style="text-align: right;"><b>(B)</b></p> $\begin{array}{cccc} \text{H} & \text{H} & \text{OH} & \text{H} \\   &   &   &   \\ \text{H}-\text{C} & -\text{C} & -\text{C} & -\text{C}-\text{H} \\   &   &   &   \\ \text{H} & \text{H} & \text{H} & \text{H} \end{array}$ <p>Ignore connectivity of the OH group unless horizontal</p> <p style="text-align: right;"><b>(A)</b></p> $\begin{array}{ccccccc} & & & \text{H} & & & \\ & & &   & & & \\ & & & \text{H}-\text{C}-\text{H} & & & \\ & & &   & & & \\ \text{H} & \text{H} & \text{O} & & \text{H} & \text{H} & \text{H} \\   &    &   & &   &   &   \\ \text{H}-\text{C} & -\text{C} & -\text{O} & -\text{C} & -\text{C} & -\text{C}-\text{H} \\   & & &   &   &   & \\ \text{H} & & & \text{H} & \text{H} & \text{H} & \end{array}$ <p>Ignore incorrect name for <b>A</b></p> <p>TE on incorrect alcohol</p> <p>Allow butan-2-ol can form a double bond either side of the C with OH / between the 1<sup>st</sup> and 2<sup>nd</sup> carbon atoms and the 2<sup>nd</sup> and 3<sup>rd</sup> carbon atoms – this can be shown on diagram / equation</p> <p>Allow OH must be on the 2<sup>nd</sup> carbon atom / secondary alcohol to form but-1-ene and but-2-ene</p> <p>Allow butan-1-ol gives but-1-ene <b>and</b> 2-methylpropan-1-ol / 2-methylpropan-2-ol gives (2-)methylpropene</p> <p>Allow the other alcohols (with formula <math>\text{C}_4\text{H}_{10}\text{O}</math>) do not give but-2-ene</p>	<p><b>(3)</b></p>

## Edexcel Chemistry A-level - Alcohols

Q22.

Question Number	Answer	Additional Guidance	Mark
(i)	<p>An explanation that makes reference to</p> <ul style="list-style-type: none"> <li>propanal is condensed back (to the pear-shaped flask) (1)</li> <li>so propanal is (further) oxidised (to propanoic acid) or propanal is more readily oxidised than propan-1-ol  (1)</li> </ul>	<p>Allow aldehyde for propanal</p> <p>Allow 'apparatus is reflux' Allow propanal is not being removed /distilled off (from the oxidising agent)</p> <p>Ignore just 'reacts further'</p> <p>Do not award reference to propanal being completely oxidised</p>	(2)

Question Number	Answer	Additional Guidance	Mark
(ii)	<ul style="list-style-type: none"> <li>(+VI)</li> </ul>	Allow (+) six / (+)6 / six (+) / 6(+)	(1)

Question Number	Answer	Additional Guidance	Mark
(iii)	<ul style="list-style-type: none"> <li>balanced equation</li> </ul>	<p>Example of equation</p> $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} \rightarrow \text{CH}_3\text{CH}_2\text{CHO} + 2\text{H}^+ + 2\text{e}^-$	(1)

Question Number	Answer	Additional Guidance	Mark
(iv)	<ul style="list-style-type: none"> <li>provides a surface for bubbles to form / enables smaller bubbles to form / provides nucleation sites for bubbles or to prevent large bubbles forming</li> </ul>	<p>Allow distribution of heat more evenly / to prevent superheating</p> <p>Ignore mixing / to stop bumping / spitting / explosion / liquid splashing out / vigorous reaction / loss of reactants</p> <p>Do not award reference to large gas molecules</p>	(1)

## Edexcel Chemistry A-level - Alcohols

Question Number	Answer	Additional Guidance	Mark
(v)	<ul style="list-style-type: none"> <li>• (M1) evaluation of number of moles of propan-1-ol (1)</li> </ul> <p>Method one using masses for percentage calculation</p> <ul style="list-style-type: none"> <li>• (M2) evaluation of maximum mass of propanal (1)</li> <li>• (M3) percentage yield (1)</li> </ul> <p>or</p> <p>Method two using moles for percentage calculation</p> <ul style="list-style-type: none"> <li>• (M2) evaluation of actual moles of propanal (1)</li> <li>• (M3) percentage yield (1)</li> </ul>	<p>Example of calculation</p> $n(\text{propan-1-ol}) = (1.50 \div 60) = 0.025 \text{ (mol)}$ $n(\text{propan-1-ol}) = n(\text{propanal})$ $\text{max } m(\text{propanal}) = (0.025 \times 58) = 1.45 \text{ (g)}$ $\% \text{Yield} = ((0.609 \div 1.45) \times 100) = 42 \%$  $n(\text{propanal}) = (0.609 \div 58) = 0.0105 \text{ (mol)}$ $\% \text{Yield} = ((0.0105 \div 0.025) \times 100) = 42 \%$ <p>Allow TE at each stage Ignore SF except 1SF Penalise incorrect <math>M_r</math> values once only Correct answer without working scores (3)</p>	(3)



## Edexcel Chemistry A-level - Alcohols

Q23.

Question Number	Acceptable Answers	Additional Guidance	Mark
(a)	<ul style="list-style-type: none"> <li>potassium dichromate(VI)/K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> <u>and</u> sulfuric acid/H<sub>2</sub>SO<sub>4</sub></li> <li>or</li> <li>sodium dichromate(VI)/Na<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> <u>and</u> (dilute) sulfuric acid/H<sub>2</sub>SO<sub>4</sub></li> </ul> <p style="text-align: right;">(1)</p> <ul style="list-style-type: none"> <li>heat/reflux</li> </ul> <p style="text-align: right;">(1)</p>	<p>Allow Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> <u>and</u> H<sup>+</sup> / acidified (potassium / sodium) dichromate(VI)</p> <p>If name and formula given, both must be correct</p> <p>Ignore concentration of acid</p> <p>Do not allow hydrochloric acid / HCl / nitric acid / HNO<sub>3</sub></p> <p>Conditional on correct reagents or near miss, provided dichromate or (per)manganate(VII) is mentioned</p> <p>Allow a specified temperature in the range 60 – 150°C</p> <p>Ignore distillation / warm</p> <p>Allow answers written on either dotted line</p>	(2)

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(i)	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li><b>Flask</b> - use of a volumetric / graduated flask</li> </ul> <p style="text-align: right;">(1)</p> <ul style="list-style-type: none"> <li><b>Weighing</b> - weigh the ethanedioic acid (in a weighed container and record the exact mass)</li> </ul> <p style="text-align: right;">(1)</p> <ul style="list-style-type: none"> <li><b>Dissolve, transfer and washings</b> – allow these in any order depending on the method used</li> </ul> <p style="text-align: right;">(1)</p> <ul style="list-style-type: none"> <li><b>Mark and mix</b> - make up to the mark / 250 cm<sup>3</sup> <u>and</u> then mix</li> </ul> <p style="text-align: right;">(1)</p>	<p>Ignore heat</p> <p>Do not allow just 'flask' / conical flask</p> <p>Ignore just 'put 1 /1.0 /1.09 g solid in beaker'</p> <p>Distilled / deionised water must be mentioned once in M3 or M4</p> <p>Allow pure water</p> <p>Allow any indication of mixing eg swirl / invert the flask</p>	(4)

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(ii)	<ul style="list-style-type: none"> <li>(From) colourless (to) pink</li> </ul>	Allow (to) red  Do not allow purple / pink/purple  Do not allow clear	(1)

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(iii)	<ul style="list-style-type: none"> <li>calculation of moles of NaOH (1)</li> <li>calculation of moles of H<sub>2</sub>C<sub>2</sub>O<sub>4</sub> in 25 cm<sup>3</sup> (1)</li> <li>calculation of moles of H<sub>2</sub>C<sub>2</sub>O<sub>4</sub> in 250 cm<sup>3</sup> (1)</li> <li>calculation of M<sub>r</sub> of crystals (1)</li> <li>calculation of value of n (1)</li> </ul>	Correct answer of 2.2582/2.258/2.26/2.3 without working scores 5 Final answer of 2, with working, resulting from a number between 2.2 and 2.3, scores 5 If no other mark is scored, an answer of just 2 scores 1  <u>Example of calculation</u> moles NaOH = $16.2 \times 0.103/1000 = 1.6686 \times 10^{-3}$  moles H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> in 25 cm <sup>3</sup> = $1.6686 \times 10^{-3}/2 = 8.343 \times 10^{-4}$ TE on mole NaOH  moles H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> in 250 cm <sup>3</sup> = $8.343 \times 10^{-4} \times 10 = 8.343 \times 10^{-3}$ TE on moles H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> in 25 cm <sup>3</sup>  M <sub>r</sub> of crystals = $1.09/8.343 \times 10^{-3} = 130.648 / 130.65 / 130.6$ TE on moles H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> in 250 cm <sup>3</sup>  For first 4 marking points ignore SF except 1 SF  $130.65 = (2 + (2 \times 12) + (4 \times 16)) + 18n$ $n = 2.2582/ 2.258/2.26/2.3/2$ TE on M <sub>r</sub> of crystals, provided n is positive	(5)

	<b>Alternative method for M4 and M5</b> <ul style="list-style-type: none"> <li>calculation of moles of H<sub>2</sub>O (1)</li> <li>calculation of value of n (1)</li> </ul>	mass H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> = $8.343 \times 10^{-3} \times 90 = 0.75087$ (g) mass H <sub>2</sub> O = $1.09 - 0.75087 = 0.3391$ (g) moles H <sub>2</sub> O = $0.3391/18 = 0.01884$  mole ratio H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> : H <sub>2</sub> O = 1 : $0.01884/8.343 \times 10^{-3} = 1 : 2.2582/2.258/2.26/2.3/2$	
--	--	--	--

## Edexcel Chemistry A-level - Alcohols

Question Number	Acceptable Answers	Additional Guidance	Mark
(b)(iv)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(damp crystals will have more water so) lower mass / moles / concentration of <math>\text{H}_2\text{C}_2\text{O}_4</math> (1)</li> <li>so titre will be lower and the value of n will be higher (1)</li> </ul>		(2)

Q24.


Question Number	Answer	Additional Guidance	Mark
	<p>A comparison that makes reference to:</p> <p>(with ethanoyl chloride)</p> <ul style="list-style-type: none"> <li>the reaction is irreversible compared to reversible (1)</li> <li>hydrogen chloride is the by-product rather than water (1)</li> <li>the reaction is very fast/occurs at room temperature so an acid catalyst is not needed (1)</li> </ul>	<p>Accept reverse arguments</p> <p>Allow steamy fumes for 'HCl'</p>	(3)


Q25.

Question Number	Answer	Additional Guidance	Mark
	<ul style="list-style-type: none"> <li>3-methylbutan-1-ol</li> </ul>	<p>Allow 'methly' for methyl</p> <p>Allow name with missing hyphens</p> <p>Allow 3-methylbutane-1-ol</p> <p>Allow 3-methylbut-1-anol</p> <p>Allow 1-hydroxy-3-methylbutane</p> <p>Do not allow 3-methylbut-1-ol</p> <p>Ignore formulae even if incorrect</p>	(1)

Edexcel Chemistry A-level - Alcohols

Q26.

Question Number	Answer	Additional Guidance	Mark
(i)	Any three of the following four structures		(3)
		<p>Accept formulae in any order</p> <p>Award (2) if 3 correct displayed/structural formulae given</p> <p>Award (1) if 2 correct displayed/structural formulae given</p>	

Question Number	Answer	Additional Guidance	Mark
(ii)	<p>An equation that has</p> <ul style="list-style-type: none"> <li>• ethanoyl chloride (1)</li> <li>• alcohol and ester+ HCl product (1)</li> </ul>	<p><u>Example of equation</u></p>  <p>Allow structural, displayed formulae in any combination</p> <p>Ignore connectivity to OH except horizontal</p> <p>Ignore state symbols even if incorrect</p> <p>If molecular formulae used then allow (1) for correct equation</p> <p>Allow (1) for a correct equation to form ester A from ethanoic acid e.g.</p> $\text{CH}_3\text{COOH} + \text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_2\text{CH}_3 \rightleftharpoons \text{CH}_3\text{COOCH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3 + \text{H}_2\text{O}$	(2)