

**Mark Scheme**

Q1.

Question Number	Acceptable Answer	Additional Guidance	Mark
	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"><li>• (add hydrochloric acid / nitric acid then) add barium chloride / barium nitrate (solution) <b>(1)</b></li><li>• white precipitate / white solid <b>(1)</b></li></ul>	<p>Ignore omission of acid Do not award just Barium ions / Ba<sup>2+</sup></p> <p>M2 is dependent on M1, with the exception of just Ba<sup>2+</sup> given as reagent</p>	<b>(2)</b>

Q2.

Question Number	Answer	Additional Guidance	Mark
	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>from chlorine to iodine / down the group, the number of electrons (in the molecule / atom) increases / changes from 34 to 106 / 17 to 53 <b>(1)</b></li> <li>so the strength of the London / instantaneous dipole-(induced) dipole forces increases / there are more London / instantaneous dipole-(induced) dipole forces <b>and</b> more energy is needed to separate the molecules <b>(1)</b></li> </ul>	<p>An answer that states 'covalent bonds break' or 'bonds between atoms break' or refers to 'ions' scores (0) overall</p> <p>Allow reverse argument for M1 and M2</p> <p>Allow iodine has more / most electron shells (than chlorine and/or bromine)</p> <p>Ignore 'the size of the atoms / molecules increases from chlorine to iodine'</p> <p>Do not allow incorrect numbers of electrons</p> <p>Allow iodine has the strongest London force <b>and</b> most energy is needed to separate the molecules</p> <p>Allow more energy is need to overcome / break the London forces / bonds instead of separate the molecules</p> <p>Allow dispersion forces / van der Waals forces for London forces</p> <p>Ignore higher temperature needed to separate the molecules</p> <p>Do not award dipole-dipole forces / just 'intermolecular forces'</p>	<b>(2)</b>

## Edexcel Chemistry A-level - Group 7

Q3.

Question Number	Acceptable Answer	Additional Guidance	Mark
	<p>An explanation that makes reference to the following points:</p> <p>Lower than 1 ppm</p> <ul style="list-style-type: none"> <li>• HClO will be low(er) <b>(1)</b></li> <li>• ineffective (as a disinfectant) <b>(1)</b></li> </ul> <p>Higher than 3 ppm</p> <ul style="list-style-type: none"> <li>• HCl will be high(er) <b>(1)</b></li> <li>• any relevant effect of increased HCl <b>(1)</b></li> </ul>	<p>Ignore reference to amount of Cl<sub>2</sub> being too low M2 dependent on correct M1</p> <p>M4 dependent on correct M3 Award effects including corrosive, alters or lowers pH NB: Do not award high(er) pH Award increases acidity / strongly acidic / toxicity Award any reasonable negative effect on swimmers e.g. irritation / irritant Ignore just 'harmful' / just 'dangerous' Ignore reference to amount of Cl<sub>2</sub> being too high and its effects</p>	<p><b>(4)</b></p>

Q4.

Question Number	Acceptable Answer	Additional Guidance	Mark
	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• add (excess) dilute ammonia / dilute NH<sub>3</sub> (to the precipitate) <b>and</b> the precipitate is insoluble / does not dissolve (1)</li> <li>• add (excess) concentrated (aqueous) ammonia / concentrated NH<sub>3</sub> (to the precipitate) <b>and</b> it is soluble / dissolves / forms a colourless solution (1)</li> </ul>	<p>Allow ammonium hydroxide for ammonia</p> <p>Ignore pure ammonia / ammonia with no concentration / ammonia gas</p> <p>Allow no change for the observation</p> <p>Allow 'if it dissolves it is not bromide'</p> <p>Allow redissolves for soluble</p> <p><b>Note</b> If no other mark is awarded allow (1) for adding dilute and concentrated ammonia with no / incorrect observation(s)</p> <p><b>Alternative test:</b> add concentrated sulfuric acid (1) brown fumes (1)</p>	(2)

Q5.

Question Number	Acceptable Answers	Additional Guidance	Mark
	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"><li>• to react with / remove any carbonate / sulfite / sulfate(IV) (ions)</li></ul>	<p>Allow equation for the reaction with acid e.g. <math>2\text{H}^+ + \text{CO}_3^{2-} \rightarrow \text{H}_2\text{O} + \text{CO}_2</math></p> <p>Allow to prevent any other ions forming a precipitate (with barium ions / <math>\text{Ba}^{2+}</math>)</p> <p>Allow to rule out the possibility of carbonate / sulfite / sulfate(IV) ions giving a false result</p> <p>Ignore just 'to remove impurities / other (an)ions' / 'react with precipitates'</p> <p>Ignore to dissolve barium carbonate / sulfite</p> <p>Ignore reference to hydrogencarbonate</p>	<b>(1)</b>

Q6.

Question Number	Answer	Additional Guidance	Mark
(i)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>balanced equation (1)</li> <li>calculation of <math>E^{\ominus}_{\text{cell}}</math> value (1)</li> <li><math>E^{\ominus}_{\text{cell}}</math> / answer is negative / &lt;0 <b>and</b> the reaction is not (thermodynamically) feasible (1)</li> </ul>	<p><u>Example of equation</u></p> <p><math>\text{Br}_2(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{HOBr}(\text{aq}) + \text{HBr}(\text{aq})</math>            Allow multiples            Allow <math>\text{H}^+(\text{aq}) + \text{Br}^-(\text{aq})</math> for <math>\text{HBr}(\text{aq})</math>            Allow reversible arrows            Ignore state symbols even if incorrect</p> <p><math>E^{\ominus}_{\text{cell}} = 1.09 - 1.57 = -0.48 \text{ (V)}</math>            Allow correct answer without calculation</p> <p>Allow 3 marks for reverse argument  <math>\text{HOBr}(\text{aq}) + \text{HBr}(\text{aq}) \rightarrow \text{Br}_2(\text{aq}) + \text{H}_2\text{O}(\text{aq})</math> (1)  <math>E^{\ominus}_{\text{cell}} = 1.57 - 1.09 = (+) 0.48 \text{ (V)}</math> (1)  <math>E^{\ominus}_{\text{cell}}</math> is positive / &gt;0 so the reverse of disproportionation is (thermodynamically) feasible (1)</p>	(3)

Question Number	Answer	Additional Guidance	Mark
(ii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>disproportionation is an equilibrium system (and although K is very small, there is still a small concentration of disproportionation products)  <b>or</b>            excess water is used  <b>or</b>            concentration is not <math>1 \text{ mol dm}^{-3}</math>  <b>or</b>            HOBr undergoes further disproportionation</li> </ul>	<p>Ignore just 'non-standard conditions'</p> <p>Ignore references to activation energy / collision theory            Ignore <math>\text{H}^+</math> / ions from the water</p>	(1)

## Edexcel Chemistry A-level - Group 7

Q7.

Question Number	Answer	Additional Guidance	Mark
(i)	<ul style="list-style-type: none"> <li>balanced equation</li> </ul>	<p><u>Example of equation</u>  <math>3\text{Cl}_2 + 6\text{OH}^- \rightarrow 5\text{Cl}^- + \text{ClO}_3^- + 3\text{H}_2\text{O}</math>                      Allow multiples                      Ignore state symbols even if incorrect</p>	(1)

Question Number	Answer	Additional Guidance	Mark
(ii) Clip with (i)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>oxidation number for chlorine changes from 0 to -1 so it is <b>reduced</b> (1)</li> <li>oxidation number for chlorine changes from 0 to +5 so it is <b>oxidised</b> (1)</li> </ul>	<p>Ignore general definitions of disproportionation</p> <p>Accept oxidation numbers and their changes shown with equation</p> <p>Allow 1 out of 2 marks for three correct oxidation numbers of the chlorine</p>	(2)

Q8.

Question Number	Acceptable Answer	Additional Guidance	Mark
	<p><math>\text{ClO}_3^-</math> (1)</p> <p>(Cl is) +5 / 5+ (1)</p>	<p>Allow <math>\text{NaClO}_3</math> / <math>\text{KClO}_3</math></p> <p>Allow (+)V</p> <p>Do not award 5 unless +5/5+ seen in the formula or as a label on the formula</p>	(2)

Q9.

Question Number	Answer	Additional Guidance	Mark
	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• iodide ions are the strongest reducing agent because iodide ions / <math>I^-</math> / (potassium) iodide reduces sulfur (in sulfuric acid) from +6 to 0 in sulfur / -2 in <math>H_2S</math> <b>(1)</b></li> <li>• (whereas) bromide ions / <math>Br^-</math> / (potassium) bromide reduces sulfur (in sulfuric acid) from +6 to +4 <b>(1)</b></li> <li>• (whereas) chloride ions / <math>Cl^-</math> / (potassium) chloride do not reduce sulfuric acid / sulfur / S (as there is no change in oxidation number of Cl or S) <b>(1)</b></li> </ul>	<p>Allow the oxidation numbers written by the species in the table</p> <p>(+)6 only needs to be mentioned once in M1 or M2</p> <p>Allow references to potassium halides / halogens / hydrogen halides instead of halide ions</p> <p>For full marks, the answer must identify iodide as the strongest reducing agent</p> <p>Only 1 oxidation number change is needed. If both are given, both must be correct</p> <p>Allow bromide ions are stronger reducing agents than chloride ions because they are oxidised from -1 to 0</p> <p>Allow just 'it is not a redox reaction'</p>	<b>(3)</b>



Q10.

Question Number	Acceptable Answer	Additional Guidance	Mark
(i)	An answer that makes reference to the following points: <ul style="list-style-type: none"> <li>the covalent bond in hydrogen chloride changes to an ionic bond in aqueous solution</li> </ul>	Both types of bond required Accept covalent bond breaks, ions are formed Accept $\text{HCl(g)} \rightarrow \text{H}^{\text{+}}(\text{aq}) + \text{Cl}^{-}(\text{aq})$ or $\text{HCl(g)} + \text{H}_2\text{O(l)} \rightarrow \text{H}_3\text{O}^{\text{+}}(\text{aq}) + \text{Cl}^{-}(\text{aq})$	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	<ul style="list-style-type: none"> <li>correct species on each side of equation (1)</li> <li>correct states for all species (1)</li> </ul>	<u>Example of equation:</u> $\text{HCl(g)} + \text{NH}_3(\text{g}) \rightarrow \text{NH}_4\text{Cl(s)} / \text{NH}_4^{\text{+}}\text{Cl}^{-}(\text{s}) / \text{NH}_4^{\text{+}}(\text{s}) + \text{Cl}^{-}(\text{s})$  Allow (aq) or (g) for reactants Do not award (liquid) for either reactant Two products will lose both marks	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
(iii)	An answer that makes reference to the following points: <ul style="list-style-type: none"> <li>first observation (1)</li> <li>second observation (1)</li> </ul>	Allow observations in any order  Sodium carbonate/ $\text{Na}_2\text{CO}_3$ /(white) solid dissolves/disappears/forms a colourless solution  Effervescence/fizzing/bubbles Ignore gas/carbon dioxide given off Do not award if any named gas other than carbon dioxide, eg hydrogen or oxygen	(2)

Edexcel Chemistry A-level - Group 7

Question Number	Acceptable Answer	Additional Guidance	Mark
(iv)	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• remove a fixed amount of one solution using a pipette into a conical flask and fill up the burette with other solution (1)</li> <li>• add a named indicator <b>and</b> colour change (1)</li> <li>• add solution from (1) burette to flask until indicator changes colour</li> <li>• technique mark (1)</li> <li>• repeat titrations (until concordant results obtained) (1)</li> </ul>	<p>Allow use of any suitable flask in place of conical flask.</p> <p>Allow any recognised acid/base indicator: methyl red / orange, phenolphthalein etc. Ignore litmus /UI. Do not award reversed colour change</p> <p>Do not penalise reverse colour change again here.</p> <p>Any one from: Rinsing burette/pipette with appropriate solution, use of white tile, adding slowly, swirling flask etc.</p> <p>Ignore mention of 'rough' or 'trial' runs etc</p>	(5)

## Edexcel Chemistry A-level - Group 7

Q11.

Question Number	Acceptable Answer	Additional Guidance	Mark
	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• ammonium (1)</li> <li>• bromide (1)</li> </ul>	<p>Mark independently</p> <p>Allow names in either order</p> <p>Ignore symbols as well as names</p> <p>Do not award ammonia</p> <p>Do not award bromine</p> <p>Allow (1) for just NH<sub>4</sub>Br</p>	(2)

Q12.

Question Number	Acceptable Answers	Additional Guidance	Mark
	<ul style="list-style-type: none"> <li>• correct equation</li> </ul>	<p><u>Example of equation</u>  <math>\text{NH}_4^+ + \text{OH}^- \rightarrow \text{NH}_3 + \text{H}_2\text{O}</math></p> <p>Allow multiples</p> <p>Allow Na<sup>+</sup> on both sides if crossed through</p> <p>Ignore state symbols even if incorrect</p>	(1)

Q13.

Question Number	Acceptable Answer	Additional Guidance	Mark
(i)	$\text{Cl}_2 + 2\text{Br}^- \rightarrow 2\text{Cl}^- + \text{Br}_2$	<p>Allow multiples</p> <p>Ignore state symbols even if incorrect</p>	(1)

## Edexcel Chemistry A-level - Group 7

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>chlorine/bromine toxic/poisonous (1)</li> <li>(Carry out the experiment in a) fume cupboard (1) OR</li> <li>bromine corrosive (1)</li> <li>wear gloves (1)</li> </ul>	<p>2nd mark dependent on first.</p> <p>Do not award harmful, but allow MP2 if correct for toxic.</p>	(2)

### Q14.

Question Number	Answer	Additional Guidance	Mark
(i)	<ul style="list-style-type: none"> <li>ionic equation (1)</li> <li>state symbols (1)</li> </ul>	<p><u>Example of equation</u>  <math>\text{Ag}^+(\text{aq}) + \text{I}^-(\text{aq}) \rightarrow \text{AgI}(\text{s})</math>            Allow multiples</p> <p>M2 dependent on M1 or near miss</p>	(2)

Question Number	Answer	Additional Guidance	Mark
(ii)	<p>An answer that includes</p> <ul style="list-style-type: none"> <li>halide ion with some justification attempt (1)</li> <li>calculation of expected mass of silver halides (1)</li> </ul>	<p>Incorrect halide scores (0)</p> <p>Bromide (ion)/Br<sup>-</sup>            Do not award 'bromine (ion)'</p> <p>0.01 mol of AgCl = 1.43 (g)            AgBr = 1.88 (g) AgI = 2.35 (g)</p> <p>OR            Mass of 1.0 mol is 188 g so subtraction of 107.9 for Ag means X = 80.1 so closest is Br            TE on incorrect formula silver halide in d(i)</p>	(2)

Edexcel Chemistry A-level - Group 7

Q15.

Question Number	Answer	Additional Guidance	Mark
(i)	<ul style="list-style-type: none"> <li>(The cation in X is) <math>\text{Fe}^{2+}</math> / iron(II) / Fe(II) (1)</li> <li>(The anion in X is) <math>\text{SO}_4^{2-}</math> / sulfate(VI) (1)</li> </ul>	Allow $\text{Fe}^{+2}$ Allow sulfate / $\text{SO}_4^{2-}$ Do not award sulfite / sulfate(IV)	(2)

Question Number	Answer	Additional Guidance	Mark
(ii)	<ul style="list-style-type: none"> <li>species and balancing (1)</li> <li>state symbols (1)</li> </ul>	<p><u>Examples of equation</u></p> $\text{Fe}^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) \rightarrow \text{Fe}(\text{OH})_2(\text{s})$ or $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) \rightarrow \text{Fe}(\text{OH})_2(\text{s}) + 6\text{H}_2\text{O}(\text{l})$ or $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) \rightarrow \text{Fe}(\text{OH})_2(\text{H}_2\text{O})_4(\text{s}) + 2\text{H}_2\text{O}(\text{l})$ Ignore missing square brackets  TE on cation that forms an insoluble hydroxide in Test 1  State symbols conditional on correct species or 'near miss' / non-ionic equation	(2)

Question Number	Answer	Additional Guidance	Mark
(iii)	An answer that makes reference to the following point: <ul style="list-style-type: none"> <li><math>\text{Fe}^{2+}</math> is oxidised (to <math>\text{Fe}^{3+}</math>) by oxygen / air</li> </ul>	Allow iron(III) hydroxide / iron(III) (ions) are formed by reaction with oxygen / air  TE on cation in Test 1  Allow just 'the precipitate / it is oxidised by oxygen / air'	(1)

Question Number	Answer	Additional Guidance	Mark
(iv)	An answer that makes reference to the following point: <ul style="list-style-type: none"> <li>to react with / remove any carbonate / sulfite / sulfate(IV) ions or to eliminate the possibility of carbonate / sulfite / sulfate(IV) ions</li> </ul>	Allow to prevent any other ions forming a precipitate with barium ions / $\text{Ba}^{2+}$	(1)

Edexcel Chemistry A-level - Group 7

Q16.

Question Number	Acceptable Answer	Additional Guidance	Mark
(i)	$\text{Cl}_2 + 2\text{e}^{(-)} \rightarrow 2\text{Cl}^-$	Allow multiples Ignore state symbols even if incorrect	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	$\text{Cl}_2 + 4\text{OH}^- \rightarrow 2\text{ClO}^- + 2\text{H}_2\text{O} + 2\text{e}^{(-)}$	Allow multiples $\text{Cl}_2 + 2\text{OH}^- \rightarrow 2\text{ClO}^- + 2\text{H}^+ + 2\text{e}^{(-)}$ Ignore state symbols even if incorrect	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
(iii)	$\text{Cl}_2 + 2\text{OH}^- \rightarrow \text{Cl}^- + \text{ClO}^- + \text{H}_2\text{O}$	Allow multiples Ignore state symbols even if incorrect Do not award mark if electrons are un-cancelled	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
(iv)	An explanation that makes reference to the following points: <ul style="list-style-type: none"> <li>(disproportionation is simultaneous) oxidation and reduction of an element (in the same species) (1)</li> <li>chlorine changes from 0 to -1 <b>and</b> +1 (1)</li> </ul>	Allow statement that chlorine is oxidised <b>and</b> reduced  This can be shown on the equation in (iii)	(2)

Q17.

Question Number	Acceptable Answers	Additional Guidance	Mark
(i)	<p>An answer that makes reference to the following points:</p> <p><b>setting up of the dipole</b></p> <ul style="list-style-type: none"> <li>uneven distribution of <b>electrons</b>/ (random) movement of <b>electrons</b> / (random) fluctuations of <b>electrons</b> (1)</li> </ul> <p><b>type of dipole</b></p> <ul style="list-style-type: none"> <li>(results in an) instantaneous dipole / temporary dipole (in the first molecule) (1)</li> </ul> <p><b>induction of a second dipole</b></p> <ul style="list-style-type: none"> <li>causes/induces a (second) dipole on another molecule (1)</li> </ul>	<p>M1 &amp; M3 could be scored for an appropriate diagram</p> <p>Allow "Change in <b>electron</b> density"</p> <p>Allow "transient dipole" / "oscillating dipole" Do not award for "permanent dipole"</p> <p>Allow neighbouring molecule / adjacent molecule Do not award for "permanent dipole"</p>	(3)

Question Number	Acceptable Answers	Additional Guidance	Mark
(ii)	<p>An explanation that makes reference to the following points:</p> <p><b>relative number of electrons</b></p> <ul style="list-style-type: none"> <li>bromine has more <b>electrons</b> (than chlorine) / bromine has one more shell of <b>electrons</b> (than chlorine) (1)</li> </ul> <p><b>relative strength of intermolecular forces</b></p> <ul style="list-style-type: none"> <li>(so) bromine has stronger (London) forces (between molecules) / more (heat) energy is needed to overcome the London forces between bromine molecules / greater temporary dipole – induced dipole forces (1)</li> </ul>	<p>Allow reverse arguments Allow correct formulae</p> <p>Bromine has 35/70 electrons and chlorine has 17/34 electrons</p> <p>Ignore comments about protons, molecular mass etc</p> <p>Do not award "more outer shells"</p> <p>Ignore comments about 'points of contact' Allow more (London) forces Allow "bonds between molecules"</p> <p>Award (0) marks overall if any implication that <b>covalent</b> bonds are broken (on boiling)</p>	(2)

## Edexcel Chemistry A-level - Group 7

Q18.

Question Number	Acceptable Answers	Additional Guidance	Mark
(i)	<p><b>Test 1</b></p> <ul style="list-style-type: none"> <li>(gas is) ammonia / <math>\text{NH}_3</math> (1)</li> <li>(cation is) ammonium / <math>\text{NH}_4^+</math> (1)</li> </ul> <p><b>Test 2</b></p> <ul style="list-style-type: none"> <li>(cation is) cobalt(II) / <math>\text{Co}^{2+}</math> / <math>[\text{Co}(\text{H}_2\text{O})_6]^{2+}</math> (1)</li> <li>(complex ion is) <math>[\text{CoCl}_4]^{2-}</math> (1)</li> </ul> <p><b>Test 3</b></p> <ul style="list-style-type: none"> <li>(precipitate is) barium sulfate / <math>\text{BaSO}_4</math> □ (1)</li> <li>(anion is) sulfate(VI) / <math>\text{SO}_4^{2-}</math> (1)</li> </ul>	<p>If name and formula are given, both must be correct Mark independently</p> <p>Do not award gas is ammonium / <math>\text{NH}_4</math> Do not award cation is ammonia / <math>\text{NH}_3^+</math></p> <p>Oxidation number of cobalt is needed in the name but allow cobalt <b>with</b> <math>\text{Co}^{2+}</math> Charge is needed on the ion Allow +2 and -2 for the charges on the ions Allow brackets around Cl Ignore missing square brackets in complex ions Do not award <math>[\text{CoCl}_6]^{4-}</math></p> <p><b>Note</b> If cation in Test 2 is identified as copper(II) / <math>\text{Cu}^{2+}</math>, do not award M3 but M4 can be awarded as TE for <math>[\text{CuCl}_4]^{2-}</math></p> <p>Oxidation number of sulfate is not needed but if given must be correct e.g. do not award sulfate(IV)</p>	(6)

Question Number	Acceptable Answers	Additional Guidance	Mark
(ii)	<ul style="list-style-type: none"> <li>correct formula</li> </ul>	<p><u>Examples of correct formula</u> <math>(\text{NH}_4)_2\text{Co}(\text{SO}_4)_2</math> <math>(\text{NH}_4)_2\text{SO}_4 \cdot \text{CoSO}_4</math></p> <p>Allow <math>\text{NH}_4</math>, Co and <math>\text{SO}_4</math> in any order</p> <p>Allow multiples</p> <p>Allow any combination of <math>\text{Co}^{2+}</math> / <math>\text{NH}_4^+</math> / <math>\text{SO}_4^{2-}</math> that gives a neutral complex</p> <p>TE on the <b>three</b> ions identified in (a)(i)</p> <p>Ignore missing dot in second formula</p> <p>Ignore any amount of water of crystallisation</p>	(1)



Edexcel Chemistry A-level - Group 7

Q19.

Question Number	Acceptable Answers	Additional Guidance	Mark
	<p>An answer that makes reference to the following points:</p> <p><b>mixing of 1<sup>st</sup> pair of solutions</b></p> <ul style="list-style-type: none"> <li>• mix Br<sub>2</sub> with KCl (1)</li> </ul> <p><b>mixing of 2<sup>nd</sup> pair of solutions</b></p> <ul style="list-style-type: none"> <li>• mix Br<sub>2</sub> with KI or mix I<sub>2</sub> with KBr (1)</li> </ul> <p><b>colours of halogen (in cyclohexane)</b></p> <ul style="list-style-type: none"> <li>• colour seen for experiment 1/ bromine is orange / yellow <b>and</b> colour seen for experiment 2/ iodine is purple / pink / violet / lilac (1)</li> </ul> <p><b>correct ionic equation</b></p> <ul style="list-style-type: none"> <li>• Br<sub>2</sub> + 2I<sup>-</sup> → 2Br<sup>-</sup> + I<sub>2</sub> (1)</li> <li>• use of ONLY two correct experiments as above (1)</li> </ul>	<p>Ignore any reference to any additional reactions, e.g. with silver nitrate</p> <p>Award mark if correct ionic equation is given</p> <p>Ignore colours before the addition of cyclohexane</p> <p>Do not award brown</p> <p>Do not award red</p> <p>Allow multiples</p> <p>Ignore state symbols even if incorrect</p>	(5)

Q20.

Question Number	Answer	Additional Guidance	Mark
(i)	<ul style="list-style-type: none"> <li>• chlorine is oxidised <b>and</b> from 0 to +1 (in NaClO) (1)</li> <li>• chlorine is reduced <b>and</b> from 0 to -1 (in NaCl) (1)</li> </ul>	<p>Check the equation</p> <p>Allow (1) for three correct oxidation numbers if no other mark is awarded.</p> <p>Allow (1) max for general definition of disproportionation</p>	(2)
(ii)	<ul style="list-style-type: none"> <li>• equation</li> </ul>	<p>6 NaOH + 3 Cl<sub>2</sub> → NaClO<sub>3</sub> + 5 NaCl + 3 H<sub>2</sub>O</p> <p>Allow multiples</p>	(1)

## Edexcel Chemistry A-level - Group 7

Question Number	Answer	Mark
(iii)	<p>The only correct answer is C (hot alkali)</p> <p>A is not correct because high temperature is required</p> <p>B is not correct because high temperature is required</p> <p>D is not correct because high temperature and not excess chlorine is required</p>	(1)

### Q21.

Question Number	Answer	Additional Guidance	Mark
(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(1) • chlorine / Cl<sub>2</sub> is simultaneously oxidised and reduced</li> <li>(1) • the oxidation number of chlorine changes from 0 to -I and (+)I / 0 to -1 and (+)1 / increases by 1 and decreases by 1</li> </ul>	<p>Allow oxidation numbers underneath or above the equation</p>	(2)

Question Number	Answer	Additional Guidance	Mark
(ii)	<ul style="list-style-type: none"> <li>(1) • substances correct in equation</li> <li>(1) • equation is balanced</li> </ul>	<p><u>Example of equation</u></p> $3\text{Cl}_2 + 6\text{NaOH} \rightarrow \text{NaClO}_3 + 5\text{NaCl} + 3\text{H}_2\text{O}$ <p>Ignore state symbols even if incorrect</p>	(2)

## Edexcel Chemistry A-level - Group 7

Q22.

Question Number	Acceptable Answers	Additional Guidance	Mark
(i)	<ul style="list-style-type: none"> <li>chlorine is a gas <b>and</b> iodine is a solid</li> </ul>	Ignore any colours, even if incorrect Do not award reference to ions once in (i)/(ii) Allow use of Cl and I	<b>(1)</b>
(ii)	<ul style="list-style-type: none"> <li>astatine is a solid <b>and</b>                      (as the number of electrons increases) the strength of the London forces increases / more London forces</li> </ul>	Allow for 'London forces' instantaneous dipole – induced dipole / van der Waals' forces / dispersion forces / induced dipole forces / temporary dipole  Ignore ID-ID References to stability The trend is increasing melting (and boiling) temperature down the group	<b>(1)</b>

Q23.

Question Number	Acceptable Answer	Additional Guidance	Mark
(i)	An answer that makes reference to the following points. <ul style="list-style-type: none"> <li>cool (the reaction mixture) <b>(1)</b></li> <li>filter off (the less soluble potassium iodate) <b>(1)</b></li> <li>any suitable method of drying (the resulting solid) <b>(1)</b></li> </ul>	Mark independently Ignore addition of extra water  Allow give time for potassium iodate to crystallise Ignore the method used to cool the solution, (ice, fridge etc.)  Ignore any details of the filtration methods  Examples of methods used to dry: 'leave to dry', warm oven, press between filter papers	<b>(3)</b>

## Edexcel Chemistry A-level - Group 7

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>add silver nitrate (solution) / <math>\text{AgNO}_3</math> (and <math>\text{HNO}_3</math> / nitric acid) (1)</li> <li>yellow <b>and</b> precipitate /ppt/solid/crystals (1)</li> </ul>	<p>Mark independently</p> <p>Do not award hydrochloric acid</p> <p>Allow a correct description of the yellow ppt, e.g. primrose coloured</p> <p>Do not award 'electrolysis'</p>	(2)

Q24.

Question Number	Answer	Additional Guidance	Mark
	<p>An answer that makes reference to:</p> <ul style="list-style-type: none"> <li>addition of (dilute/strong) name/formula of acid (1)</li> <li>effervescence/bubbling/fizzing (1)</li> </ul>	<p>Allow weak acids If formula given then must be correct</p> <p>Allow Gas given off which turns limewater cloudy</p> <p>Do not award just 'gas/ <math>\text{CO}_2</math> given off' Do not award incorrect observations such as precipitate forming due to addition of acid</p> <p>M2 dependent on M1 or 'near miss'</p>	(2)

## Edexcel Chemistry A-level - Group 7

Q25.

Question Number	Answer	Additional Guidance	Mark
	<p>An answer that makes reference to:</p> <ul style="list-style-type: none"> <li>addition of barium chloride/nitrate (solution) (1)</li> <li>white precipitate forms (1)</li> </ul>	<p>Accept formulae BaCl<sub>2</sub>/Ba(NO<sub>3</sub>)<sub>2</sub></p> <p>Ignore addition of acids such as HCl or HNO<sub>3</sub> but do not award M1 if addition of sulfuric acid</p> <p>Allow white solid If ppt identified then must be correct</p> <p>M2 dependent on M1 or 'near miss'</p>	(2)

Q26.

Question Number	Acceptable Answers	Additional Guidance	Mark
	$\text{Cl}_2 + 2\text{NaOH} \rightarrow \text{NaCl} + \text{NaClO} + \text{H}_2\text{O}$ <p>(1)</p> <ul style="list-style-type: none"> <li>(Type of reaction) disproportionation (1)</li> </ul>	<p>Accept multiples and ionic equations. Allow NaOCl</p> $\text{Cl}_2 + 2\text{OH}^- \rightarrow \text{Cl}^- + \text{ClO}^- + \text{H}_2\text{O}$ <p>Ignore state symbols even if incorrect</p> <p>Ignore redox Mark independently</p>	(2)

Q27.

Question Number	Answer	Mark
(i)	<p>The only correct answer is C (Disproportionation)</p> <p><i>A is not correct because oxidation and reduction are occurring</i></p> <p><i>B is not correct because oxidation and reduction are occurring</i></p> <p><i>D is not correct because two reactants are involved</i></p>	(1)

## Edexcel Chemistry A-level - Group 7

Question Number	Acceptable Answer	Additional Guidance	Mark
(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(trigonal) pyramidal (1)</li> <li>(predicted bond angle) <math>107^\circ</math> (1)</li> <li>three groups / three pairs of bonding electrons and one lone pair OR lone pair – bond pair repulsion &gt; bond pair – bond pair repulsion (1)</li> <li>(electron pairs / groups repel to positions of) minimum repulsion / maximum separation (1)</li> </ul>	<p>For M1, this shape must be <b>named</b></p> <p>Allow answers in the range <math>106.5^\circ</math> to <math>107.5^\circ</math> (allow actual value <math>110^\circ</math>) Allow M2 on an annotated diagram</p> <p>Allow 'regions' for 'groups' or 'pairs'</p> <p>Allow statements such as "lone pair repulsion greater than bond pair repulsion"</p>	(4)

Q28.

Question Number	Answer	Mark
(i)	<p><b>The only correct answer is C (redox)</b></p> <p><i>A is not correct because chlorine is reduced and iodide is oxidised</i></p> <p><i>B is not correct because chlorine is reduced and iodide is oxidised</i></p> <p><i>D is not correct because different species are oxidised and reduced</i></p>	(1)

Question Number	Answer	Mark
(ii)	<p><b>The only correct answer is A (pale yellow)</b></p> <p><i>B is not correct because the question refers to the aqueous layer</i></p> <p><i>C is not correct because the question refers to the aqueous layer</i></p> <p><i>D is not correct because this would be the colour of <math>Cl_2(aq)</math> in the absence of <math>I_2(aq)</math></i></p>	(1)

## Edexcel Chemistry A-level - Group 7

Q29.

Question Number	Answer	Additional Guidance	Mark
(i)	An answer that makes reference to the following point: <ul style="list-style-type: none"> <li>the oxidation number / state does not change for any element</li> </ul>	Accept there is no transfer of electrons	(1)

Question Number	Answer	Additional Guidance	Mark
(ii)	An explanation that makes reference to the following points: <ul style="list-style-type: none"> <li>(because) sulfur in sulfuric acid is reduced further by hydrogen iodide than hydrogen bromide (and hydrogen chloride)</li> <li>SO<sub>2</sub> / S(IV) produced in the reaction with HBr</li> <li>more negative oxidation states of sulfur / S / H<sub>2</sub>S / S<sup>2-</sup> are produced in the reaction with HI</li> </ul>	Mark independently (1) Allow potassium salt / halide ion for hydrogen halide (1) May be shown in an equation, but ignore incorrect state symbols and/or balancing (1) May be shown in an equation, but ignore incorrect state symbols and/or balancing	(3)

Q30.

Question Number	Answer	Additional Guidance	Mark
(i)	<ul style="list-style-type: none"> <li>equation</li> </ul>	<u>Example of equation</u> $\text{Cl}_2 + 2\text{KBr} \rightarrow \text{Br}_2 + 2\text{KCl}$ Accept ionic equation $\text{Cl}_2 + 2\text{Br}^- \rightarrow \text{Br}_2 + 2\text{Cl}^-$ Allow multiples Ignore state symbols even if incorrect	(1)

## Edexcel Chemistry A-level - Group 7

Question Number	Answer	Additional Guidance	Mark
(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• <b>(M1)</b> the precipitate is a mixture of silver chloride and silver bromide or not all of the bromide ions were oxidised <b>(1)</b></li> <li>• <b>(M2)</b> silver chloride/AgCl dissolves in dilute ammonia <b>(1)</b></li> <li>• <b>(M3)</b> silver bromide/AgBr does not dissolve in dilute ammonia <b>(1)</b></li> </ul>	<p>Allow Some bromide ions are still present/bromide ions were in excess/both chloride and bromide ions are present</p> <p>Do not award references to Cl<sup>-</sup> dissolving</p> <p>Silver bromide only dissolves in concentrated ammonia</p>	<b>(3)</b>

Question Number	Answer	Additional Guidance	Mark
(iii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>• iodine is a weaker oxidising agent than chlorine or iodine cannot oxidise bromide ions or iodine is a stronger reducing agent</li> </ul>	<p>Accept reverse arguments</p> <p>Ignore Just references to reactivity/displacement e.g. iodine is less reactive/cannot displace</p>	<b>(1)</b>

Q31.

Question Number	Answer	Mark	
(i)	<ul style="list-style-type: none"> <li>• silver nitrate (solution) / chlorine</li> </ul>	<p>Allow correct formula/AgNO<sub>3</sub> If both name and formula are given both must be correct Allow acidified silver nitrate (solution) Ignore addition of nitric acid Do not award sulfuric acid / hydrochloric acid</p> <p>Do not award conc. sulfuric acid here but allow TE in dii</p>	<b>(1)</b>



## Edexcel Chemistry A-level - Group 7

Question Number	Acceptable Answer	Additional Guidance	Mark
	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• cream/off-white precipitate <b>(1)</b></li> <li>• AgBr <b>(1)</b></li> </ul>	<p>Do not accept just 'white' or 'yellow' Accept (very) pale yellow</p> <p>Ignore name Ignore unbalanced equation</p> <p>Award (2) marks for use of chlorine: orange / brown fumes / solution Br<sub>2</sub>(gas / aq)</p> <p>Allow TE (2) marks for use of conc. sulfuric acid in 3di choking fumes SO<sub>2</sub> (g)</p>	<p><b>(2)</b></p>