

Mark Scheme

Q1.

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---------------------|------|
| | An answer that includes <ul style="list-style-type: none"> (estimated value) between 1100 – 1380 (kJ mol⁻¹) | | (1) |

Q2.

| Question Number | Answer | Mark |
|-----------------|--|------|
| (i) | The only correct answer is C ($S(g) \rightarrow S^+(g) + e^-$) <i>A is not correct because the sulfur must be in the gas phase</i> <i>B is not correct because the sulfur must be individual atoms and in the gas phase</i> <i>D is not correct because the sulfur must be individual atoms</i> | (1) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| (ii) | An explanation that makes reference to the following points: <ul style="list-style-type: none"> first ionisation energy decreases down the group because although the number of protons is increasing (1) the electron being removed is (one shell of electrons) further from the nucleus (1) (with one shell of electrons) giving more shielding from the nucleus (1) | Allow greater repulsion between inner electron shells | (3) |

Edexcel Chemistry A-level - Periodicity

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| (iii) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> because in sulfur the nuclear charge / atomic number / proton number / number of protons has is less (by 1) (1) and the electron being removed is from the same sub-shell / a (3)p electron / has similar shielding / is further from the nucleus / (1) | <p>Do not award just 'the charge has decreased (by 1) in sulfur' Allow effective nuclear charge has decreased by 1 in sulfur</p> <p>Allow has the same shielding Allow atomic radius is larger Do not award ionic radius is larger Ignore same shell</p> <p>Allow reverse arguments for chlorine</p> | (2) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| (iv) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> because in sulfur (spin) pairing has occurred (for the first time in the 3p sub-shell) or electron being removed from an orbital containing two electrons (1) (resulting in an increase in) repulsion between electrons (so the electron is lost more easily) (1) | <p>Ignore half-filled (sub-) shell is more stable in phosphorus</p> <p>Ignore reference to shielding and distance to the nucleus</p> | (2) |

Q3.

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> a trend/pattern of repeating (physical and chemical) properties (with increasing atomic number) (1) atomic radii decrease from left to right/ across the period (1) the pattern /atomic radius trend is repeated in period 3 (1) | <p>Do not award for trend in group</p> <p>Allow a sketch of the trend</p> <p>Allow even if the trend is incorrect</p> | (3) |

Q4.

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|---|--|------------|
| | <p>An answer that makes reference to the following:</p> <p>(in favour)</p> <ul style="list-style-type: none"> • electronic structure of hydrogen is s^1 / $1s^1$ / has one electron in s orbital / form $1+$ ions (1) <p>(against) any two from</p> <ul style="list-style-type: none"> • the rest of Group 1 are (alkali) metals / metallic (hydrogen is not) (1) • hydrogen does not react in the same way as / has different reactivity to the rest of Group 1 / has different chemical properties (1) • forms a H^- ion (1) | <p>Allow 1 electron in outer shell / has 1 valence electron Do not award 'last electron is in s orbital' unless it is clear there is only one Do not award just 'single unpaired electron'</p> <p>Allow hydrogen is not a metal Ignore hydrogen is a gas but Group 1 elements are solid</p> <p>Do not award just 'different properties' or 'different behaviour' Allow hydrogen forms covalent bonds as a chemically different property Ignore trends in physical properties</p> <p>Allow hydrogen can gain one electron to form a stable ion / become stable / fill its outer shell</p> | (3) |

Edexcel Chemistry A-level - Periodicity

Q5.

| Question Number | Acceptable Answer | Additional Guidance | Mark | | | | | | | | | | | | | | | | |
|--|---|--|---|---|---|-----|---|-----|---|---|---|---|---|--|--|--|---|--|-----|
| * | <p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="379 600 860 869"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table border="1" data-bbox="379 987 860 1279"> <thead> <tr> <th></th> <th>Number of marks awarded for structure and sustained lines of reasoning</th> </tr> </thead> <tbody> <tr> <td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning</td> <td>2</td> </tr> </tbody> </table> | Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | 6 | 4 | 5-4 | 3 | 3-2 | 2 | 1 | 1 | 0 | 0 | | Number of marks awarded for structure and sustained lines of reasoning | Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning | 2 | <p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning, scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> <p>In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.</p> <p>If there is any incorrect chemistry, deduct mark(s) from the reasoning.</p> | (6) |
| Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | | | | | | | | | | | | | | | | | | |
| 6 | 4 | | | | | | | | | | | | | | | | | | |
| 5-4 | 3 | | | | | | | | | | | | | | | | | | |
| 3-2 | 2 | | | | | | | | | | | | | | | | | | |
| 1 | 1 | | | | | | | | | | | | | | | | | | |
| 0 | 0 | | | | | | | | | | | | | | | | | | |
| | Number of marks awarded for structure and sustained lines of reasoning | | | | | | | | | | | | | | | | | | |
| Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning | 2 | | | | | | | | | | | | | | | | | | |

Edexcel Chemistry A-level - Periodicity

| | | | | |
|--|---|---|---|--|
| | <p>demonstrated throughout.</p> | | <p>If no reasoning mark(s) awarded do not deduct mark(s). Comment: Look for the indicative marking points first, then consider the mark for the structure of the answer and sustained line of reasoning.</p> | |
| | <p>Answer is partially structured with some linkages and lines of reasoning.</p> | 1 | | |
| | <p>Answer has no linkages between points and is unstructured.</p> | 0 | | |
| | <p>Indicative points:</p> <ul style="list-style-type: none"> • IP1: at the start of the period / on the LHS / Li to Be the bonding is metallic • IP2: metallic bonding gets stronger as the number of delocalised electrons in a metal (atom) increases or metallic bonding gets stronger as radius of cation decreases or metallic bonding gets stronger as the charge on the cation increases • IP3 in the middle of the period / (B and) C has / have a giant structure of atoms • IP4 A lot of energy is needed to break (strong) covalent bonds, (in graphite and diamond) | | <p>Ignore statements about boron</p> <p>Allow a description of a giant structure, e.g. each C atom is bonded to 4 other (in diamond)</p> | |
| | <ul style="list-style-type: none"> • IP5: at the end of the period / on the RHS / N to Ne are simple molecules or N₂, O₂ and F₂ are simple molecules, • IP6: weak London forces (between molecules) | | <p>Do not award London forces</p> <p>Ignore reference / lack of reference to Ne unless incorrect</p> | |

Edexcel Chemistry A-level - Periodicity

Q6.


| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|---|---|------------|
| (i) | <p>G above F AND H between G and F (1)</p> <p>I above H and below A AND J above I and below B (1)</p> <p>K below C (1)</p> | <p>Points which are not joined with lines are perfectly acceptable.</p> <p>Do not penalise I below G if MP1 not awarded</p> | (3) |

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|--|--|------------|
| (ii) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> D has one more proton / has a higher nuclear charge (1) the electron being removed in C and D are from the same subshell / s-subshell / (s) orbital (1) | <p>Allow same shell / energy level Allow the electron in D is closer to the nucleus than C / atomic radius decreases</p> <p>Ignore references to shielding, and full s-orbital which is more stable.</p> | (2) |

Edexcel Chemistry A-level - Periodicity

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|---|---|------------|
| (iii) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • (the electron being removed from E) is from a new subshell / p-subshell / p-orbital (1) • which is more shielded from the nucleus than the s-subshell (from which the electron is removed in D) <p>OR</p> <ul style="list-style-type: none"> • which is further from the nucleus than the s-subshell / orbital (in E) (1) | <p>Do not award 'in a new quantum shell' Allow electron removed from a higher energy level.</p> <p>Do not award clear reference to the outer electron in E being further from the nucleus than outer electron in D/atomic radius increasing from D to E</p> <p>Do not award clear reference to the outer electron in E being further from the nucleus than outer electron in D/atomic radius increasing from D to E</p> | (2) |

Q7.

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|---|---|------------|
| | <p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> • Workable method + time / compare (1) • Same heat applied (1) • Same amount of each nitrate in separate test tubes (1) • safety precaution: fume cupboard/hood (1) | <p><u>Examples of workable methods</u></p> <ul style="list-style-type: none"> • First one to re-light a glowing splint / produce brown fumes. Accurate timing not essential. • Use of light sensor / meter to measure colour of gas • Use of gas syringe and measure rate of production of gas / time to produce specific volume • Bubble gas into indicator solution – time to change colour • Collection of gases over water and volume measured <p>Reward any workable alternative. e.g. use the same Bunsen Award if implied by diagram</p> <p>Award 'equal masses'.</p> <p><u>Example diagram:</u></p>  <p>Ignore well ventilated room / face mask / goggles / gloves / lab coat This is the only acceptable safety precaution.</p> | <p>(4)</p> |

Edexcel Chemistry A-level - Periodicity

Q8.

| Question Number | Acceptable Answers | Additional Guidance | Mark | | | | | | | | | | | | |
|--|--|--|---|---|---|-----|---|-----|---|---|---|---|---|---|-------------------|
| | <p>This question assesses the student's ability to show a coherent and logically structured answer with linkages and fully sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="387 645 922 913"> <thead> <tr> <th data-bbox="387 645 655 763">Number of indicative marking points seen in answer</th> <th data-bbox="655 645 922 763">Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td data-bbox="387 763 655 792">6</td> <td data-bbox="655 763 922 792">4</td> </tr> <tr> <td data-bbox="387 792 655 822">5-4</td> <td data-bbox="655 792 922 822">3</td> </tr> <tr> <td data-bbox="387 822 655 851">3-2</td> <td data-bbox="655 822 922 851">2</td> </tr> <tr> <td data-bbox="387 851 655 880">1</td> <td data-bbox="655 851 922 880">1</td> </tr> <tr> <td data-bbox="387 880 655 909">0</td> <td data-bbox="655 880 922 909">0</td> </tr> </tbody> </table> | Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | 6 | 4 | 5-4 | 3 | 3-2 | 2 | 1 | 1 | 0 | 0 | <p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, a response with four indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there were no linkages between the points, then the same indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and zero marks for linkages).</p> | <p>(6)</p> |
| Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | | | | | | | | | | | | | | |
| 6 | 4 | | | | | | | | | | | | | | |
| 5-4 | 3 | | | | | | | | | | | | | | |
| 3-2 | 2 | | | | | | | | | | | | | | |
| 1 | 1 | | | | | | | | | | | | | | |
| 0 | 0 | | | | | | | | | | | | | | |

Edexcel Chemistry A-level - Periodicity

| | | | |
|--|--|--|--|
| | The following table shows how the marks should be awarded for structure and lines of reasoning | | <p>In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.</p> <p>Reasoning marks may be reduced for extra incorrect chemistry</p> |
| | | Number of marks awarded for structure of answer and sustained lines of reasoning | |
| | Answer shows a coherent logical structure with linkages and fully sustained lines of reasoning demonstrated throughout | 2 | |
| | Answer is partially structured with some linkages and lines of reasoning | 1 | |
| | Answer has no linkages between points and is unstructured | 0 | |

| | | | |
|--|--|--|--|
| | <p>Indicative content</p> <ul style="list-style-type: none"> the sum of the first two ionisation energies for barium is lower / barium loses (its outer) electrons more easily barium is a bigger atom/barium has a larger atomic radius/barium has more shells of electrons barium has more shielding these outweigh/exert a greater influence than barium has more proton/greater nuclear charge barium reacts faster/barium is more reactive | <p>Allow reverse argument for strontium</p> <p>Allow max 5 IPs for reference to general trends only down group 2</p> <p>Allow any reference to single ionisation</p> <p>Do not award for barium 'molecule'</p> | |
|--|--|--|--|

Q9.

| Question Number | Answer | Additional guidance | Mark |
|-----------------|---|---|------|
| | <p>A discussion that makes reference to the following points:</p> <ul style="list-style-type: none"> • both elements / atoms have the last added electron in the d-subshell / d orbital (so are d-block elements) (1) • but neither forms a (stable) ion with an incomplete d-subshell / d orbital (so are not transition metals) (1) • Zn^{2+} is $1s^22s^22p^63s^23p^63d^{10}$ (so d subshell is full) (1) • Sc^{3+} is $1s^22s^22p^63s^23p^6$ (so d subshell is empty) (1) | <p>Do not award just 'contains d electrons'</p> <p>Allow 'transition elements form a (stable) ion with an incomplete d-subshell / d orbital'</p> <p>Allow $[\text{Ar}]3d^{10}$</p> <p>Allow $[\text{Ar}]$</p> | (4) |

Edexcel Chemistry A-level - Periodicity

Q10.

| Question Number | Acceptable Answers | Additional Guidance | Mark | | | | | | | | | | | | |
|--|--|--|---|---|---|-----|---|-----|---|---|---|---|---|--|-----|
| * | <p>This question assesses a student's ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="379 779 751 1104"> <thead> <tr> <th data-bbox="379 779 552 954">Number of indicative marking points seen in answer</th> <th data-bbox="552 779 751 954">Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td data-bbox="379 954 552 987">6</td> <td data-bbox="552 954 751 987">4</td> </tr> <tr> <td data-bbox="379 987 552 1021">5-4</td> <td data-bbox="552 987 751 1021">3</td> </tr> <tr> <td data-bbox="379 1021 552 1055">3-2</td> <td data-bbox="552 1021 751 1055">2</td> </tr> <tr> <td data-bbox="379 1055 552 1088">1</td> <td data-bbox="552 1055 751 1088">1</td> </tr> <tr> <td data-bbox="379 1088 552 1122">0</td> <td data-bbox="552 1088 751 1122">0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> | Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | 6 | 4 | 5-4 | 3 | 3-2 | 2 | 1 | 1 | 0 | 0 | <p>Guidance on how the mark scheme should be applied: The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning). If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> <p>In general it would be expected that 5 or 6 indicative points would score 2 reasoning marks, and 3 or 4 indicative points would score 1 reasoning mark. A total of 2, 1 or 0 indicative points would score 0 marks for reasoning.</p> <p>Reasoning marks may be subtracted for extra incorrect chemistry.</p> | (6) |
| Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | | | | | | | | | | | | | | |
| 6 | 4 | | | | | | | | | | | | | | |
| 5-4 | 3 | | | | | | | | | | | | | | |
| 3-2 | 2 | | | | | | | | | | | | | | |
| 1 | 1 | | | | | | | | | | | | | | |
| 0 | 0 | | | | | | | | | | | | | | |

Edexcel Chemistry A-level - Periodicity

| | Number of marks awarded for structure of answer and sustained line of reasoning | |
|---|---|--|
| Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout. | 2 | |
| Answer is partially structured with some linkages and lines of reasoning. | 1 | |
| Answer has no linkages between points and is unstructured. | 0 | |

| | |
|---|--|
| <p>Indicative content (IPs)</p> <p>IP1:</p> <ul style="list-style-type: none"> (transition metal) forms an ion with an incomplete <i>d</i> sub-shell <p>IP2:</p> <ul style="list-style-type: none"> scandium and zinc are not transition metals <p>IP3:</p> <ul style="list-style-type: none"> Sc³⁺ and 1s² 2s² 2p⁶ 3s² 3p⁶ <p>IP4:</p> <ul style="list-style-type: none"> Zn²⁺ and 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ <p>IP5:</p> <ul style="list-style-type: none"> Sc³⁺ and <i>d</i> sub-shell empty / <i>d</i>-orbitals empty <p>IP6:</p> <ul style="list-style-type: none"> Zn²⁺ and <i>d</i> sub-shell full / ALL <i>d</i>-orbitals are full | <p>Allow 'partially-filled' for incomplete Allow <i>d</i>-orbital(s) Do not award "d-shell" Allow "D" for "d" throughout</p> <p>Allow if only Sc and Zn are used to illustrate <i>d</i>-block elements that are not transition metals</p> <p>Allow 4s⁰ and/or 3d⁰ Penalise use of [Ar] once only</p> <p>Allow "Sc³⁺ has no <i>d</i> sub-shell"</p> <p>Allow 'd orbital is full' if clarified by 3d¹⁰</p> |
|---|--|

Edexcel Chemistry A-level - Periodicity

Q11.

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|--|---|------|
| (i) | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> Al below Mg but above /equal to Na (1) rise from Al to Si and then to P and rise from S to Cl to Ar (1) S below P but above / equal to Si (1) | <p><u>Example of chart</u></p> <p>First ionisation energies of the Period 3 elements</p> <p>Allow use of dots (·) or other alternatives to X</p> <p>Ignore any lines connecting the crosses (X)</p> | (3) |

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|--|---|------|
| (ii) | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> big increase/jump between 1st and 2nd electrons removed and between 9th and 10th electrons removed (1) one / first electron in the outer most / third shell and eight electrons / electron 2 - 9 in the next / second shell and two electrons / electrons 10 & 11 in the inner most/ first shell (1) | <p>Allow answers in terms of energy levels</p> <p>Allow Na is a group 1 element</p> <p>Allow electronic configuration of Na is 2, 8, 1</p> <p>Allow an answer that relates jump in energy to existence of (new) shells</p> <p>Allow there are three shells of electrons</p> | (2) |

Q12.

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|--|---|------------|
| (i) | <p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> equation (1) state symbol, (g), on both H and H⁺ (1) | <p>$H(g) \rightarrow H^+(g) + e^{(-)}$</p> <p>or</p> <p>$H(g) - e^{(-)} \rightarrow H^+(g)$</p> <p>Ignore state symbol for electron</p> <p>$H_2(g) \rightarrow H_2^+(g) + e^{(-)}$ scores only M2</p> <p>$H_2(g) - e^{(-)} \rightarrow H_2^+(g)$ scores only M2</p> <p>$H_2(g) \rightarrow 2H^+(g) + 2e^{(-)}$ scores 0</p> <p>$X(g) \rightarrow X^+(g) + e^{(-)}$ scores only M2</p> | (2) |

| Question Number | Acceptable Answer | Additional Guidance | Mark |
|-----------------|---|---|------------|
| (ii) | <p>An explanation that makes reference to the following points: <u>H < He</u>:</p> <ul style="list-style-type: none"> He more protons than H / He greater nuclear charge than H (1) in helium the outer electron is in the same shell as hydrogen (1) <p><u>H > Li</u>:</p> <ul style="list-style-type: none"> in lithium the outer electron is in a higher energy level / a new shell / further from the nucleus / in a 2s orbital (1) (and) is shielded by inner electrons / 1s² electrons (1) | <p>Ignore references to shielding for H and He</p> <p>Ignore references to atomic radius or electrons being closer to or the same distance from the nucleus in helium</p> <p>Allow lithium has more shells of electrons</p> <p>Allow (outer electron of) lithium has more shielding than hydrogen / is shielded</p> | (4) |