

Q1. Fluorine and iodine are elements in Group 7 of the Periodic Table.

(a) Explain why iodine has a higher melting point than fluorine.

.....
.....
.....
(Extra space).....
.....

(2)

(b) (i) Draw the shape of the NHF_2 molecule and the shape of the BF_3 molecule.

Include any lone pairs of electrons that influence the shape. In each case name the shape.

Shape of NHF_2

Shape of BF_3

Name of shape of NHF_2

Name of shape of BF_3

(4)

(ii) Suggest a value for the F—N—F bond angle in NHF_2

.....


(1)

(c) State the strongest type of intermolecular force in a sample of NHF_2

.....

(1)

- (d) A molecule of NHF_2 reacts with a molecule of BF_3 as shown in the following equation.



State the type of bond formed between the N atom and the B atom in F_2HNBF_3 .

Explain how this bond is formed.

Name of type of bond

How bond is formed


.....

.....

(2)
(Total 10 marks)

Q2. Water can be found as ice, water and steam.

- (a) The following diagram shows the arrangement of some of the water molecules in a crystal of ice.



With reference to the structure shown above give **one** reason why ice is less dense than water.

.....
.....
.....

(1)

(b) Water and methane have similar relative molecular masses and both contain the element hydrogen.

The table below gives some information about water and methane.

	H ₂ O	CH ₄
<i>M_r</i>	18.0	16.0
Melting point / K	273	91

(i) State the strongest type of intermolecular force holding the water molecules together in the ice crystal.

.....

(1)

(ii) State the strongest type of intermolecular force in methane.

.....

(1)

(iii) Give **one** reason why the melting point of ice is higher than the melting point of methane.

.....
.....
.....

(1)

(c) A molecule of H₂O can react with an H⁺ ion to form an H₃O⁺ ion.

- (i) Draw and name the shape of the H_3O^+ ion. Include any lone pairs of electrons.

Shape of the H_3O^+ ion

Name of shape

(2)

- (ii) Suggest a value for the bond angle in the H_3O^+ ion.

.....

(1)

- (iii) Identify **one** molecule with the same number of atoms, the same number of electrons and the same shape as the H_3O^+ ion.

.....

(1)



- (d) Water can also form the hydroxide ion.
State the number of lone pairs of electrons in the hydroxide ion.

.....

(1)

(Total 9 marks)

- Q3.** (a) Two organic compounds with similar relative molecular masses are shown below.



Ethanol

Propane

- (i) State the type of bond present between the C and H atoms in both of these molecules. Explain how this type of bond is formed.

Type of bond

Explanation

(2)


- (ii) State the strongest type of intermolecular force present in each compound.

Liquid ethanol

Liquid propane

(2)

- (b) Ethanol dissolves in water. Draw a diagram to show how one molecule of ethanol interacts with one molecule of water in the solution. Include partial charges and all lone pairs. The ethanol molecule has been drawn for you.



(3)

- (c) Ethanol was the fuel used in the first mass-produced car, the Model T Ford.

- (i) Write an equation which shows how ethanol burns completely in air to form carbon dioxide and water as the only products.

.....

(1)

- (ii) Suggest **one** environmental problem caused by incomplete combustion of ethanol in a car engine.

.....

.....

(1)

- (iii) Suggest **one** economic problem for the car user caused by incomplete combustion of ethanol in the car engine.

.....

.....

(1)

- (d) Propane is also used as a fuel, although sometimes it can be contaminated with sulfur-containing impurities. When this propane burns, these impurities form sulfur dioxide.

- (i) State how the sulfur dioxide can be removed from the waste gases produced when this propane is burned on a large scale in industry. Suggest a reason why the method you have stated may not be 100% efficient.

How removed

.....

Reason for less than 100% efficiency

.....

(2)

- (ii) Although propane has a boiling point of $-42\text{ }^{\circ}\text{C}$, it is usually supplied as a liquid for use in camping stoves. Suggest why it is supplied as a liquid.

.....

.....

(1)

(Total 13 marks)

Q4. The table below shows the boiling points of some hydrogen compounds formed by Group 6 elements.

	H ₂ O	H ₂ S	H ₂ Se	H ₂ Te
Boiling point / K	373	212	232	271

- (a) State the strongest type of intermolecular force in water and in hydrogen sulfide (H₂S).

Water

Hydrogen sulfide

.....

(2)

- (b) Draw a diagram to show how two molecules of water are attracted to each other by the type of intermolecular force you stated in part (a). Include partial charges and all lone pairs of electrons in your diagram.

(3)

- (c) Explain why the boiling point of water is much higher than the boiling point of hydrogen sulfide.

.....

.....

(1)

- (d) Explain why the boiling points increase from H₂S to H₂Te

.....

.....

(2)

- (e) When H⁺ ions react with H₂O molecules, H₃O⁺ ions are formed.

Name the type of bond formed when H^+ ions react with H_2O molecules.
Explain how this type of bond is formed in the H_3O^+ ion.

Type of bond

Explanation

.....

(2)

- (f) Sodium sulfide (Na_2S) has a melting point of 1223 K.
Predict the type of bonding in sodium sulfide and explain why its melting point is high.

Type of bonding

Explanation

.....

.....

(3)

(Total 13 marks)

Q5. Fluorine forms many compounds that contain covalent bonds.

- (a) (i) State the meaning of the term *covalent bond*.

.....

.....

(1)

- (ii) Write an equation to show the formation of one molecule of ClF_3 from chlorine and fluorine molecules.

.....

(1)

- (b) Draw the shape of a dichlorodifluoromethane molecule (CCl_2F_2) and the shape of a

chlorine trifluoride molecule (ClF₃). Include any lone pairs of electrons that influence the shape.

Shape of CCl₂F₂

Shape of ClF₃

(2)

(c) Suggest the strongest type of intermolecular force between CCl₂F₂ molecules.

.....

(1)

(d) BF₃ is a covalent molecule that reacts with an F⁻ ion to form a BF₄⁻ ion.

(i) Name the type of bond formed when a molecule of BF₃ reacts with an F⁻ ion. Explain how this bond is formed.

Type of bond

Explanation

.....

.....

.....


(3)

(ii) State the bond angle in the BF₄⁻ ion

.....

(1)

(e) An ultrasound imaging agent has the formula C₄F₁₀. It can be made by the reaction of butane and fluorine as shown in the following equation.



Calculate the percentage atom economy for the formation of C_4F_{10} in this reaction.
Give your answer to three significant figures.

.....
.....

(2)
(Total 11 marks)