Q1. (a)	Nickel is a r	Nickel is a metal with a high melting point.					
	(i)	State the block in the Periodic Table that contains nickel.					
			(1)				
	(ii)	Explain, in terms of its structure and bonding, why nickel has a high melting point.					
			(2)				
	(iii)	Draw a labelled diagram to show the arrangement of particles in a crystal of nickel. In your answer, include at least six particles of each type.					
			(2)				
	(iv)	Explain why nickel is ductile (can be stretched into wires).					
			(1)				

(b) Nickel forms the compound nickel(II) chloride (NiCl $_2$).

	(i)	Give the full electron configuration of the Ni ²⁺ ion.	
			(1)
	(ii)	Balance the following equation to show how anhydrous nickel(II) chloride can be obtained from the hydrated salt using SOCl ₂ Identify one substance that could react with both gaseous products.	I
	1	$NiCl_2.6H_2O(s) +SOcl_2(g) \longrightarrowNiCl_2(s) +SO_2(g) +HCl(g)$	
		Substance(Total 9	(2) 9 marks)
Q2. This qu	ıestioı	n is about some Period 3 elements and their oxides.	
(a)	heat	scribe what you would observe when, in the absence of air, magnesium is ted strongly with water vapour at temperatures above 373 K. e an equation for the reaction that occurs.	
	Obse	ervations	
	Equa	ation	(3)
(b)	Exp	lain why magnesium has a higher melting point than sodium.	
	(Fxtr	ra space)	

		(2)
(c)	State the structure of, and bonding in, silicon dioxide. Other than a high melting point, give two physical properties of silicon dioxide that are characteristic of its structure and bonding.	
	Structure	
	Bonding	
	Physical property 1	
	Physical property 2	(4)
(d)	Give the formula of the species in a sample of solid phosphorus(V) oxide. State the structure of, and describe fully the bonding in, this oxide.	
	Formula	
	Structure	
	Bonding	
		(4)
(e)	Sulfur(IV) oxide reacts with water to form a solution containing ions.	
	Write an equation for this reaction.	
		(1)
(f)	Write an equation for the reaction between the acidic oxide, phosphorus(V) oxide, and the basic oxide, magnesium oxide.	
	(Total 15	(1) marks)

Q3.	 Group 2 metals and their compounds are used commercially in a variety of proces and applications. 				
	(a)	Stat	e a use of magnesium hydroxide in medicine.		
				(1)	
	(b)		cium carbonate is an insoluble solid that can be used in a reaction to lower the ity of the water in a lake.		
		Expl the la	ain why the rate of this reaction decreases when the temperature of the water in ake falls.		
				(3)	
	(c)	Stro	ntium metal is used in the manufacture of alloys		
	(c)	(i)	ntium metal is used in the manufacture of alloys. Explain why strontium has a higher melting point than barium.		
		()			
				(2)	

Write an equation for the reaction of strontium with water.

(ii)

				(1)
	(d)	Mag	nesium can be used in the extraction of titanium.	
		(i)	Write an equation for the reaction of magnesium with titanium(IV) chloride.	
				(1)
		(ii)	The excess of magnesium used in this extraction can be removed by reacting it with dilute sulfuric acid to form magnesium sulfate.	
			Use your knowledge of Group 2 sulfates to explain why the magnesium sulfate formed is easy to separate from the titanium.	
			(Total 9 ma	(1) rks
Q4.	com	There pound:	are several types of crystal structure and bonding shown by elements and s.	
	(a)	(i)	Name the type of bonding in the element sodium.	
				(1)
		(ii)	Use your knowledge of structure and bonding to draw a diagram that shows how the particles are arranged in a crystal of sodium. You should identify the particles and show a minimum of six particles in a two-dimensional diagram.	

,	
1	_

(b)	Sodi	um reacts wit	h chlorine t	to form sodium ch	loride.			
	(i)	Name the ty	Name the type of bonding in sodium chloride.					
								(1
	(ii)	Explain why	the melting	g point of sodium	chloride is hig	ıh.		
		(Extra space	•)					
								(2
(-)	The	tabla balawal	41			- 1: -1		
(c)	rne	table below si	nows the m	nelting points of so	me soaium r	ialides.	ı	
		Na	aCl	NaBr	Na	ıl		
elting	point /	K 10	74	1020	92	0		
	Suggest why the melting point of sodium iodide is lower than the melting point of sodium bromide.				ing point of			
							(Total	(1 7 marks

Q5. Group 2 metals and their compounds are used commercially in a variety of processes.

(a) Strontium is extracted from strontium oxide (SrO) by heating a mixture of powdered strontium oxide and powdered aluminium.

Consider these standard enthalpies of formation.

	SrO(s)	Al ₂ O ₃ (s)
ΔH _i ∘ / kJ mol⁻¹	– 590	– 1669

$$3SrO(s) + 2AI(s) \longrightarrow 3Sr(s) + AI2O3(s)$$

Use these data and the equation to calculate the standard enthalpy change for this extraction of strontium.

The use of powdered strontium oxide and powdered aluminium increases the surface area of the reactants.

Suggest **one** reason why this increases the reaction rate.

Suggest one major reason why this method of extracting strontium is expensive.

(5)

(b) Explain why calcium has a higher melting point than strontium.

		(2)
(c)	Magnesium is used in fireworks. It reacts rapidly with oxygen, burning with a white light. Magnesium reacts slowly with cold water.	bright
	Write an equation for the reaction of magnesium with oxygen.	
	Write an equation for the reaction of magnesium with cold water.	
	Give a medical use for the magnesium compound formed in the reaction of magnesium with cold water.	
	т	(3) otal 10 marks)

Q6.Group 2 elements and their compounds have a wide range of uses.

(a) For parts (a)(i) to (a)(iii), draw a ring around the correct answer to complete each sentence.

decreases.

	(i) From Mg(OH) ₂ to Ba(OH) ₂ , the solubility in water	increases. stays the same.	(1)
	(ii) From Mg to Ba, the first ionisation energy	decreases. increases. stays the same.	(1)
	(iii) From Mg to Ba, the atomic radius	decreases. increases. stays the same.	(1)
(b)	Explain why calcium has a higher melting point than stro	ontium.	(1)
	(Extra space)		(2)
(c)	Acidified barium chloride solution is used as a reagent to	o test for sulfate ions.	(2)

State why sulfuric acid should ${f not}$ be used to acidify the barium chloride.

(i)

		(1)
(ii)	Write the simplest ionic equation for the reaction that occurs when acidified barium chloride solution is added to a solution containing sulfate ions.	
		(1)
	(Total 7 ma	arks)