

A-Level Chemistry

Organic Analysis (Multiple Choice)

Question Paper

Time available: 12 minutes Marks available: 11 marks

www.accesstuition.com



1.



Which compound produces this spectrum?



2. Which can be used to distinguish between these two compounds?

(CH₃)₂CHCH₂CHO and (CH₃)₃CCHO



(Total 1 mark)

Three reagents are added separately to four organic compounds.

Which row shows the correct observations?

		Sodium hydrogen carbonate	Acidified potassium dichromate(VI)	Tollens' reagent	
A	Propan-1-ol	effervescence	orange solution turns green	no visible change	0
в	Propanal	no visible change	orange solution turns green	silver mirror	0
С	Propanone	no visible change	no visible change	silver mirror	0
D	Propanoic acid	effervescence	no visible change	silver mirror	0

(Total 1 mark)

3.



Which compound produces this spectrum?

Α	butanone	0
в	ethanol	0
С	pent-2-ene	0
D	propanoic acid	0

5.

(Total 1 mark)

Which compound forms a molecular ion with a different precise molecular mass from the other three?



(Total 1 mark)

6.





www.accesstuition.com



(Total 1 mark)

8. Which of the following compounds would form an orange-red precipitate when heated with Fehling's solution?

Α	CH ₃ CH ₂ CN	0
в	CH ₃ CH ₂ COOH	0
С	CH₃CHO	0
D	CH ₃ COCH ₃	0

(Total 1 mark)

9.

This question is about the reaction between propanone and an excess of ethane-1,2-diol, the equation for which is given below.

$$CH_3COCH_3 + HOCH_2CH_2OH \implies (CH_3)_2 C \swarrow \begin{matrix} O - CH_2 \\ I \\ O - CH_2 \end{matrix} + H_2O$$

In a typical procedure, a mixture of 1.00 g of propanone, 5.00 g of ethane-1,2-diol and 0.100 g of benzenesulphonic acid, $C_6H_5SO_3H$, is heated under reflux in an inert solvent. Benzenesulphonic acid is a strong acid.

The products would not have an absorption in the infra-red at

- A 1050 cm⁻¹
- **B** 1720 cm⁻¹
- **C** 2950 cm⁻¹
- **D** 3400 cm⁻¹



A It has stereoisomers.

11.

- **B** It shows a strong absorption in the infra-red at about 1700 cm⁻¹.
- **C** It will turn an acidified solution of potassium dichromate(VI) green.
- **D** It can be dehydrated by concentrated sulphuric acid.

(Total 1 mark)

Certain chemical tests were performed on the pain-relief drug ibuprofen. The results of these tests are given in the table below.

Test	Result
Aqueous sodium carbonate	Effervescence
Bromine water	Remained orange
Acidified potassium dichromate(VI) and heat	Remained orange
Fehling's solution and heat	Remained blue

Which one of the following functional groups do these results suggest that ibuprofen contains?



(Total 1 mark)