



1 Chemists often use two different structures to represent a molecule of benzene, as shown below.



structure A

below.



structure B

(a) (i) Describe, with the aid of suitable diagrams showing orbital overlap, the difference in bonding between structure **A** and structure **B**.



In your answer, you should use appropriate technical terms, spelled correctly.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[4]

- (ii) The table below shows the enthalpy changes for the reactions of cyclohexene, C_6H_{10} , and benzene, C_6H_6 , with hydrogen.

reaction	enthalpy change/ kJ mol^{-1}
$C_6H_{10} + H_2 \rightarrow C_6H_{12}$	-119
$C_6H_6 + 3H_2 \rightarrow C_6H_{12}$	-208


Using this information, suggest and explain whether structure **A** or structure **B** is a better representation of benzene.

.....


 [2]

- (b) Benzene compounds can undergo nucleophilic substitution reactions.

Add curly arrows to the diagram below to show the two-step mechanism of $C_6H_5N_2^+$ with F^- .



- (c) Benzene can react with halogenoalkanes in the same way as with bromine, as shown in **reaction 1** below.




Write an equation to show the formation of the electrophile that reacts with benzene in **reaction 1**.


(d) The types of reaction in (b) and (c) can be used to synthesise compound **D**, as shown in the flowchart below.

(i) Complete the boxes below to suggest formulae for the reactants involved in the synthesis of compound **D**.


Give structures for organic compounds.



(ii) In a synthesis of compound **D** from 1,3-diaminobenzene shown in the flowchart, 1.73 g of compound **D** was prepared. These structures have been repeated below:



1,3-diaminobenzene



compound D

The overall percentage yield of compound **D** was 40.0%.

M_r of compound **D** = 346.0

Calculate the mass of 1,3-diaminobenzene needed for this synthesis.

mass = g [3]


(iii) Compound **D** has been developed for possible use as a drug to treat heart conditions. When compound **D**, prepared in this synthesis, was given to patients, only 25% of the dose was effective in treating their heart conditions.

Explain why only 25% of the dose was effective. Suggest how the synthesis of compound **D** might be changed to make the dose more effective.

.....
.....
.....
.....
..... [3]

[Total: 18]

2 4-Aminophenol is an organic compound that can behave as an acid and a base.



4-aminophenol

(a) State how 4-aminophenol can behave as a base.

.....
..... [1]


(b) 4-Aminophenol is produced by the reduction of 4-nitrophenol.

Write an equation to show the production of 4-aminophenol from 4-nitrophenol.
Use [H] to represent the reducing agent.

[1]


(c) 4-Nitrophenol can be produced from 4-bromophenol.

- Complete the mechanism for this reaction.
- Use $^+\text{NO}_2$ as the electrophile. Include any intermediate and the products.
- In the mechanism for this reaction, NO_2 substitutes for Br on the ring.



[4]

(d) The flowchart below shows some reactions of 4-aminophenol.



(i) Identify the reagent in **reaction I**.

..... [1]

(ii) Name the organic product of **reaction II**.

..... [1]


(iii) Write the equation for **reaction II**.

[1]


(iv) In the box on the flowchart, draw the structure of the organic compound formed by **reaction III**. [1]

- (e) The Sandmeyer reaction can be used to replace a diazonium group, N_2^+ , with a halogen atom, **X**, on an aromatic ring.

The reagent used for the reaction is a copper(I) halide, CuX .



Compound **C**, shown below, can be synthesised using **only** 4-aminophenol and other standard laboratory reagents. The flowchart on the next page shows this synthesis.



compound C


- (i) State a possible use for compound **C**.

..... [1]

- (ii) On the flowchart on the next page:

- state the reagents and conditions used for **reaction 1**
- suggest the structure of compound **B**
- suggest the reagent used for **reaction 2**
- state the conditions used for **reaction 3**.

[5]




4-aminophenol

reaction 1

reagents

conditions



compound A


reaction 2

reagent

compound B

reaction 3

conditions



compound C

[Total: 16]

3 A student was investigating the reactions and uses of organic amines.

(a) The student found that amines such as ethylamine, $C_2H_5NH_2$, and phenylamine, $C_6H_5NH_2$, both behave as bases.

(i) Explain why amines can behave as bases.

.....
..... [1]


(ii) The student reacted an excess of $C_2H_5NH_2$ with two different acids.

Write the formulae of the salts that would be formed when an **excess** of $C_2H_5NH_2$ reacts with:

sulfuric acid,

ethanoic acid. [2]

(b) The student reacted phenylamine with a mixture of $NaNO_2(aq)$ and $HCl(aq)$ whilst keeping the temperature below $10^\circ C$. A diazonium ion was formed. The student then reacted the diazonium ion with compound **B**. After neutralisation, compound **A** was formed.



compound **A**

(i) Draw the structures of the diazonium **ion** and compound **B**.

Display the functional group in the diazonium ion.

diazonium ion	compound B
----------------------	-------------------

[2]

- (ii) State the conditions required for the reaction of the diazonium ion with compound **B** and state a possible use for compound **A**.

conditions

possible use for compound **A**. [1]

- (iii) The student added Na_2CO_3 to a solution of compound **A**.

Draw the structure of the organic product and state the formulae of any other products from this reaction.

[2]

- (c) The student repeated the experiment in part (b) but allowed the temperature to rise above 10°C .

Under these conditions, the diazonium ion in (b)(i) reacts with water to produce phenol. A gas with molar mass of 28.0 g mol^{-1} and one other product are also formed.

Construct an equation for this reaction.


[1]

[Total: 9]

4 Hydroxyamines are organic compounds containing hydroxyl and amino functional groups.

(a) Salbutamol is a hydroxyamine used in the treatment of asthma and bronchitis. Salbutamol is an example of a chiral drug.

(i) Draw a circle around the chiral carbon in the structure of salbutamol shown below.



salbutamol

[1]


(ii) Suggest possible problems of making a chiral drug such as salbutamol and describe **two** ways that the pharmaceutical industry might overcome these problems.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

[4]

- (b) Monoethanolamine, MEA, $\text{H}_2\text{NCH}_2\text{CH}_2\text{OH}$, is a hydroxyamine that is used in aqueous solution as a gas scrubber to remove acidic gases from emissions in incinerators.

MEA is prepared industrially by reacting ammonia with epoxyethane.



epoxyethane

- (i) Write an equation for the industrial preparation of MEA.

[1]

- (ii) During the manufacture of MEA, a compound with molecular formula $\text{C}_4\text{H}_{11}\text{NO}_2$ is also formed.

Draw the structure of the compound with molecular formula $\text{C}_4\text{H}_{11}\text{NO}_2$.

[1]

- (c) The combustion of some polymers produces emissions containing toxic acidic gases such as HCl and H_2S . MEA can remove HCl and H_2S from the emissions.

Give the formula of the organic salts formed when MEA removes:

- (i) HCl ,

[1]

- (ii) H_2S .

[1]

(d) MEA, $\text{H}_2\text{NCH}_2\text{CH}_2\text{OH}$, can be oxidised to form an α -amino acid.

(i) Explain what is meant by an α -amino acid.


.....
.....
..... [1]

(ii) Write an equation for the oxidation of MEA to form an α -amino acid.

Use [O] to represent the oxidising agent.

..... [1]

(e) Isomers **F** and **G** are hydroxyamines each with the molecular formula $\text{C}_4\text{H}_{11}\text{NO}$.

- Isomer **F** can be dehydrated to form the cyclic compound 
- Isomer **G** has two chiral centres.

Identify and draw the structural isomers **F** and **G**.

isomer F	isomer G
-----------------	-----------------

[2]

[Total: 13]