

1. (a) (i) Calculate correctly $\frac{0.0880 \times 25.0}{1000} = 2.20 \times 10^{-3} \text{ mol}$

OR 0.00220 mol ✓

ALLOW 0.0022 **OR** $2.2 \times 10^{-3} \text{ mol}$

1

(ii) Calculates correctly $\frac{0.00220}{2} = 1.10 \times 10^{-3} \text{ mol}$

OR 0.00110 mol ✓

ALLOW 0.0011 **OR** $1.1 \times 10^{-3} \text{ mol}$

ALLOW ECF for answer (i)/2 as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes

1

(iii) $\frac{0.00110 \times 1000}{17.60} = 0.0625 \text{ mol dm}^{-3}$

OR $6.25 \times 10^{-2} \text{ mol dm}^{-3}$ ✓

ALLOW 0.063 **OR** $6.3 \times 10^{-2} \text{ mol dm}^{-3}$

ALLOW ECF for answer (ii) $\times 1000/17.60$

OR

ECF from (i) for answer (i)/2 $\times 1000/17.60$ as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes

1

(b) (i) (The number of) Water(s) of crystallisation ✓

IGNORE hydrated **OR** hydrous

1

(ii) 142.1 ✓

ALLOW 142

ALLOW M_r expressed as a sum

ALLOW ECF from incorrect M_r and x is **calculated correctly**

$$x = \frac{(322.1 - 142.1)}{18.0} = 10 \text{ ✓}$$

ALLOW ECF values of x from nearest whole number to calculator value

ALLOW 2 marks if final answer is 10 **without any working**

2

[6]

2. (i) O goes from -2 to 0 ✓

Oxidation numbers may be seen with equation

N goes from +5 to +4 ✓

N is reduced **AND** O is oxidised ✓

Third mark is dependent upon seeing a reduction in oxidation number of N and an increase in oxidation number of O

***ALLOW** ECF for third mark for N is oxidised **and** O is reduced if incorrect oxidation numbers support this*

***IGNORE** references to strontium*

***IGNORE** references to electron loss **OR** gain*

***DO NOT ALLOW** 'One increases and one decreases'*

3

- (ii) Calculates correctly:

$$\text{Mol of Sr(NO}_3)_2 = \frac{5.29}{211.6} = 0.0250 \quad \checkmark$$

***ALLOW** 0.025*

Calculates correctly:

$$\text{Mol of gas} = 5/2 \times 0.0250 = 0.0625 \quad \checkmark$$

***ALLOW** ECF for first answer $\times 2.5$ as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes*

Calculates correctly:

$$\text{Volume of gas} = 24.0 \times 0.0625 = 1.50 \text{ dm}^3 \quad \checkmark$$

***ALLOW** ECF for second answer $\times 24(.0)$ as calculator value or correct rounding to 2 significant figures or more but ignore trailing zeroes*

***DO NOT ALLOW** ECF of first answer $\times 24(.0)$ (which gives $0.6(0) \text{ dm}^3$) as this has not measured the volume of any gas, simply 0.0250 mol of solid $\text{Sr(NO}_3)_2$ converted into a gas*

*i.e. This answer would give **one** mark*

***ALLOW** 1.5 dm^3*

***ALLOW** ECF producing correct volume of NO_2 only*

*i.e. $1.2(0) \text{ dm}^3$ would give **two** marks*

OR

***ALLOW** ECF producing correct volume of O_2 only*

*i.e. $0.3(0) \text{ dm}^3$ would give **two** marks*

3

[6]

3. (i) 0.0268 **OR** 0.027 **OR** 0.02675 mol ✓
NO OTHER ACCEPTABLE ANSWER 1

(ii) 1.61×10^{22} ✓
ALLOW 1.6×10^{22} up to calculator value
ALLOW
ECF answer to (i) $\times 6.02 \times 10^{23}$
ALLOW any value for N_A in the range:
 $6.0 \times 10^{23} - 6.1 \times 10^{23}$ 1

[2]

4. (a) BaO ✓
Ba₃N₂ ✓
Treat any shown charges as working and ignore.
Treat B for Ba as a slip 2

(b) (i) $\frac{0.11}{137.3}$ ✓
mark is for the working out which MUST lead to the correct answer of 8×10^{-4} up to calculator value 1

(ii) 19.2
OR
calculated answer to (b)(i) $\times 24000$ ✓
ALLOW 19 up to calculator value. 1

(iii) 8.0×10^{-3}
OR
calculated answer to (b)(i) $\times 10$ ✓
ALLOW 8.01×10^{-3} up to calculator value 1

(iv) any pH > 7 but < 15 ✓
ALLOW a correct range of pH. 1

(c) Less barium to react **OR**

some barium has already reacted ✓
ALLOW less volume because contains some BaO or Ba₃N₂

1

- (d) reactivity increases (down the group) ✓
atomic radii increase **OR**
there are more shells ✓
there is **more** shielding **OR more** screening ✓
the nuclear attraction decreases **OR**
Increased shielding and distance outweigh the
increased nuclear charge ✓
easier to remove (outer) electrons **OR**
ionisation energy decreases ✓

*USE annotations with ticks, crosses, ecf, etc for
this part.*

DO NOT ALLOW more orbitals OR more sub-shells

'More' is essential

ALLOW 'more electron repulsion from inner shells'

ALLOW 'nuclear pull'

IGNORE any reference to 'effective nuclear charge'

ALLOW easier to form positive ion

5

[12]

5. (i) mol HCl = 1.50×10^{-2} ✓

volume HCl(aq) = 75.0 ✓

ALLOW answers to 2 significant figures

ALLOW ecf from wrong number of moles

i.e. $\frac{\text{moles of HCl} \times 1000}{0.200}$

ALLOW one mark for 37.5 (from incorrect 1:1 ratio)

2

- (ii) 180 ✓

No other acceptable answer

1

[3]

6. (i) Molar mass of $\text{CaCO}_3 = 100.1 \text{ g mol}^{-1}$ (1) 2
 $2.68/100.1 = 0.0268/0.027$ (1)
- (ii) $0.0268 \text{ mol} \times 24,000 = 643 \text{ cm}^3$ (1) 1
- (iii) moles $\text{HNO}_3 = 2 \times 0.0268$
 $= 0.0536 / 0.054 \text{ mol}$ (1)
(i.e. answer to (i) $\times 2$)
- volume of $\text{HNO}_3 = 0.0536 \times 1000/2.50 = 21.4 \text{ cm}^3$ (1) 2
- [5]**
7. (i) **Simplest (whole number) ratio** of atoms/moles/elements ✓ 1
- (ii) ratio Rb : Ag : I = $7.42/85.5 : 37.48/108 : 55.10/127$
or $0.0868 : 0.347 : 0.434$
or $1 : 4 : 5$ ✓
= RbAg_4I_5 ✓ 2
- [3]**
8. (a) (i) $12 \times 50/1000 = 0.600 \text{ mol}$ ✓ 1
- (ii) $4 \text{ mol HCl} \rightarrow 1 \text{ mol Cl}_2$ / moles $\text{Cl}_2 = 0.15 \text{ mol}$ ✓
vol of $\text{Cl}_2 = 0.15 \times 24 = 3.60 \text{ dm}^3$ ✓ 2
2nd mark is consequential on molar ratio given
- (b) Evidence that the oxidation number of Mn has reduced
and one of the oxidation numbers correct (ie MnO_2 : ox no
of Mn = +4 or MnCl_2 : ox no of Mn = +2 ✓
The **other** oxidation number of Mn is correct,
ie in MnO_2 : ox no of Mn = +4
or in MnCl_2 : ox no of Mn = +2 ✓ 2
- [5]**

9. (i) $\text{mass} = 0.0500 \times 23.0 = 1.15 \text{ g}$ ✓ 1
- (ii) $\text{moles H}_2 = 0.0250$ ✓
 $\text{volume H}_2 = 0.0250 \times 24 = 0.600 \text{ dm}^3$ ✓ 2
 ecf from calculated moles H₂
 $0.0500 \text{ mol in } 50.0 \text{ cm}^3$ 1
- (iii) $\text{concentration} = 0.0500 \times 20 = 1.00 \text{ mol dm}^{-3}$ ✓
- [4]

10. (i) $2\text{Na} + \text{O}_2 \rightarrow \text{Na}_2\text{O}_2$ ✓ 1
- (ii) $\text{Na}_2\text{O}_2 + 2\text{H}_2\text{O} \rightarrow \text{H}_2\text{O}_2 + 2\text{NaOH}$ ✓ 1
 correct covalent bonds shown ✓
- (iii) electron count (14) for rest of molecule correct ✓ 2
- [4]

11. $M(\text{BaO}) = 137 + 16 = 153$ ✓
 $\text{moles BaO} = 500/153$ or 3.268 mol ✓
 $\text{moles Ba} = 3.268/2$ or 1.634 ✓
 $\text{mass Ba formed} = 1.634 \times 137 = 224 \text{ g}$ ✓
 accept 223.856209/223.86/223.9 g.
 if 6 mol BaO forms 3 mol Ba, award 3rd mark
- Alternative method
 $\text{mass } 6\text{BaO} = 918 \text{ g}$ ✓
 $\text{mass } 3\text{Ba} = 411 \text{ g}$ ✓
 1g BaO forms 411/918 g Ba ✓
 500 g BaO forms 223.856209/223.86/223.9 g Ba ✓
- [4]

12. (i) $\text{ratio N : H : S : O} = \frac{24.12}{14} : \frac{6.94}{1} : \frac{27.61}{32.1} : \frac{41.33}{16} : \checkmark$
 $= 2 : 8 : 1 : 3$
 Empirical formula = $\text{N}_2\text{H}_8\text{SO}_3$ ✓
 $\text{N}_2\text{H}_4\text{SO}_3$ is worth 1 mark from consistent use of at nos. 2
- (ii) $\text{H}_2\text{O} + 2\text{NH}_3 + \text{SO}_2 \rightarrow (\text{NH}_4)_2\text{SO}_3$ ✓ 1
(Award mark for N₂H₈SO₃)
- [3]

13. (a) (i) Amount of substance that has the same number of particles as there are atoms in 12 g of ^{12}C /
 6×10^{23} / Avogadro's Number ✓ 1
- (ii) moles = $\frac{0.275 \times 120}{1000} = 0.0330 \text{ mol}$ ✓
 moles $\text{Cl}_2 = \frac{0.0330}{2} = 0.0165 \text{ mol}$ ✓ 1
- (iii) volume $\text{Cl}_2 = 0.0165 \times 24000 = 396 \text{ cm}^3$ ✓ / 0.396 dm^3
 792 cm^3 worth 1 mark (no molar ratio)
 1584 cm^3 worth 1 mark (x 2)
 units needed. 2
- (iv) bleach / disinfectant /sterilising /killing germs ✓ 1
- (b) NaClO_3 ✓ 1

[6]

14. (i) Mass Sb_2S_3 in stibnite = 5% of 500 kg = 25.0 kg ✓
 Moles $\text{Sb}_2\text{S}_3 = \frac{25.0 \times 10^3}{340} / 73.5 / 73.529 / 73.53 / 74 \text{ mol}$ ✓
 (calculator value: 73.52941176)
 If 5% is not used, 1471 mol; ecf for 2nd mark
 (calculator value: 1470.588235)
 If 5% is used 2nd, 73.6 mol: OK for both marks 2
- (ii) moles Sb = $2 \times 73.5 \text{ mol}$ ✓
 mass Sb = $2 \times 73.5 \times 122 \text{ g} = 17.9 \text{ kg}$ ✓
 If the 2 isn't used, answer = $73.5 \times 122 = 8.95$ ✓
ecf ans from (i) x 2
ecf ans above x 2
- OR**
 % Sb = $244/340 = 71.7\%$ ✓
 mass Sb = $25.0 \times 71.7/100 = 17.9 \text{ kg}$ ✓ (ecf as above) 2

[4]

15. (i) Molar mass CaO = 56.1 (g mol⁻¹) ✓ (anywhere) 2
 moles CaO = $\frac{1.50}{56.1} = 0.0267/0.027$ ✓ calc: 0.0267379
 Allow 56 which gives 0.0268
- (ii) moles HNO₃ = 2 × 0.0267
 = 0.0534 or 0.0535 /0.053 mol ✓
(i.e. answer to (i) x 2)
 volume of HNO₃ = $\frac{0.0534 \text{ (or 5)} \times 1000}{2.50} = 21.4 \text{ cm}^3$ ✓ 2
 calc from value above = 21.3903743
 If 0.053 mol, answer is 21 cm³ but accept 21.2 cm³
 If 0.054 mol, answer is 22 cm³ but accept 21.6 cm³

[4]

16. (i) dative covalent, bonded pair comes from same atom/
 electron pair is donated from one atom/
 both electrons are from the same atom ✓ 1
- (ii) Ca(NO₃)₂ ✓ → CaO + 2NO₂ + ½O₂ ✓
or double equation with 2/2/4/1 1

[2]

17. (i) 203.3 g mol⁻¹ ✓
 Accept 203 1
- (ii) white precipitate / goes white ✓ 1
- (iii) Ag⁺(aq) + Cl⁻(aq) → AgCl(s)
 equation ✓
 state symbols ✓ 2
 AgCl dissolves in NH₃(aq) ✓
- (iv) AgBr dissolves in **conc** NH₃(aq)/
 partially soluble in NH₃(aq) ✓
 AgI insoluble in NH₃(aq) ✓ 3

[7]

18. (i) moles $\text{CO}_2 = 1000 / 44 \text{ mol} = 22.7 \text{ mol} \checkmark$
 volume CO_2 in 2000 = $22.7 \times 24 = 545 \text{ dm}^3 \checkmark$
- (ii) reduction = $545 \times 60/100 = 327 \text{ dm}^3 \checkmark$
- [3]**
-
19. (i) moles $\text{HCl} = 2.0 \times 50/1000 = 0.10 \checkmark$ 1
- (ii) moles $\text{Ca} = \frac{1}{2} \times \text{moles HCl} = 0.050 \checkmark$
 mass $\text{Ca} = 40.1 \times 0.050 = 2.00 \text{ g} / 2.005 \text{ g} \checkmark$ 2
 (accept $40 \times 0.050 = 2.0 \text{ g}$)
 (mass Ca of 4.0 g would score 1 mark as 'ecf' as molar ratio has not been identified)
- (iii) Ca has reacted with water \checkmark
 $\text{Ca} + 2\text{H}_2\text{O} \rightarrow \text{Ca}(\text{OH})_2 + \text{H}_2 \checkmark\checkmark$
state symbols not required
- 1st mark for H_2 3
 2nd mark is for the rest of the balanced equation
- [6]**
-
20. (i) moles $\text{Ti} = 1.44/47.9 = 0.0301 \text{ mol}/0.03 \text{ mol}$ 1
 (accept use of answer from (b))
- (ii) mass of $\text{Cl} = 5.70 - 1.44 = 4.26 \text{ g} \checkmark$
 moles $\text{Cl} = 4.26/35.5 = 0.120 \text{ mol} \checkmark$ 2
 $5.70/35.5 = 0.161 \text{ mol}$ gets 1 mark
- (iii) $\text{Ti}:\text{Cl} = 0.0301 : 0.12 = 1:4$.
 Empirical formula = $\text{TiCl}_4 \checkmark$
 $0.0301 : 0.161 \text{ mol}$ gives TiCl_5 for 1 mark 1
- (iv) $\text{Ti} + 2\text{Cl}_2 \rightarrow \text{TiCl}_4 \checkmark$ 1
 (ecf possible from (iii))
 covalent \checkmark
- [5]**
-
21. (a) $\dots\text{Mg}(\text{OH})_2(\text{s}) + 2\dots\text{HCl}(\text{aq}) \rightarrow \dots\text{MgCl}_2(\text{aq}) + 2\dots\text{H}_2\text{O}(\text{l}) \checkmark$ 1

- (b) (i) moles HCl = $0.108 \times 500/1000 = 0.054$ ✓ 1
- (ii) moles Mg(OH)₂ = $\frac{1}{2} \times$ moles HCl = 0.027 ✓
 molar mass of Mg(OH)₂ = $24.3 + 17 \times 2 = 58.3$ ✓
 (do not penalise 24)
 mass Mg(OH)₂ = $58.3 \times 0.027 = 1.57$ g / 1.5741 g ✓
 (accept ans from (ii) $\times 0.027 = 1.566$ g)
 (mass Mg(OH)₂ of 3.15 g would score 2 marks as 'ecf' as
 molar ratio has not been identified) 3
- (iii) Too much **if** 2.42 g (dose) > ans to (ii) ✓
 (If answer to (ii) > 2.42 g then 'correct' response here would
 be 'Not enough' 1

[6]

22. (i) Number **AND** type of atoms (making up a
 molecule)/number of atoms of each element ✓ 1
Not ratio
- (ii) P₄ + 6 Br₂ → 4 PBr₃ ✓ 1
- (iii) ratio P : Br = 16.2/31 : 83.8/79.9
 /= 0.52 : 1.05
 /= 1 : 2 ✓
 Empirical formula = PBr₂ ✓
 Correct compound = P₂Br₄ /phosphorus(II) bromide but
not PBr₂ ✓ 3

[5]

23. (i) mass of Ni = 2.0g ✓
 moles of Ni = $2.0/58.7$ mol = 0.0341/0.034 mol ✓
 (1 mark would typically result from no use of 25% → 0.136 mol)
 2nd mark is for the mass of Ni divided by 58.7 2
- (ii) number of atoms of Ni = $6.02 \times 10^{23} \times 0.0341$ 1
 = $2.05 \times 10^{22} / 2.1 \times 10^{22}$ atoms ✓
 Can be rounded down to 2.1 or 2.0 or 2 (if 2.0)
 From 8 g, ans = $8.18/8.2 \times 10^{22}$
 (and other consequential responses)

[3]