


Q1. Water dissociates slightly according to the equation:



The ionic product of water, K_w , is given by the expression

$$K_w = [\text{H}^+][\text{OH}^-]$$

K_w varies with temperature as shown in the table.

Temperature / °C	$K_w / \text{mol}^2 \text{dm}^{-6}$
25	1.00×10^{-14}
50	5.48×10^{-14}

(a) Explain why the expression for K_w does **not** include the concentration of water.

.....

.....

.....

.....

.....

(2)

(b) Explain why the value of K_w increases as the temperature increases.

.....

.....

.....

.....

.....

(2)

(c) Calculate the pH of pure water at 50 °C.
Give your answer to 2 decimal places.

.....
.....
.....
.....
.....
.....
.....
.....

(3)

(d) Calculate the pH of 0.12 mol dm⁻³ aqueous NaOH at 50 °C.
Give your answer to 2 decimal places.

.....
.....
.....
.....
.....
.....
.....
.....

(3)

(Total 10 marks)

Q2. This question is about alkalis and carboxylic acids.

In this question, all data are quoted at 25 °C.

(a) Carboxylic acids are weak acids.

State the meaning of the term **weak** as applied to carboxylic acids.

.....
.....

(1)

(b) Write an equation for the reaction of propanoic acid with sodium carbonate.

.....

(1)

(c) Calculate the pH of a $0.0120 \text{ mol dm}^{-3}$ solution of calcium hydroxide.
The ionic product of water $K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$.
Give your answer to 2 decimal places.

.....
.....
.....
.....
.....
.....

(Extra space).....

.....
.....

(3)

(d) The value of the acid dissociation constant K_a for benzenecarboxylic acid ($\text{C}_6\text{H}_5\text{COOH}$) is $6.31 \times 10^{-5} \text{ mol dm}^{-3}$.

(i) Write an expression for the acid dissociation constant K_a for benzenecarboxylic acid.

.....
.....

(1)

(ii) Calculate the pH of a $0.0120 \text{ mol dm}^{-3}$ solution of benzenecarboxylic acid.
Give your answer to 2 decimal places.

.....
.....

.....
.....
.....
.....
.....
.....
.....

(Extra space).....
.....
.....

(3)

- (iii) A buffer solution with a pH of 4.00 is made using benzenecarboxylic acid and sodium benzenecarboxylate.

Calculate the mass of sodium benzenecarboxylate ($M_r = 144.0$) that should be dissolved in 1.00 dm^3 of a $0.0120 \text{ mol dm}^{-3}$ solution of benzenecarboxylic acid to produce a buffer solution with a pH of 4.00

The value of the acid dissociation constant K_a for benzenecarboxylic acid ($\text{C}_6\text{H}_5\text{COOH}$) is $6.31 \times 10^{-5} \text{ mol dm}^{-3}$.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

(Extra space)
.....
.....
.....

- (e) Two solutions, one with a pH of 4.00 and the other with a pH of 9.00, were left open to the air.

The pH of the pH 9.00 solution changed more than that of the other solution.

Suggest what substance might be present in the air to cause the pH to change. Explain how and why the pH of the pH 9.00 solution changes.

Substance present in air

Explanation

.....
.....
.....
.....

(3)
(Total 17 marks)

Q3. This question involves calculations about two strong acids and one weak acid. All measurements were carried out at 25 °C.

- (a) A 25.0 cm³ sample of 0.0850 mol dm⁻³ hydrochloric acid was placed in a beaker and 100 cm³ of distilled water were added. Calculate the pH of the new solution formed. Give your answer to 2 decimal places.

.....
.....
.....
.....

(Extra space)

.....

(2)

- (b) HX is a weak monobasic acid.

(i) Write an expression for the acid dissociation constant, K_a , for HX.

.....
.....

(1)

(ii) The pH of a $0.0850 \text{ mol dm}^{-3}$ solution of HX is 2.79
Calculate a value for the acid dissociation constant, K_a , of this acid.
Give your answer to 3 significant figures.

.....
.....
.....
.....
.....
.....
.....

(Extra space)
.....
.....
.....

(3)

(c) A 25.0 cm^3 sample of $0.620 \text{ mol dm}^{-3}$ nitric acid was placed in a beaker and 38.2 cm^3 of $0.550 \text{ mol dm}^{-3}$ aqueous sodium hydroxide were added.
Calculate the pH of the solution formed.
Give your answer to 2 decimal places.

The ionic product of water $K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ at $25 \text{ }^\circ\text{C}$.

.....
.....
.....
.....
.....
.....

.....
.....
.....
.....

(Extra space)
.....
.....
.....
.....
.....

(6)

(Total 12 marks)

Q4. This question is about the pH of some solutions containing potassium hydroxide and ethanoic acid.

Give all values of pH to 2 decimal places.

(a) (i) Write an expression for pH.

.....

(1)

(ii) Write an expression for the ionic product of water, K_w

.....

(1)

(iii) At 10 °C, a 0.154 mol dm⁻³ solution of potassium hydroxide has a pH of 13.72. Calculate the value of K_w at 10 °C.

.....
.....
.....
.....
(Extra space)
.....
.....

(2)

(b) At 25 °C, the acid dissociation constant K_a for ethanoic acid has the value $1.75 \times 10^{-5} \text{ mol dm}^{-3}$.

(i) Write an expression for K_a for ethanoic acid.

.....
.....

(1)

(ii) Calculate the pH of a $0.154 \text{ mol dm}^{-3}$ solution of ethanoic acid at 25 °C.

.....
.....
.....
.....
.....
.....
.....
.....

(Extra space)
.....
.....

(3)

(c) At 25 °C, the acid dissociation constant K_a for ethanoic acid has the value $1.75 \times 10^{-5} \text{ mol dm}^{-3}$.

- (i) Calculate the pH of the solution formed when 10.0 cm³ of 0.154 mol dm⁻³ potassium hydroxide are added to 20.0 cm³ of 0.154 mol dm⁻³ ethanoic acid at 25 °C.

.....
.....
.....
.....
.....
.....
.....
.....

(Extra space)
.....
.....
.....

(4)

- (ii) Calculate the pH of the solution formed when 40.0 cm³ of 0.154 mol dm⁻³ potassium hydroxide are added to 20.0 cm³ of 0.154 mol dm⁻³ ethanoic acid at 25 °C.

At 25 °C, K_w has the value $1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

(Extra space)
.....

.....
.....

(4)
(Total 16 marks)