

Q1. (a) By reference to the forces between molecules, explain why ammonia is very soluble in water.

.....
.....

(2)

(b) Aqueous solutions of ammonia have a pH greater than 7.

(i) Write an equation for the reaction of ammonia with water.

.....

(ii) Explain why the pH of a solution containing 1.0 mol dm^{-3} of ammonia is less than 14 at 298 K.

.....
.....

(3)

(c) An ammonium ion in aqueous solution can behave as a Brønsted–Lowry acid. State what is meant by the term *Brønsted–Lowry acid*.

.....

(1)

(d) State what is meant by the term *buffer solution*. Identify a reagent which could be added to a solution of ammonia in order to form a buffer solution.

Buffer solution

.....

Reagent

(3)

(e) An acidic buffer solution is obtained when sodium ethanoate is dissolved in aqueous ethanoic acid.

(i) Calculate the pH of the buffer solution formed at 298 K when 0.125 mol of sodium ethanoate is dissolved in 250 cm³ of a 1.00 mol dm⁻³ solution of ethanoic acid.

The acid dissociation constant, K_a , for ethanoic acid is 1.70×10^{-5} mol dm⁻³ at 298 K.

.....
.....
.....
.....
.....
.....

(ii) Write an ionic equation for the reaction which occurs when a small volume of dilute hydrochloric acid is added to this buffer solution.

.....

(5)
(Total 14 marks)

Q2. An aqueous solution contains 4.0 g of sodium hydroxide in 250 cm³ of solution.
($K_w = 1.00 \times 10^{-14}$ mol² dm⁻⁶)

The pH of the solution is

- A 13.0
- B 13.3
- C 13.6
- D 13.9

(Total 1 mark)

Q3. (a) At 50°C, the ionic product of water, K_w , has the value $5.48 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$.

(i) Define the term K_w

.....

(ii) Define the term pH

.....

(iii) Calculate the pH of pure water at 50 °C. Explain why pure water at 50 °C is still neutral even though its pH is not 7.

Calculation

.....

.....

Explanation

.....

(5)

(b) At 25°C, K_w has the value $1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$. Calculate the pH at 25 °C of

(i) a $0.150 \text{ mol dm}^{-3}$ solution of sodium hydroxide,

.....

.....

.....

(ii) the solution formed when 35.0 cm^3 of this solution of sodium hydroxide is mixed with 40.0 cm^3 of a $0.120 \text{ mol dm}^{-3}$ solution of hydrochloric acid.

.....

.....

.....

.....
.....
.....
.....

(8)

(c) In a $0.150 \text{ mol dm}^{-3}$ solution of a weak acid HX at $25 \text{ }^\circ\text{C}$, 1.80% of the acid molecules are dissociated into ions.

(i) Write an expression for K_a for the acid HX.

.....
.....

(ii) Calculate the value of K_a for the acid HX at this temperature and state its units.

.....
.....
.....
.....
.....
.....

(5)

(Total 18 marks)