


- M1.** (a) (i)  $\text{MnO}_2$  (+) 4 1
- (ii)  $\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \longrightarrow \text{Mn}^{2+} + 2\text{H}_2\text{O}$   
*Or multiples*  
*Ignore state symbols*  
*Credit electrons subtracted from RHS*  
*Ignore absence of charge on e* 1
- (iii) Iodide ion(s) is/are oxidised because they have lost electron(s)  
*Do not penalise reference to iodine; the mark is for electron loss* 1
- (b) (i) **M1**  $\text{Cl}_2$  0  
**M2**  $\text{HClO}$  (+) 1 2
- (ii) **M1** Equilibrium will shift/move to the right  
OR L to R  
OR to favour the forward reaction  
OR to produce more HClO  
**M2** Consequential on correct M1  
To oppose the loss of HClO  
OR replaces the HClO (that has reacted)  
*for M2*  
*NOT just "to oppose the change"* 2
- (c) (i) The answers can be in either order  
**M1**  $2\text{Br}^- \longrightarrow \text{Br}_2 + 2\text{e}^-$   
**M2**  $4\text{H}^+ + \text{SO}_4^{2-} + 2\text{e}^- \longrightarrow \text{SO}_2 + 2\text{H}_2\text{O}$

OR




*NOT multiples*

*Ignore state symbols*

*Credit electrons subtracted from incorrect side*

*Ignore absence of charge on e*

2



OR



*Credit ionic equations*

1

- (iii) For M1 and M2, chloride ions are weaker reducing agents than bromide ions, because

**M1 Relative size of ions**

Chloride ions are smaller than bromide ions OR  
chloride ion electron(s) are closer to the nucleus  
OR chloride ion has fewer (electron) shells/levels  
OR chloride ion has less shielding (or converse for  
bromide ion)

**M2 Strength of attraction for electron being lost**

Outer shell/level electron(s) OR electron(s) lost  
from a chloride ion is more strongly held by the  
nucleus compared with that lost from a bromide  
ion (or converse for bromide ion)

*If the forces are described as intermolecular or Van der  
Waals then CE = 0*

*Ignore general reference to Group 7 trend*


*For M1 accept reference to chlorine/bromine or reference to  
atoms of these but NOT "chloride/bromide atoms" or  
"chlorine/bromine molecules"*

*For M2 insist on reference to the correct ions*

*This is the expected answer, but award credit for a candidate  
who gives a correct explanation in terms of hydration  
enthalpy, electron affinity and atomisation enthalpy.*

2

**M2.** (a) (i) **M1 Initiation**




**M2 First propagation**



**M3 Second propagation**



**M4 Termination (must make  $\text{C}_4\text{H}_{10}$ )**



*Penalise absence of dot once only.*

*Penalise + or – charges every time*

*Penalise incorrect position of dot on ethyl radical once only.*

*Penalise  $\text{C}_2\text{H}_5\cdot$  once only*

*Accept  $\text{CH}_3\text{CH}_2\cdot$  with the radical dot above/below/to the side of the  $\text{CH}_2$*

*Mark independently*

4

(ii) **M1** ultra-violet/uv/sun light  
OR (very) high temperature OR  $500\text{ }^\circ\text{C} \geq T \leq 1000\text{ }^\circ\text{C}$

**M2** (free-)radical substitution

*Ignore "heat" for M1*


*Both words needed for M2*

*For M2, ignore the word "mechanism"*

2

(b) (i)  $\text{Cl}_2 + \text{H}_2\text{O} \longrightarrow \text{HClO} + \text{HCl}$

**OR**



*Accept HOCl or ClOH*

*Accept other ionic or mixed representations*

- (ii) **M1** Any one from
- in swimming pools
  - in drinking water
  - to sterilise/disinfect/sanitise water
  - in water treatment
- Ignore the manufacture of bleach*  
*Ignore "to clean water"*  
*Ignore "water purification"*
- M2** The (health) benefit outweighs the risk or wtte  
OR a clear statement that once it has done its job,  
little of it remains OR used in (very) dilute concentrations/  
small amounts/low doses
- Mark independently but M1 can score from (M2) explanation*

2

- (iii) Sodium chlorate(I) or sodium hypochlorite
- Must be named*  
*Ignore (in)correct formulae*  
*Insist on the (I) in the name*

1

- (c) (i)  $\text{Cl}_2 + 2\text{Br}^- \longrightarrow \text{Br}_2 + 2\text{Cl}^-$
- Or half this equation*  
*Ignore state symbols*

1

- (ii) **M1** **The relative size (of the molecules/atoms)**  
Bromine is larger than chlorine OR has more  
electrons/electron shells  
**OR** It is larger/It has a larger atomic radius/it is a  
larger molecule/atom
- M2** **How size of the intermolecular force affects  
energy needed**  
The forces between bromine/ $\text{Br}_2$  molecules are  
stronger (than the forces between chlorine/ $\text{Cl}_2$   
molecules leading to more energy needed to  
separate the molecules) (or converse)  
**OR** bromine/ $\text{Br}_2$  has stronger/more (VdW) intermolecular  
forces.  
(or converse)

For M1 ignore whether it refers to molecules or atoms.

CE = 0 for reference to (halide) ions

Ignore molecular mass

**QoL** for clear reference to the difference in size of the force between molecules

Penalise M2 if covalent bonds are broken

2

[13]

**M3.** (a) **M1** Cl<sub>2</sub> (provides the pale green colour)

*M1 requires the formula*

**M2** NaOH reacts with the acid(s)/the HCl/the HClO/H<sup>+</sup>

*Ignore "reacts with the products"*

*Ignore "reacts with chloride ion"*

*Ignore "reacts with chlorine"*

**M3 requires a correct answer in M2**

Equilibrium shifts (from left ) to right **OR** wtte

3

(b) **M1** A reducing agent is an electron donor OR (readily) loses/ gives away electrons

*Penalise M1 if "electron pair donor"*

**M2** Cl<sub>2</sub> + 2e<sup>-</sup> → 2Cl<sup>-</sup>

For M3 and M4, iodide ions are stronger reducing agents than chloride ions, because

*Ignore state symbols in M2 Accept no charge on the electron*

*Credit the electrons being lost on the RHS*

**M3 Relative size of ions/atomic radius/ionic radius**

Iodide ions are larger/have more (electron) shells/levels than chloride ions (or converse for chloride ion) OR electron(s) to be lost/outer shell/level is further from the nucleus (or converse for chloride ion) OR greater/more shielding

*For M3 insist on "iodide ions"*


**M4 Strength of attraction for electron(s) being lost**

Electron(s) lost from an iodide ion is less strongly held by the nucleus compared with that lost from a chloride ion

*M3 and M4 must be comparative and should refer to electrons.*

(assume argument refers to iodide ions but accept converse argument for chloride ions)

4



*Or multiples*

**M2** silver chloride ONLY

*M2 requires a name*

**M3** The solid/precipitate would dissolve

**OR** is soluble

**OR** (It) forms a (colourless) solution

*Mark M3 independently*


*Ignore "disappears"*

3

(d) Electrophilic addition

1

Mechanism:



*M2 Penalise partial charges if wrong way around, otherwise ignore*

*Max 3 marks **for the mechanism** for wrong reactant and/or "sticks" (wrong reactant could be HBr or Br<sub>2</sub> or incorrect alkene)*

**M1** must show an arrow from the double bond towards one of the Cl atoms on a Cl-Cl molecule.

**M2** must show the breaking of the Cl-Cl bond.

**M3** is for the structure of the carbocation with Cl substituent.

**M4** must show an arrow from the lone pair of electrons on a negatively charged chloride ion towards the positively charged carbon atom.

4

[15]