

Q1. A mass M hangs in equilibrium on a spring. M is made to oscillate about the equilibrium position by pulling it down 10 cm and releasing it. The time for M to travel back to the equilibrium position for the first time is 0.50 s. Which line, **A** to **D**, is correct for these oscillations?

	amplitude/cm	period/s
A	10	1.0
B	10	2.0
C	20	2.0
D	20	1.0

(Total 1 mark)

Q2. Which one of the following statements is true when an object performs simple harmonic motion about a central point O?

- A** The acceleration is always away from O.
- B** The acceleration and velocity are always in opposite directions.
- C** The acceleration and the displacement from O are always in the same direction.
- D** The graph of acceleration against displacement is a straight line.

(Total 1 mark)

Q3. A girl of mass 40 kg stands on a roundabout 2.0 m from the vertical axis as the roundabout rotates uniformly with a period of 3.0 s. The horizontal force acting on the girl is approximately

- A zero.
- B 3.5×10^2 N.
- C 7.2×10^2 N.
- D 2.8×10^4 N.

(Total 1 mark)

Q4. For a particle moving in a circle with uniform speed, which one of the following statements is **incorrect**?

- A The velocity of the particle is constant.
- B The force on the particle is always perpendicular to the velocity of the particle.
- C There is no displacement of the particle in the direction of the force.
- D The kinetic energy of the particle is constant.

(Total 1 mark)

Q5. A simple pendulum and a mass-spring system are taken to the Moon, where the gravitational field strength is less than on Earth. Which line, **A** to **D**, correctly describes the change, if any, in the period when compared with its value on Earth?

	period of pendulum	period of mass-spring system
A	decrease	decrease
B	increase	increase
C	no change	decrease
D	increase	no change

(Total 1 mark)


Q6. A body moves with simple harmonic motion of amplitude A and frequency $\frac{b}{2\pi}$.

What is the magnitude of the acceleration when the body is at maximum displacement?

- A** zero
- B** $4\pi^2 Ab^2$
- C** Ab^2
- D** $\frac{4\pi^2 A}{b^2}$

(Total 1 mark)

Q7.



A ball of mass m , which is fixed to the end of a light string of length l , is released from rest at X. It swings in a circular path, passing through the lowest point Y at speed v . If the tension in the string at Y is T , which one of the following equations represents a correct application of Newton's laws of motion to the ball at Y?

A $T = \frac{mv^2}{l} - mg$

B $T - mg = \frac{mv^2}{l}$

C $mg - T = \frac{mv^2}{l}$

D $T + \frac{mv^2}{l} = mg$

(Total 1 mark)

Q8. A body is in simple harmonic motion of amplitude 0.50 m and period 4π seconds. What is the speed of the body when the displacement of the body is 0.30 m?

- A** 0.10ms^{-1}
- B** 0.15ms^{-1}
- C** 0.20 m s^{-1}
- D** 0.40 m s^{-1}


(Total 1 mark)

Q9. Which one of the following statements always applies to a damping force acting on a vibrating system?

- A** It is in the same direction as the acceleration.
- B** It is in the same direction as the displacement.
- C** It is in the opposite direction to the velocity.
- D** It is proportional to the displacement.

(Total 1 mark)

Q10.



A simple pendulum consists of a bob of mass m on the end of a light string of length l . The bob is released from rest at X when the string is horizontal. When the bob passes through Y its velocity is v and the tension in the string is T . Which one of the following equations gives the correct value of T ?

A $T = mg$

B $T = \frac{mv^2}{l}$

C $T + mg = \frac{mv^2}{l}$

D $T - mg = \frac{mv^2}{l}$

(Total 1 mark)

Q11. A particle of mass m executes simple harmonic motion in a straight line with amplitude A and frequency f . Which one of the following expressions represents the total energy of the particle?

- A** $2\pi^2 m f A^2$
- B** $2\pi^2 m f^2 A^2$
- C** $4\pi^2 m^2 f^2 A$
- D** $4\pi^2 m f^2 A^2$

(Total 1 mark)

Q12. A simple pendulum and a mass-spring system both have the same time period T at the surface of the Earth. If taken to another planet where the acceleration due to gravity was half that on Earth, which line, **A-D**, in the table gives correctly the new periods?

	simple pendulum	mass-spring
A	$T\sqrt{2}$	T
B	$\frac{T}{\sqrt{2}}$	T
C	$T\sqrt{2}$	$\frac{T}{\sqrt{2}}$
D	$\frac{T}{\sqrt{2}}$	$T\sqrt{2}$

(Total 1 mark)

Q13. A body undergoes forced oscillation. Which one of the following will **not** be increased by increasing the amplitude of the oscillatory driving force?

- A the amplitude of the driven oscillation
- B the energy of the driven oscillation
- C the frequency of the driven oscillation
- D the power required to maintain the driven oscillation

(Total 1 mark)

Q14. Which one of the following statements is **not** true for a body vibrating in simple harmonic motion when damping is present?

- A The damping force is always in the opposite direction to the velocity.
- B The damping force is always in the opposite direction to the acceleration.
- C The presence of damping gradually reduces the maximum potential energy of the system.
- D The presence of damping gradually reduces the maximum kinetic energy of the system.

(Total 1 mark)


Q15. For which of the following relationships is the quantity y related to the quantity x by the

relationship $x \propto \frac{1}{y}$?

	x	y
A	energy stored in a spring	extension of the spring
B	gravitational field strength	distance from a point mass
C	de Broglie wavelength of an electron	momentum of the electron
D	period of a mass-spring system	spring constant (stiffness) of the spring

(Total 1 mark)

Q16. The diagrams show the variation of velocity and acceleration with time for a body undergoing simple harmonic motion.



Which one of the following is proportional to the change in momentum of the body during the time covered by the graphs?

- A** The area enclosed by the velocity-time graph and the time axis
- B** The gradient of the velocity-time graph at the point **P**
- C** The area enclosed by the acceleration-time graph and the time axis
- D** The gradient of the acceleration-time graph at the point **Q**

(Total 1 mark)

Q17. A particle is oscillating with simple harmonic motion described by the equation:

$$s = 5 \sin (20\pi t)$$

How long does it take the particle to travel from its position of maximum displacement to its mean position?

A $\frac{1}{40}$ s


B $\frac{1}{20}$ s

C $\frac{1}{10}$ s

D $\frac{1}{5}$ s





(Total 1 mark)

Q18. The diagram shows two pendulums suspended from the same thread, **PQ**.



X is a heavy pendulum, the frequency f_x of which can be varied. **Y** is a lighter pendulum of fixed frequency f_y . As the frequency of oscillation of **X** is increased by shortening the thread, the amplitude of the oscillation of **Y** changes.

Which one of the following graphs best represents the relationship between the amplitude a_y of the oscillation of **Y** and the frequency f_x of **X**?



(Total 1 mark)