

- M1.(a) (i) the minimum energy required by an electron ✓  
to escape from a (metal)surface ✓

*if refer to atom / ionisation zero marks*

2

- (ii) the (minimum) energy to remove an electron(from an atom) ✓  
from the ground state ✓

2

- (b) (use of  $hf = eV$ )  
 $6.63 \times 10^{-34} \times f = 5.15 \times 1.60 \times 10^{-19}$  ✓

$$f = \frac{5.15 \times 1.60 \times 10^{-19}}{6.63 \times 10^{-34}} \quad \checkmark = 1.24 \times 10^{15} (\text{Hz})$$

*if no working and  $1.24 \times 10^{15} (\text{Hz})$  1 mark*

2

- (c) (use of  $hf = E_k + \Phi$ )  
 $\Phi = 2.28 \times 1.60 \times 10^{-19} = 3.648 \times 10^{-19} (\text{J})$  ✓  
 $E_k = 5.15 \times 1.60 \times 10^{-19} - 3.648 \times 10^{-19} = 4.59 \times 10^{-19} \text{ J}$  ✓ ✓

*3 sig figs*

*if clearly used  $1.2 \times 10^{15}$  then final answer must be to 2 sig. figs. for last mark to be awarded*

*accept 4.57 in place of 4.59*

3

- (d) (use of  $c = f\lambda$ )

$$\lambda = \frac{3.0 \times 10^8}{1.24 \times 10^{15}} = 2.42 \times 10^{-7} \quad \checkmark$$

$$v = h / m\lambda = 6.63 \times 10^{-34} / (9.11 \times 10^{-31} \times 2.42 \times 10^{-7})$$

$$v = 3010 \text{ m s}^{-1} \quad \checkmark \checkmark$$

*first mark minimum working – determination of wavelength*

*bald answer gets 2 marks*

*range to 3 sig figs 2900 – 3030*

3

[12]

**M2.(a)** The process involves the ejection of electrons which are negatively charged. ✓ ✓

1

Any electrons ejected will only make the positive charge greater. ✓

1

- (b) **The mark scheme gives some guidance as to what statements are expected to be seen in a 1 or 2 mark (L1), 3 or 4 mark (L2) and 5 or 6 mark (L3) answer. Guidance provided in section 3.10 of the ‘Mark Scheme Instructions’ document should be used to assist in marking this question.**

Mark	Criteria	QoWC
6	Both ideas fully analysed, with full discussion of alternatives.	The student presents relevant information coherently, employing structure, style and sp&g to render meaning clear. The text is legible.
5	Both ideas analysed with supporting discussion but without alternatives	
4	Both ideas analysed, with one dealt with satisfactorily and the other with some supporting discussion	The student presents relevant information and in a way which assists the communication of meaning. The text is legible. Sp&g are sufficiently accurate not to obscure meaning.
3	Both ideas analysed, with only one dealt with satisfactorily	
2	One idea analysed with some supporting discussion	The student presents some relevant information in a simple form. The text is usually legible. Sp&g allow meaning to be derived although errors are sometimes obstructive.
1	One idea analysed, with little supporting discussion	
0	Unsupported combination or no relevant analysis	The student's presentation, spelling, punctuation and grammar seriously obstruct

		understanding.
--	--	----------------

*The following statements are likely to be present.*

*To demonstrate threshold frequency:*

*The metal should be kept the same, and the light source varied.*

*Using any metal, and light sources 1 and 3,*

*no charge will be lost with light source 1*

*but charge will be lost with light source 3*

*because light source three has a greater photon energy*

*and therefore frequency (from  $E=hf$ )*

*and is above the threshold frequency  
as the photon energy is greater than the work function of the metal*

*but light source 1 has a photon energy less than the work function of the metal*

*so its frequency is below the threshold frequency.*

*To demonstrate work function*

*The light source should be kept the same, and the metal varied*

*Use light source 2 as the other two will either cause all three metals to lose their charge, or none of the metals to lose their charge.*

*Use each metal in turn, so that zinc loses its charge, due to its low work function, but copper and iron do not lose their charge.*

6

- (c) Work function in joules =  $1.6 \times 10^{-19} \times 4.3 = 6.9 \times 10^{-19} \text{ J}$  ✓

*The first mark is for converting the work function into J*

1

Use of  $hf = \text{work function} + KE_{\text{max}}$

*The second mark is for substituting into the photoelectric equation*

1

$$KE_{\text{max}} = hf - \text{work function}$$

$$= (6.63 \times 10^{-34}) \times (1.2 \times 10^{15}) + 6.9 \times 10^{-19} \quad \checkmark$$

$$= 7.9 \times 10^{-19} - 6.9 \times 10^{-19}$$

$$= 1.0 \times 10^{-19} \text{ J} \quad \checkmark$$

*The third mark is for the final answer*

*Allow 1.1*

1

- (d) The work function is the minimum amount of energy needed to remove the electron from the zinc surface ✓

*Alternative*

*Reference to max  $ke$  corresponding to emission of surface electrons whilst electrons from deeper in the metal will be emitted with smaller  $ke$*

1  
[12]

M3.D

[1]

M4.B

[1]

M5.C

[1]

- M6.(a) energy of photon is constant / fixed OR energy given to electron is fixed ✓  
energy required for electron to leave / escape / emit from the surface / metal  
OR electron has to overcome work function ✓  
maximum kinetic energy is the energy of photon minus the work function ✓  
deeper electrons require energy to get to the surface OR have less  $E_k$  than surface electrons ✓

*mention of energy levels means can only score first mark  
photoelectric equation alternative for third mark if  $\phi$  and  $hf$  defined*

3 max

- (b) (i) (use of  $E = hf$ )  
energy of photon =  $6.63 \times 10^{-34} \times 3.0 \times 10^{15}$  ✓ =  $1.989 \times 10^{-18}$  (J)  
work function =  $hf - E_k = 1.989 \times 10^{-18} - 1.7 \times 10^{-18} = 2.89 \times 10^{-19}$  ✓  
work function =  $2.89 \times 10^{-19} / 1.6 \times 10^{-19}$  ✓ = (1.8 eV)

*$hf$  gets first mark even if in wrong equation*

3

- (ii) work function =  $hf_0$   
 $f_0 = 1.8 \times 1.6 \times 10^{-19} / 6.63 \times 10^{-34} \checkmark = 4.3 \times 10^{14} \checkmark$  (Hz)  $\checkmark$  (2 sig figs)  
 2 sig . fig stand alone mark  
 Accept  $4.4 \times 10^{14}$

3

- (c) (i) decrease the energy of( incident) photons  $\checkmark$   
 decrease the maximum kinetic energy of electrons  $\checkmark$   
 OR  
 decrease the energy of( incident) photons  $\checkmark$   
 hence fewer deeper electrons escape  $\checkmark$   
 OR  
 below threshold frequency  $\checkmark$   
 no electrons emitted  $\checkmark$   
 OR  
 as energy of each photon decreases but intensity is constant ( there are more photons / sec)  $\checkmark$   
 number of emitted electrons(/sec) must increase  $\checkmark$   
*for last two alternatives must get first mark before can qualify for second mark*

2

- (ii) increase in photons cause increase in (emitted) electrons  $\checkmark$   
double number of electrons / photons OR reference to rate /per second  $\checkmark$   
*if refer to energy levels / atoms can only award first mark*

2

[13]

**M7.(a)** Minimum energy to remove an electron

**B1**

from a (metal) surface

**B1**

2

- (b) Converts 2.28 (e V) to  $3.6 \times 10^{-19}$  (J) /  $2.28 \times 1.6 \times 10^{-19}$

**C1**

*Condone minus sign here on energy or*

*charge*

Use of  $hf = \varphi_0$

e.g.  $f = 2.28 / h$  (will need to see subject)

or  $2.28 = 6.6(3) \times 10^{-34} \times f$  or  $f =$

$2.28 / 6.6(3) \times 10^{-34}$  (will need to see subject )

*Makes  $f$  subject or substitutes correctly for  $h$  and  $\varphi_0$ .*

**C1**

allow equivalent substitution into  $hf = \varphi_0 + KE_{max}$  where  $KE = 0$

*Penalise minus sign on answer*

$(f =) 5.5(0) \times 10^{14}$  (Hz) cao

**A1**

3

[5]