

M1.(a) (it takes) 130 J / this energy to raise (the temperature of) a mass of 1 kg (of lead) by 1 K / 1 °C (without changing its state) ✓

1 kg can be replaced with unit mass.

Marks for 130J or energy.

+1 kg or unit mass.

+1 K or 1 °C.

Condone the use of 1 °K

1

(b) (using $Q = mc\Delta T + ml$)
= $0.75 \times 130 \times (327.5 - 21) + 0.75 \times 23000$ ✓
(= 29884 + 17250)
= 47134 ✓
= 4.7×10^4 (J) ✓

For the first mark the two terms may appear separately i.e. they do not have to be added.

Marks for substitution + answer + 2 sig figs (that can stand alone).

3

[4]

M2.(a) (i) Appreciates pV should be constant for isothermal change (by working or statement) $W = p\Delta V$ is TO

Allow only products seen where are approximately 150 for 1 mark

Penalise J as unit here

M1

Demonstrates $pV = \text{constant}$ using 2 points (on the line) set equal to each other or conclusion made or **shows** that for V doubling that p halves (worth 2 marks)

need to see values for p and V

Products should equal 150 to 2 sf

Accept statement that products are slightly different so not quite isothermal

A1

Demonstrates $pV = \text{constant}$ using 3 points (on the line) with conclusion

Need to see values for p and V

Products should equal 150 to 2 sf
 Accept statement that products are slightly
 different so not quite isothermal

A1

3

- (ii) Adiabatic therefore no heat transfer **or**
 Adiabatic therefore $Q = 0$

B1

Work is done by gas therefore W is negative **or**
 Work is done by gas therefore energy is removed from
 the system

B1

ΔU is negative therefore internal energy of gas
 decreases **or** energy is removed from the system
therefore internal energy of gas decreases or work
 done by the gas so internal energy decreases

Allow

$$-\Delta U = -W \text{ or } \Delta U = -W$$

B1

3

- (iii) Uses $pV / T = \text{constant}$ or uses $pV = nRT$ or uses
 $pV = NkT$
 e.g. makes T subject or substitutes into an equation
 with p_A and V_A or p_C and V_C (condone use of $n = 1$) or
 $\frac{(pV)_A}{(pV)_C}$
 their $\frac{(pV)_A}{(pV)_C}$

V_a read off range

$$= 2.5 \text{ to } 2.6 (\times 10^{-4})$$

$$p_A = 600 \times 10^3$$

V_c read off range

$$= 8.5 \text{ to } 8.6 (\times 10^{-4})$$

$$p_C = 140 \times 10^3$$

C1

Correct substitution of coordinates (inside range) into

$$\frac{(pV)_A}{(pV)_C}$$

$$\frac{(pV)_A}{(pV)_C}$$

With consistent use of powers of 10

$(pV)_A$ range is 150 to 156 and $(pV)_C$ range is 119 to 120.4

C1

1.2(5) Allow range from 1.2 to 1.3

Accept decimal fraction : 1

A1

3

- (b) Energy per large square = 10(J) **or** states that work done is equal to area under curve (between A and B)
or energy per small square = 0.4(J)
or square counting seen on correct area

Must be clear that area represents energy either by subject of formula or use of units on 10 or 0.4

Alternative:

$W = \text{area of a trapezium}$

(with working)

or $W = P_{\text{mean}} \times \Delta V$ **or**

$W = 450 \times 10^3 \times 2.5 \times 10^{-4}$

or $W = \text{area of a rectangle} + \text{area of a triangle}$ (with working)

B1

Number of large squares = 10.5 to 11.5 seen and (\bar{W}) =
number of squares \times area of one square (using numbers)

Range = 105 to 115 (J)

Or

Number of small squares = 263 to 287 seen and (\bar{W}) =
number of squares \times area of one square (using numbers)

Range = 105 to 115 (J)

States that actual work done would be lower because of curvature of line

B1

2

- (c) (Total energy removed per s =) 4560 (J)
or number of cycles per s = 40
or (Mass per second =) $114 \div 68400$ in rearranged form
or their energy \div ($c \Delta T$) **or** their energy \div 68400

C1

0.067 (kg) seen Allow 0.066 (kg) here
or allow $V / t = 1.67 \times 10^{-3} \div 1100$

or $\left(\frac{V}{t}\right) = \frac{E}{\rho \Delta \theta}$ and correct **substitution** seen

Condone $E = 114$ (J) or temperature = 291(K)

C1

= 0.061×10^{-3} or 6.06×10^{-5} (m³)

A1

3

[14]

M3.B

[1]

M4.A

[1]

M5.(a) the energy required to change the state of a unit mass of water to steam / gas ✓
when at its boiling point temperature / 100°C / without a change in temperature) ✓

allow 1 kg in place of unit

allow liquid to vapour / gas without reference to water

don't allow 'evaporation' in first mark

2

(b) (i) thermal energy given by copper block ($= mc\Delta T$)
= $0.047 \times 390 \times (990 - 100)$
= 1.6×10^4 (J) ✓
2 sig figs ✓

can gain full marks without showing working

a negative answer is not given credit

sig fig mark stands alone

2

- (ii) thermal energy gained by water and copper container
 ($= mc\Delta T_{\text{water}} + mc\Delta T_{\text{copper}}$)
 $= 0.050 \times 4200 \times (100 - 84) + 0.020 \times 390 \times (100 - 84)$
 or
 $= 3500 \text{ (J)} \checkmark \text{ (3485 J)}$
 available heat energy ($= 1.6 \times 10^4 - 3500$) $= 1.3 \times 10^4 \text{ (J)} \checkmark$

allow both 12000 J and 13000 J

allow CE from (i)

working must be shown for a CE

take care in awarding full marks for the final answer – missing out the copper container may result in the correct answer but not be worth any marks because of a physics error

(3485 is a mark in itself)

ignore sign of final answer in CE

(many CE's should result in a negative answer)

2

- (iii) (using $Q = ml$)
 $m = 1.3 \times 10^4 / 2.3 \times 10^6$
 $= 0.0057 \text{ (kg)} \checkmark$
 Allow 0.006 but not 0.0060 (kg)

allow CE from (ii)

answers between 0.0052 \rightarrow 0.0057 kg resulting from use of 12000 and 13000 J

1

[7]

M6.(i) (heat supplied by glass = heat gained by cola)

(use of $m_g c_g \Delta T_g = m_c c_c \Delta T_c$)

1st mark for RHS or LHS of substituted equation

$$0.250 \times 840 \times (30.0 - T_f) = 0.200 \times 4190 \times (T_f - 3.0) \checkmark$$

2nd mark for 8.4°C

$$(210 \times 30 - 210 t_f = 838 T_f - 838 \times 3)$$

$$T_f = 8.4(1) \text{ (°C)} \checkmark$$

Alternatives:

8°C is substituted into equation (on either side shown will get mark)✓

resulting in 4620J~4190J ✓

or

8°C substituted into LHS ✓ (produces $\Delta T = 5.5^\circ\text{C}$ and hence)

$$= 8.5^\circ\text{C} \sim 8^\circ\text{C} \checkmark$$

8°C substituted into RHS ✓
 (produces $\Delta T = 20^\circ\text{C}$ and hence)
 = $10^\circ\text{C} \sim 8^\circ\text{C}$ ✓

2

- (ii) (heat gained by ice = heat lost by glass + heat lost by cola)
NB correct answer does not necessarily get full marks

(heat gained by ice = $mc\Delta T + ml$)
 heat gained by ice = $m \times 4190 \times 3.0 + m \times 3.34 \times 10^5$ ✓
 (heat gained by ice = $m \times 346600$)

3rd mark is only given if the previous 2 marks are awarded

heat lost by glass + heat lost by cola
 = $0.250 \times 840 \times (8.41 - 3.0) + 0.200 \times 4190 \times (8.41 - 3.0)$ ✓
 (= 5670 J)

(especially look for $m \times 4190 \times 3.0$)

the first two marks are given for the formation of the substituted equation not the calculated values

$m (=5670 / 346600) = 0.016$ (kg) ✓

if 8°C is used the final answer is 0.015 kg

or (using cola returning to its original temperature)

(heat supplied by glass = heat gained by ice)

(heat gained by glass = $0.250 \times 840 \times (30.0 - 3.0)$)

heat gained by glass = 5670 (J) ✓

(heat used by ice = $mc\Delta T + ml$)

heat used by ice = $m(4190 \times 3.0 + 3.34 \times 10^5)$ ✓ (= $m(346600)$)

$m (=5670 / 346600) = 0.016$ (kg) ✓

3

[5]

M7.

(a)
$$\Delta T = \left(\frac{\Delta Q}{mc} \right) = \frac{8.5 \times 10^3}{4200 \times 0.12}$$
 ✓

17 K ✓

2

(b) $\left(\frac{\Delta T}{\Delta t} = \frac{\Delta Q}{mc} \right) = \frac{100 - 26}{\Delta t} = \frac{8.5 \times 10^3}{0.41 \times 4200}$ ✓

$t = 15 \text{ s}$ ✓

2

[4]