

**Q1.** A liquid flows continuously through a chamber that contains an electric heater. When the steady state is reached, the liquid leaving the chamber is at a higher temperature than the liquid entering the chamber. The difference in temperature is  $\Delta t$ .

Which of the following will increase  $\Delta t$  with no other change?

- A Increasing the volume flow rate of the liquid
- B Changing the liquid to one with a lower specific heat capacity
- C Using a heating element with a higher resistance
- D Changing the liquid to one that has a higher density

(Total 1 mark)

**Q2.** The temperature of a hot liquid in a container falls at a rate of 2 K per minute just before it begins to solidify. The temperature then remains steady for 20 minutes by which time all the liquid has all solidified.

What is the quantity  $\frac{\text{Specific heat capacity of the liquid}}{\text{Specific latent heat of fusion}}$  ?

- A  $\frac{1}{40} \text{ K}^{-1}$
- B  $\frac{1}{10} \text{ K}^{-1}$
- C  $10 \text{ K}^{-1}$
- D  $40 \text{ K}^{-1}$

(Total 1 mark)

**Q3.** A fixed mass of gas occupies a volume  $V$ . The temperature of the gas increases so that the root mean square velocity of the gas molecules is doubled. What will the new volume be if the pressure remains constant?

A  $\frac{V}{2}$


B  $\frac{V}{\sqrt{2}}$

C  $2V$

D  $4V$

(Total 1 mark)

**Q4.** **X** and **Y** are two gas bottles that are connected by a tube that has negligible volume compared with the volume of each bottle.



Initially the valve **W** is closed.

**X** has a volume  $2V$  and contains hydrogen at a pressure of  $p$ .

**Y** has a volume  $V$  and contains hydrogen at a pressure of  $2p$ .

**X** and **Y** are both initially at the same temperature.

**W** is now opened. Assuming that there is no change in temperature, what is the new gas pressure?

A  $\frac{2}{3}p$





B  $\frac{5}{3}p$

C  $\frac{4}{3}p$

D  $\frac{3}{2}p$


(Total 1 mark)

**Q5.** Which one of the graphs below shows the relationship between the internal energy of an ideal gas ( $y$ -axis) and the absolute temperature of the gas ( $x$ -axis)?



(Total 1 mark)

**Q6.** The diagram shows the  $p$ - $V$  diagram of an ideal hot-air engine. **WX** and **YZ** are isothermal changes.



Which line of the table below correctly indicates the nature of the work done **on** or **by** the air in each part of the cycle?

	<b>WX</b>	<b>XY</b>	<b>YZ</b>	<b>ZW</b>
<b>A</b>	zero	by	zero	on
<b>B</b>	by	zero	on	zero
<b>C</b>	zero	on	zero	by
<b>D</b>	on	zero	by	zero

(Total 1 mark)

**Q7.** The temperature of a room increases from 283K to 293K. The r.m.s. speed of the air molecules in the room increases by a factor of

- A** 1.02
- B** 1.04
- C** 1.41
- D** 2.00

(Total 1 mark)

**Q8.** A fixed mass of an ideal gas initially has a volume  $V$  and an absolute temperature  $T$ . Its initial pressure could be doubled by changing its volume and temperature to

- A  $V/2$  and  $4T$
- B  $V/4$  and  $T/2$
- C  $2V$  and  $T/4$
- D  $4V$  and  $2T$

(Total 1 mark)

**Q9.** A car of mass  $M$  travelling at speed  $V$  comes to rest using its brakes. Energy is dissipated in the brake discs of total mass  $m$  and specific heat capacity  $c$ . The rise in temperature of the brake discs can be estimated from

- A  $\frac{mV^2}{2Mc}$
- B  $\frac{2MV^2}{mc}$
- C  $\frac{MV^2}{2mc}$
- D  $\frac{2mc}{MV^2}$


(Total 1 mark)

**Q10.** Which one of the following is **not** an assumption about the properties of particles in the simple kinetic theory?

- A**  $\frac{1}{2} m \langle c^2 \rangle$  is the average speed of the particles
- B** The forces between the particles are negligible except when particles collide
- C** The time spent by particles in collision is negligible compared with the time spent between collisions
- D** The volume of the particles is negligible compared to the volume of the container

(Total 1 mark)

**Q11.** The diagram shows a  $p$ - $V$  graph for a fixed mass of gas. The volume increases from  $V_1$  to  $V_2$  while the pressure falls from  $p_1$  to  $p_2$ .




Which one of the paths **A**, **B**, **C** or **D** will result in the greatest amount of work being done by the gas?

(Total 1 mark)

**Q12.** In the diagram the dashed line **X** shows the variation of pressure,  $p$ , with absolute temperature,  $T$ , for 1 mol of an ideal gas in a container of fixed volume.

Which line, **A**, **B**, **C** or **D** shows the variation for 2 mol of the gas in the same container?



(Total 1 mark)


**Q13.** A raindrop of mass  $m$  falls to the ground at its terminal speed  $v$ . The specific heat capacity of water is  $c$  and the acceleration of free fall is  $g$ . Given that 25% of the energy is retained in the raindrop when it strikes the ground, what is the rise in temperature of the raindrop?

- A  $\frac{mv^2}{8c}$
- B  $\frac{v^2}{4mc}$
- C  $\frac{mg}{4c}$
- D  $\frac{v^2}{8c}$

(Total 1 mark)



**Q14.** The graph shows the relation between the product *pressure × volume*,  $pV$ , and temperature,  $\theta$ , in degrees celsius for 1 mol of an ideal gas for which the molar gas constant is  $R$ .



Which one of the following expressions gives the gradient of this graph?

- A  $\frac{1}{273}$
- B  $\frac{pV}{\theta}$
- C  $\frac{pV}{(\theta - 273)}$
- D  $R$

(Total 1 mark)

**Q15.** At a certain temperature, the root-mean-square speed of the molecules of a fixed volume of an ideal gas is  $C$ . The temperature of the gas is changed so that the pressure is halved. The root-mean-square speed of the molecules becomes

- A  $\frac{c}{4}$
- B  $\frac{c}{2}$
- C  $\frac{c}{\sqrt{2}}$
- D  $2c$

(Total 1 mark)

**Q16.** A  $1.0 \text{ k}\Omega$  resistor is thermally insulated and a potential difference of  $6.0 \text{ V}$  is applied to it for  $2.0$  minutes. The thermal capacity of the resistor is  $9.0 \text{ J K}^{-1}$ . The rise in temperature, in  $\text{K}$ , is

- A**  $1.3 \times 10^{-3}$
- B**  $8.0 \times 10^{-3}$
- C**  $0.48$
- D**  $0.80$

**(Total 1 mark)**