Question number	Answer	Mark
1(a)	В	(1)

Question number	Answer	Marks
1(b)	An answer that combines the following points to provide a plan: • measure known volume of sodium hydroxide solution (1) • add same volume of each of the acids (1) • stir the mixture (1) • record the initial and final temperatures/temperature change (1)	(4)

Question number	Answer	Mark
1(c)	heat energy $ \underbrace{ \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\$	
	progress of reaction	
	 product line, labelled (2)HCI/product(s), to right of and lower than reactant line, labelled H² + CI²/reactants (1) curve drawn on diagram (1) activation energy labelled (1) 	(3)

Question number	Answer	Additional guidance	Mark
1(d)	 calculates energy needed to break bonds (1) calculates energy released in forming bonds (1) calculates energy change (1) evaluation of final answer with negative sign (1) 	Example of calculation Bonds broken = 436 + 243 = 679 (kJ mol ⁻¹) Bonds formed = 2 × 432 = 864 (kJ mol ⁻¹)	
		Energy change = 679 - 864 = -185 (kJ mol ⁻¹) Award full marks for correct numerical answer without working	(4)

Question Number	Answer	Acceptable answers	Mark
2(a)	An explanation linking two of the following temperature decreases (1) {heat / energy} taken in (1) (so process) endothermic (1)	ignore references to bond breaking / making heat given out / exothermic = 1 max.	(2)

Question Number	Answer	Acceptable answers	Mark
2(b)	Shown correctly on diagram: horizontal line to right of reactant (1) product line below reactant line (1)	ignore any connecting lines product label not needed	(2)

Question	Answer	Acceptable answers	Mark
Number			
2 (c)	D : heat energy is required heat		(1)
	energy is released		

Questi		Indicative Content	Mark
QWC	*2(d)	An explanation including some of the following points	
		smaller pieces of solid of same mass larger surface area more frequent collisions higher rate of reaction	
		higher temperature particles move faster more frequent collisions particles have more energy more collisions have required energy to react / activation energy more collisions successful higher rate of reaction	(6)
		ORA	
Leve I	0	No rewardable content	
1	1 - 2	a limited explanation of one of factors e.g. at higher temper higher rate e.g. when particles smaller size higher rate the answer communicates ideas using simple language and limited scientific terminology spelling, punctuation and grammar are used with limited according.	uses
2	3 - 4	a simple explanation e.g. at higher temperature particles move faster, more collisions so higher rate e.g. smaller sized particles (of same mass) have greater surface area so higher rate the answer communicates ideas showing some evidence of clarity and organisation and uses scientific terminology appropriately spelling, punctuation and grammar are used with some accuracy	
3	5 - 6	a detailed explanation e.g. (when particles collide they) only react when they have sufficient energy/activation energy and at a higher temperature more of the particles have sufficient energy/activation energy so more collisions will be successful and when particles smaller size higher rate the answer communicates ideas clearly and coherently uses a range of scientific terminology accurately spelling, punctuation and grammar are used with few errors	

Question Number	Answer	Acceptable answers	Mark
3(a)	A use hydrochloric acid which is more dilute		(1)

Question Number	Answer	Acceptable answers	Mark
3 (b)	An explanation linking two of		
	M1 {particles/reactants/collisions} have more energy (1)	atoms/ions/molecules as alternatives to particles	
		reject electrons	
		particles move faster	
	M2 more frequent collisions (1)	more collisions per unit time ignore collisions are more likely/greater chance/probability of collisions/faster collisions	
	M3 more {productive/successful/effective} collisions (1)	more particles have required activation energy	(2)

Question Number	Answer	Acceptable answers	Mark
3 (c)(i)			
	mass of catalyst A O time		
	mass of catalyst B or b time		
	mass of ↑ catalyst □ C		
	mass of catalyst D time		
	mass of catalyst A O time		
	mass of catalyst B organization by time		
	mass of catalyst C O time		
	mass of catalyst		
	0 time		(1)

Question Number	Answer	Acceptable answers	Mark
3 (c)(ii)	$2H_2O_2 \rightarrow 2H_2O + O_2$ (2)	multiples or halves	
	all formulae correct (1)	reject other reactants or products	
	balancing correct formulae (1)	ignore heat on arrow or elsewhere ignore state symbols ignore use of lower case h, lower case o, or use of superscripts or	
		large numbers inside the formulae	(2)

Question Number	Answer	Acceptable answers	Mark
3(d)	An explanation linking		
	M1 energy needed to break bonds / energy released when bonds formed (1)	bond breaking is endothermic / bond making is exothermic if any contradictory statements made in M1, the mark cannot be awarded (and so M2 cannot be awarded either)	
	M2 more heat / energy is released than needed (1) M2 dependent on scoring M1	ignore numbers of bonds eg more bonds formed than broken "more energy is released forming bonds than needed to break bonds" (2)	
			(2)

Question Number	Answer	Acceptable answers	Mark
4(a)(i)	Zn +H ₂ SO ₄ →ZnSO ₄ + H ₂ reactants (1) products (1)	Accept multiples If not correctly balanced max 1 Must be subscripts where relevant	(2)

Question		Indicative Content	Mark	
Number				
QWC	*4(a)	A description including some of the following points		
		general points		
		 reactions occur when particles collide more frequent collisions cause higher rate of reaction mass and size of zinc pieces same so no effect on rate of reaction because same surface area two factors have been altered in the same experiment cannot be certain of effect of each concentration experiment 2 higher/triple concentration of acid 		
		 so more particles (in same volume) so more frequent collisions between particles more successful collisions 		
		 temperature experiment 2 higher temperature particles move faster particles have more energy so more frequent collisions between particles (so 		
		 increased rate) more successful collisions so more energetic collisions between particles more particles have enough energy to react (activation energy) when they collide 	(6)	
Level	0	No rewardable content	<u>I</u>	
1	1 - 2	 a limited description e.g. temperature is higher and concentration is higher so reaction is faster e.g. temperature is higher so particles move faster so reaction is faster the answer communicates ideas using simple language and uses limited scientific terminology spelling, punctuation and grammar are used with limited accuracy 		
2	3 - 4	a simple description e.g. temperature is higher so particles move faster a concentration is higher so more particles so reaction is fas eg when concentration is higher there will be more	ster	
		e.g. when temperature is higher particles move fast more successful collisions so faster reaction		
		the answer communicates ideas showing some evide clarity and organisation and uses scientific terminolo		

		 appropriately spelling, punctuation and grammar are used with some accuracy
3	5 - 6	 a detailed description e.g. higher concentration of acid so more particles so more frequent collisions so faster reaction and higher temperature so particles have more energy so more successful collisions so faster reaction the answer communicates ideas clearly and coherently uses a range of scientific terminology accurately spelling, punctuation and grammar are used with few errors

Question Number	Answer	Acceptable answers	Mark
4(b)(i)	B displacement		(1)

Question	Answer	Acceptable answers	Mark
Number			
4(b) (ii)	 Shown on diagram horizontal reactant line above product line (1) horizontal product line to right of reactant line (1) 	lines must be correctly labelled eg reactants/Zn + CuSO ₄ and products/ CuSO ₄ and Cu ignore any extra lines/curves/labels if not drawn lines but just labels in correct relative positions max 1 If two lines drawn in correct	
		positions but no labels max 1	(2)