

Please write clearly in	block capitals.
Centre number	Candidate number
Surname	
Gamame	
Forename(s)	
Candidate signature	

# AS CHEMISTRY

Paper 1: Inorganic and Physical Chemistry

Friday 27 May 2016 Morning Time allowed: 1 hour 30 minutes

#### **Materials**

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a calculator, which you are expected to use where appropriate.

### **Instructions**

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.


#### Information

- The maximum mark for this paper is 80.
- The Periodic Table/Data Sheet is provided as in insert.


#### Advice

You are advised to spend about 65 minutes on Section A and 25 minutes on Section B.

Section A	<b>1</b>
Answer <b>all</b> questions i	n this section.
This question is about electron configur	ation.
0 1 . 1 Give the full electron configuration of a	n AI atom and of a Cr <sup>3+</sup> ion.
	[2 marks]
Al atom	
Cr <sup>3+</sup> ion	
0 1 . 2 Deduce the formula of the ion that has configuration as krypton.	a charge of 2+ with the same electron
	[1 mark]
0 1 . 3 Deduce the formula of the compound the have the same electron configuration a	nat contains 2+ ions and 3- ions that both s argon.
_	[1 mark]



2	This question is about Period 3 of the Periodic Table.	
0 2 . 1	Deduce which of Na <sup>+</sup> and Mg <sup>2+</sup> is the smaller ion.  Explain your answer.	[2 marks]
	Smaller ion	
	Explanation	
0 2 . 2		nisation
	energy for sodium is measured.	[1 mark]



This question is about a white solid, MHCO<sub>3</sub>, that dissolves in water and reacts with hydrochloric acid to give a salt.

$$MHCO_3 + HCI \rightarrow MCI + H_2O + CO_2$$

A student was asked to design an experiment to determine a value for the  $M_r$  of MHCO<sub>3</sub>. The student dissolved 1464 mg of MHCO<sub>3</sub> in water and made the solution up to 250 cm<sup>3</sup>.

up to 250 cm<sup>3</sup>. 25.0 cm<sup>3</sup> samples of the solution were titrated with 0.102 mol dm<sup>-3</sup> hydrochloric acid. The results are shown in **Table 1**.

Table 1

	Rough	1	2	3
Initial burette reading / cm <sup>3</sup>	0.00	10.00	19.50	29.25
Final burette reading / cm <sup>3</sup>	10.00	19.50	29.25	38.90
Titre / cm <sup>3</sup>	10.00	9.50	9.75	9.65

0 3 . 1	Calculate the mean titre and use this to determine the amount, in moles that reacted with 25.0 $\text{cm}^3$ of the MHCO $_3$ solution.	, of HCI
0 3 . 2	Calculate the amount, in moles, of MHCO $_3$ in 250 cm $^3$ of the solution. Then calculate the experimental value for the $M_{\rm r}$ of MHCO $_3$ . Give your answer to the appropriate number of significant figures.	[3 marks]

0 3 . 3	The student identified use of the burette as the largest source of uncertainty in the experiment.
	Using the same apparatus, suggest how the procedure could be improved to reduce the percentage uncertainty in using the burette.
	Justify your suggested improvement.  [2 marks]
	Suggestion
	Justification
0 3 . 4	Another student is required to make up 250 cm <sup>3</sup> of an aqueous solution that contains a known mass of MHCO <sub>3</sub> . The student is provided with a sample bottle containing the MHCO <sub>3</sub> .
	Describe the method, including apparatus and practical details, that the student should use to prepare the solution.
	[6 marks]
	More answer space is available on page 8

4 Table 2 shows some data about the elements bromine and magnesium.

Table 2


Element	Melting point / K	Boiling point / K
Bromine	266	332
Magnesium	923	1383

In terms of structure and bonding explain why the boiling point of bromine different from that of magnesium. Suggest why magnesium is a liquid ove greater temperature range compared to bromine. [5]	is r a much 5 marks]

5 Figure 2 represents two glass flasks, P and Q, connected via a tap.

Flask **Q** (volume =  $1.00 \times 10^3 \text{ cm}^3$ ) is filled with ammonia (NH<sub>3</sub>) at 102 kPa and 300 K. The tap is closed and there is a vacuum in flask **P**. (Gas constant R =  $8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ )

Figure 2



0	5	1	Calculate the mass of ammonia in flask <b>Q</b> .  Give your answer to the appropriate number of significant figures.	
				[3 marks]


Turn over for the next question	0 5 . 2	When the tap is opened, ammonia passes into flask <b>P</b> . The temperature decreases by 5 °C. The final pressure in both flasks is 75.0 kPa. Calculate the volume, in cm³, of flask <b>P</b> .	
Turn over for the next question			[3 marks]
Turn over for the next question			
Turn over for the next question			
Turn over for the next question			
Turn over for the next question			
Turn over for the next question			
		Turn over for the next question	

6		
0 6 . 1	Explain how ions are accelerated, detected and have their abundance of in a time of flight (TOF) mass spectrometer.	determined [3 marks]
0 6 . 2	Calculate the mass, in kg, of a single $^{52}\mathrm{Cr^+}$ ion. Assume that the mass of a $^{52}\mathrm{Cr^+}$ ion is the same as that of a $^{52}\mathrm{Cr}$ atom.	
	(The Avogadro constant L = $6.022 \times 10^{23} \text{ mol}^{-1}$ )	[1 mark]
-		
-		
0 6 . 3	] In a TOF mass spectrometer the kinetic energy (KE) of a $^{52}\text{Cr}^+$ ion was 1.269 x $10^{-13}\text{J}$	
	Calculate the velocity of the ion using the equation.	
	$KE = \frac{1}{2}mv^2$	
	$(m = \text{mass/kg and } v = \text{velocity/ms}^{-1})$	[2 marks]

**0 6 . 4** Bromine has two isotopes, <sup>79</sup>Br and <sup>81</sup>Br, in approximately equal abundance. In a TOF mass spectrometer bromine forms ions with formula [Br<sub>2</sub>]<sup>+</sup>

Sketch the pattern of peaks you would expect to see in the mass spectrum of a sample of bromine.

[2 marks]



 $oxed{0}$  .  $oxed{5}$  A sample of xenon has  $A_r = 131.31$ . The sample consists of four isotopes. The abundances of three of the isotopes are shown in **Table 3**. The data for one of the isotopes,  ${}^m\!Xe$ , is missing.

Table 3

Isotope	<sup>129</sup> Xe	<sup>131</sup> Xe	<sup>132</sup> Xe	<sup>m</sup> Xe
% abundance	28.0	25.0	27.0	To be calculated

Use the data to calculate the abundance of isotope "Xe and calculate m, the mass number of "Xe. Show your working.

[4 marks]

7	Ammonia reacts with aluminium chloride as shown by the equation:				
	$NH_3 + AICI_3 \rightarrow H_3NAICI_3$				
0 7 . 1	Draw diagrams to illustrate the shapes of NH <sub>3</sub> molecules and of AlCl <sub>3</sub> molecules.				
	Include in your diagrams any lone pairs of electrons that influence the shape.				
	Indicate the values of the bond angles.				
	[3 marks]				

0 7 . 2			
	this bond is formed.	[2 marks]	
	Type of bond		
	Explanation		
0 7 . 3	Explain how the value of the CI-AI-CI bond angle in AICI <sub>3</sub> changes, if at	all, on	
	formation of the compound H <sub>3</sub> NAICI <sub>3</sub>	[2 marks]	
	Turn over for the next question		
	1		

8	A student oxidised a solution of hydrochloric acid with a few drops of sodium chlorate(I) solution. The reaction mixture effervesced and turned pale green. The gas formed bleached universal indicator paper.
0 8 . 1	Write a half-equation for the oxidation of chloride ions.  [1 mark]
0 8 . 2	Write a half-equation for the reduction of chlorate(I) ions to chlorine in acidic conditions.  [1 mark]
08.3	Write an overall equation for the redox reaction of chlorate(I) ions with hydrochloric acid.  [1 mark]
08.4	A solution of sodium chlorate(I) was added to a colourless solution of potassium iodide. Suggest what is observed.  Explain the reaction that leads to this observation.  [3 marks]

9				
0 9 . 1	. 1 A student was given a powder made from a mixture of anhydrous barium chloride and anhydrous magnesium chloride. The student dissolved 1.056 g of the powder in water in a conical flask and added an excess of sulfuric acid. A white precipitate formed and was filtered off, washed and dried. The mass of this solid was 0.764 g.			
	Identify the white precipitate and calculate the percentage, by mass, of magnesium chloride in the powder.	[4 marks]		
	Turn over for the next question			

## Section B

## Answer all questions in the spaces provided

	Allower all questions in the spaces provided	
	wer per question is allowed. wer completely fill in the circle alongside the appropriate answer.	
CORRECT METHOD		
If you want to	change your answer you must cross out your original answer as sh	nown.
If you wish to r shown.	return to an answer previously crossed out, ring the answer you now w	ish to select as
	our working out in the blank spaces around the questions but this will radiational sheets for this working.	not be marked.
1 0	Which element is in the d-block of the Periodic Table?	
		[1 mark]
4	A Selenium	0
1	<b>B</b> Antimony	0
(	C Tantalum	0
1	<b>D</b> Lead	0
1 1 1	Which species contains an element with an oxidation state of +4?	
	A NO <sub>2</sub> <sup>+</sup>	[1 mark]
	B CIO₃¯	
	C H <sub>2</sub> SO <sub>3</sub>	
1	D PCI <sub>5</sub>	
		<del></del>

V A E	There are 392 mol of pure gold in a bar measuring 10 cm by 10 cm What is the density of gold in kg dm <sup>-3</sup> ?  193 19.3 1.93 0.193	[1 mark]
W	<sup>53</sup> Fe <sup>2+</sup> has fewer protons than <sup>56</sup> Fe <sup>2+</sup>	[1 mark]

1 4 The successive ionisation energies for element X are shown in **Figure 3**. Figure 3 30000 25000 20000 15000 Ionisation energy /kJ mol<sup>-1</sup> 10000 5000 0 0 2 6 7 8 9 Number of ionisation Which element is X? [1 mark] A Nitrogen **B** Phosphorus **C** Aluminium **D** Boron 1 5 Which of these decreases down Group 2? [1 mark] A First ionisation energy **B** Atomic radius C Number of protons **D** Reactivity with water

	Refer to the unbalanced equation below when answering question	ns <b>16</b> and <b>17</b> .	
	$K_2Cr_2O_7 + 3H_2C_2O_4 + _H_2SO_4 \rightarrow Cr_2(SO_4)_3 + _H_2O + 6CO$	<sub>2</sub> + K <sub>2</sub> SO <sub>4</sub>	
1 6	In the balanced equation the mole ratio for sulfuric acid to wa	ater is	
	<b>A</b> 1:4 <b>B</b> 1:2	[1 mark	]
	<b>C</b> 4:7		
	<b>D</b> 4:9		
1 7	What is the reducing agent in this reaction?	[4 mayl	- ا
	A H <sup>+</sup>	[1 mark	
	<b>B</b> C <sub>2</sub> O <sub>4</sub> <sup>2-</sup>	0	
	<b>C</b> K <sup>+</sup>	0	
	<b>D</b> $Cr_2O_7^{2-}$	0	

1 8	Which substance exists as a macromolecule?		
		[1 mark]	
	A Cu	0	
	B SiO <sub>2</sub>		
	<b>C</b> P <sub>4</sub> O <sub>10</sub>		
	<b>D</b> MgO	0	
1 9	A pale brown mixture of $NO_2$ and $N_2O_4$ is allowed to reach equilibring gas syringe according to the following equation.	ium in a sealed	
	$2NO_2(g) \rightleftharpoons N_2O_4(g)$		
	When the plunger is pushed further into the syringe the pressure in mixture becomes paler in colour.	ncreases and the	
	When the syringe is placed in a hot oven the mixture becomes darker in colour.		
	Which of the following statements is correct?		
	<b>A</b> NO <sub>2</sub> is brown and the forward reaction is exothermic.	[1 mark]	
	<b>B</b> NO <sub>2</sub> is brown and the forward reaction is endothermic.		
	<b>C</b> NO <sub>2</sub> is colourless and the forward reaction is exothermic.		
	<b>D</b> NO <sub>2</sub> is colourless and the forward reaction is endothermic.	0	

2 0	Which molecule has the largest dipole?		
	A CIF <sub>3</sub>	0	[1 mark]
	<b>B</b> BF <sub>3</sub>	$\circ$	
	C SF <sub>6</sub>	$\bigcirc$	
	D CF <sub>4</sub>	$\bigcirc$	
2 1	In a molecule of a hydrocarbon, the fraction by mass of carbon is $\frac{c}{1}$	<del>9</del> 1	
	What is the empirical formula of the hydrocarbon?		[4 -]
	A CH	$\bigcirc$	[1 mark]
	B CH <sub>3</sub>	$\bigcirc$	
	<b>C</b> C <sub>3</sub> H <sub>8</sub>	$\bigcirc$	
	<b>D</b> C <sub>5</sub> H <sub>12</sub>	$\bigcirc$	

2 2	30 cm <sup>3</sup> of xenon are mixed with 20 cm <sup>3</sup> of fluorine. The gases react according to the following equation. Assume that the temperature and pressure remain constant		
	$Xe(g) + F_2(g) \rightarrow XeF_2(g)$		
	What is the final volume of gas after the reaction is complete?	-	
	<b>A</b> 50 cm <sup>3</sup>	[1	mark]
	<b>B</b> 40 cm <sup>3</sup>	$\bigcirc$	
	<b>C</b> 30 cm <sup>3</sup>	$\bigcirc$	
	<b>D</b> 20 cm <sup>3</sup>	$\bigcirc$	
2 3	Which of the following solutions would react exactly with a solution 0.0500 mol sulfuric acid?	n containin	g
	<b>A</b> 50.0 cm <sup>3</sup> of 1.00 mol dm <sup>-3</sup> KOH	[1	l mark]
	<b>B</b> 100.0 cm <sup>3</sup> of 2.00 mol dm <sup>-3</sup> KOH		
	<b>C</b> 100.0 cm <sup>3</sup> of 2.00 mol dm <sup>-3</sup> Ba(OH) <sub>2</sub>		
	<b>D</b> 50.0 cm <sup>3</sup> of 1.00 mol dm <sup>-3</sup> Ba(OH) <sub>2</sub>	$\bigcirc$	


2 4	In a car airbag, sodium azide (NaN $_{\rm 3}$ ) decomposes to form sodium metal and nitrogen gas.
	$2NaN_3(s) \rightarrow 2Na(s) + 3N_2(g)$
	The sodium metal then reacts with potassium nitrate to produce more nitrogen gas.
	$10Na(s) + 2KNO3(s) \rightarrow N2(g) + 5Na2O(s) + K2O(s)$
	If 2.00 mol of sodium azide react in this way, how many molecules of $N_2$ will be formed? (The Avogadro constant $L = 6.022 \times 10^{23} \text{ mol}^{-1}$ )
	[1 mark
	<b>A</b> $2.41 \times 10^{24}$

**B** 1.93 x 10<sup>24</sup>

**C**  $1.81 \times 10^{24}$ 

**D**  $9.63 \times 10^{23}$ 

**END OF QUESTIONS** 





There are no questions printed on this page
DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED
Copyright Information
For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.
Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.
Copyright © 2016 AQA and its licensors. All rights reserved.