

Please write clearly in	block capitals.		
Centre number		Candidate number	
Surname			
Forename(s)			
Candidate signature			

A-level BIOLOGY

Paper 2

Thursday 13 June 2019

Morning

Materials

For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 91.

Time allowed: 2 hours

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
TOTAL		





Answer all questions in the spaces provided.		
0 1.1	Succession occurs in natural ecosystems. Describe and explain how succession occurs.	
	[4 marks]	
	·	




	Figure 1 shows percentages of energy transferred from sunlight to a zebra in a grassland ecosystem.	Do not write outside the box
	Figure 1	
	Sunlight 2.1% Vegetation 12.7% Zebra 58.2% Faeces and urine 5.7% New tissue	
01.2	Use Figure 1 to calculate the percentage of sunlight energy that would be transferred into the faeces and urine of a zebra. Give your answer to 3 significant figures. [1 mark]	
	Answer = %	
0 1.3	In this ecosystem the net productivity of the vegetation is 24 525 kJ m ^{-2} year ^{-1}	
	Use this information and Figure 1 to calculate the energy stored in new tissues of the zebra in kJ m^{-2} year ⁻¹	
	[2 marks]	
	Answer =kJ m ⁻² year ⁻¹	7



Turn over ►

02	Sickle cell disease (SCD) is a group of inherited disorders. People with SCD have sickle-shaped red blood cells. A single base substitution mutation can cause one type of SCD. This mutation causes a change in the structure of the beta polypeptide chains in haemoglobin.	Do not w outside t box
02.1	Explain how a single base substitution causes a change in the structure of this polypeptide.	
	Do not include details of transcription and translation in your answer. [3 marks]	
	Haematopoietic stem cell transplantation (HSCT) is a long-term treatment for SCD. In HSCT, the patient receives stem cells from the bone marrow of a person who does not have SCD. The donor is often the patient's brother or sister. Before the treatment starts, the patient's faulty bone marrow cells have to be destroyed.	
02.2	Use this information to explain how HSCT is an effective long-term treatment for SCD. [3 marks]	







	The student: • removed the shoot tip from • placed 10 lengths of shoot • added to each Petri dish an • added to each Petri dish 40 • left the Petri dishes at 20 °C • removed the shoots after 5 • determined the mean change Table 1 shows her results.	into each c i identical v) cm ³ of a d C in the dar days and i	of 5 Petri di volume of 5 lifferent cor k with their measured 1	shes 5% glucose acentration 1 lids on for them	solution of IAA solu 5 days	ution
		1	Table 1			
	IAA concentration added to Petri dish / parts per million	10 ⁻⁵	10 ⁻³	10 ⁻¹	1	10
	Mean change in length of shoot / mm	0.0	0.1	1.3	2.4	3.1
03.1	Explain why the student remo	oved the sh	oot tip from	n each seed	lling.	[2 marks]
	[Extra space]					



A student investigated the effects of indoleacetic acid (IAA) on the growth of oat seedlings (young plants).

03.2	Explain why the student added glucose solution to each Petri dish.	2 marks]
03.3	Explain why the lids were kept on the Petri dishes.	2 marks]
0 3.4	Describe and explain the results shown in Table 1 and suggest how the result have differed if lengths of root had been used.	ts might 3 marks]
03.4	have differed if lengths of root had been used.	



Do not write outside the box

0 3.5

The student produced the different concentrations of IAA using a stock 1 g dm^{-3} solution of IAA ($1 \text{ g dm}^{-3} = 1$ part per thousand) and distilled water.

8

Complete **Table 2** with the volumes of stock IAA solution and distilled water required to produce 40 cm^3 of 10 ppm (parts per million) IAA solution.

[1 mark]

Table 2


Concentration of IAA solution / parts per million	Volume of stock IAA solution / cm ³	Volume of distilled water / cm ³
10		

10

Do not write outside the

box







Scientists investigated the effect of a decrease in pH on muscle contraction. The

Do not write outside the

box

A - mouse muscle fibres at typical pH of mouse muscle tissue (control 1).

scientists did the investigation with four different preparations of isolated muscle

B - mouse muscle fibres at 0.5 pH units below typical pH.


tissue: A, B, C and D.

C - rabbit muscle fibres at typical pH of rabbit muscle tissue (control 2).

D - rabbit muscle fibres at 0.5 pH units below typical pH.

They measured the force of muscle contraction of the muscle fibres at 12 °C, 22 °C and 32°C

Figure 3 shows the results the scientists obtained for B and D compared with the appropriate control.








0 4 . 1	A student looked at the results and concluded that a decrease in pH does cause a	Do not write outside the box
	decrease in the force of muscle contraction.	
	Use Figure 3 to evaluate this conclusion.	
	[4 marks]	
	[Extra space]	
	Question 4 continues on the result serve	
	Question 4 continues on the next page	



		Do not write
04.2	Another group of scientists suggested that a decrease in the force of muscle contraction is caused by an increase in the concentration of inorganic phosphate, Pi, in muscle tissues.	outside the box
	Their hypothesis is that an increase in the concentration of Pi prevents the release of calcium ions within muscle tissues.	
	Explain how a decrease in the concentration of calcium ions within muscle tissues could cause a decrease in the force of muscle contraction.	
	[3 marks]	
04.3	In muscles, pyruvate is converted to lactate during prolonged exercise.	
	Explain why converting pyruvate to lactate allows the continued production of ATP by anaerobic respiration. [2 marks]	
		9








0 5.1	Describe the role of glucagon in gluconeogenesis.	
	Do not include in your answer details on the second messenger model of glucagon action.	
		[2 marks]
05.2	The gene that codes for glucagon is 9.531 kilobases in length. The DNA he one complete turn every 10 base pairs. Every complete turn is 3.4 nm in le	
	Use this information to calculate the length in micrometres (μm) of the generative glucagon. Give your answer to 3 significant figures.	e for
		[2 marks]
	A power –	
	Answer =	µm



Do not write outside the box

		Do not write
	Metformin is a drug commonly used to treat type II diabetes. Metformin's ability to lower the blood glucose concentration involves a number of mechanisms including:	outside the box
	increasing a cell's sensitivity to insulininhibiting adenylate cyclase.	
0 5.3	Explain how increasing a cell's sensitivity to insulin will lower the blood glucose concentration.	
	[2 marks]	
0 5 . 4	Evaluin how inhibiting adapted avalage may help to lower the blood gluccos	
0 5.4	Explain how inhibiting adenylate cyclase may help to lower the blood glucose concentration. [3 marks]	
]
		9







0 6.3	Explain one piece of evidence fr not on the X chromosome.	rom Figure 4 to show	that t	he gene for body	colour is
	not on the X chromosome.				[2 marks]
0 6.4	A heterozygous grey-bodied, wh black-bodied, red-eyed male fly.		as cro	ossed with a	
	Complete the genetic diagram b of phenotypes expected in the o			le genotypes and	d the ratio
			•••		[3 marks]
	Phenotypes of parents:	Grey-bodied, white-eyed female		Black-bodied, red-eyed male	
	Genotypes of parents:		_×		
	Genotypes of offspring				
	Phenotypes of offspring				
	Ratio of phenotypes				
	Question 6 cont	tinues on the next pa	ige		

Turn over ►





5 A population of fruit flies contained 64% grey-bodied flies. Use the Hardy–Weinberg equation to calculate the percentage of flies heterozygous for gene **G**.

[2 marks]

Do not write outside the

box

Answer = _____






0 7.1	In photosynthesis, which chemicals are needed for Tick (\checkmark) one box.		Do not write outside the box
	Reduced NADP, ADP, Pi, water and oxygen.	[1 mark]	
	NADP, ATP and water.		
	Reduced NADP, ATP, water and carbon dioxide.		
	NADP, ADP, Pi and water.		
0 7.2	Describe what happens during photoionisation in t	the light-dependent reaction. [2 marks]	
		4	
	Question 7 continues on the nex	tt page	




IB/M/Jun19/7402/2






		Do not write outside the
0 7 . 5	Calculating the R_f values of the pigments can help to identify each pigment. An R_f value compares the distance the pigment has moved from the origin with the distance the solvent front has moved from the origin.	box
	distance pigment has moved from the origin	
	$R_f = \frac{\text{distance pigment has moved from the origin}}{\text{distance solvent front has moved from the origin}}$	
	The distance each pigment has moved is measured from the middle of each spot.	
	Pigment A has an R _f value of 0.95	
	Use Figure 5 to calculate the R _f value of pigment C . [1 mark]	
	R _f value of pigment C =	
0 7.6	The pigments in leaves are different colours. Suggest and explain the advantage of	
	having different coloured pigments in leaves. [1 mark]	
		8
	Turn over for the next question	
	Turn over ►	











Turn over

box

	The lengths of the viral DNA fragments were:	Do not write outside the box
	 600 base pairs 250 base pairs 535 base pairs 300 base pairs 500 base pairs. 	
08.5	Which volunteers had at least one of the viral DNA fragments with 250 base pairs or 535 base pairs? [1 mark]	8

		Do not write
09	The sundew is a small flowering plant, growing in wet habitats such as bogs and marshes. The soil in bogs and marshes is acidic and has very low concentrations of some nutrients. The sundew can trap and digest insects.	outside the box
09.1	Describe how you could estimate the size of a population of sundews in a small marsh.	
	[5 marks]	
09.2	Suggest and explain how digesting insects helps the sundew to grow in soil with very low concentrations of some nutrients.	
	[2 marks]	
		7

Turn over ►

			Do not writ
10	Guillain–Barré syndrome is a rare disease in which the immune system damages the myelin sheath of neurones. Myelin sheath damage can cause a range of symptoms, for example numbness, muscular weakness and muscula paralysis. Sometimes, neurones of the autonomic nervous system are affected, causing heart rate irregularities. Huntington's disease is a disorder caused when a protein called huntingtin damages the brain. Huntingtin is produced because of a dominant, mutant allele.		outside the box
	The first successful drug trial to reduce concentrations of huntingtin in the human brain involved 46 patients. The patients received the drug for 4 months. The concentration of huntingtin was reduced in all the patients. The drug was injected at the base of the spine into the cerebrospinal fluid bathing the brain and spinal cord. The drug contains single-stranded DNA	10	
	molecules. These single-stranded molecules inhibit the mRNA needed to produce huntingtin. Symptoms of Huntington's disease can start at any time, but usually develop between 30 and 50 years of age. The likelihood and age when symptoms start are linked to the number of CAG base sequence repeats in the gene for	15 art	
	Huntington's disease. However, recent studies have suggested that epigenetics may also affect the age when symptoms first start.	20	
1 0 . 1	Damage to the myelin sheath of neurones can cause muscular paralysis (line	s 2–4).	
	Explain how.	3 marks]	



1 0.2	Sometimes Guillain–Barré syndrome causes heart rate irregularities (lines 4–5).	Do no outsic bo
	Suggest and explain why. [3 marks]	
1 0.3	The first successful drug trial to reduce concentrations of huntingtin in the brain used single-stranded DNA molecules (lines 13–14).	
	Suggest and explain how this drug could cause a reduction in the concentration of the	
	protein huntingtin. [3 marks]	



1 0.4	Scientists from the first successful drug trial to reduce concentrations of huntingtin (lines 9–11) reported that the drug is not a cure for Huntington's disease.	
	Suggest two reasons why the drug should not be considered a cure.	
	Do not include repeats of the drug trial in your answer. [2 marks]	
	1	
	2	
10.5	Suggest two reasons why people had the drug injected into the cerebrospinal fluid (lines 12–13) rather than taking a pill containing the drug. [2 marks]	
	1	
	2	
	۲	
1 0.6	Suggest and explain one way epigenetics may affect the age when symptoms of Huntington's disease start.	
	[2 marks]	
	END OF QUESTIONS	
Copyright informati	on	
booklet rather than in	rposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate cluding them on the examination paper or support materials. This booklet is published after each examination series and is available n www.aqa.org.uk after the live examination series.	
	uce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House,	
Copyright © 2019 AQA and its licensors. All rights reserved.		



Do not write outside the box