



GCSE
COMBINED SCIENCE: TRILOGY
8464/P/1F

Physics Paper 1F

Mark scheme

June 2019

Version: 1.0 Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

Information to Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the Examiner make his or her judgement
- the Assessment Objectives, level of demand and specification content that each question is intended to cover.

The extra information is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent.

2. Emboldening and underlining

- 2.1** In a list of acceptable answers where more than one mark is available ‘any **two** from’ is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.
- 2.2** A bold **and** is used to indicate that both parts of the answer are required to award the mark.
- 2.3** Alternative answers acceptable for a mark are indicated by the use of **or**. Different terms in the mark scheme are shown by a / ; e.g. allow smooth / free movement.
- 2.4** Any wording that is underlined is essential for the marking point to be awarded.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general principle to be followed in such a situation is that 'right + wrong = wrong'.

Each error / contradiction negates each correct response. So, if the number of error / contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (indicated as * in examples 1 and 2) are not penalised.

Example 1: What is the pH of an acidic solution?

[1 mark]

Student	Response	Marks awarded
1	green, 5	0
2	red*, 5	1
3	red*, 8	0

Example 2: Name two planets in the solar system.

[2 marks]

Student	Response	Marks awarded
1	Neptune, Mars, Moon*	1
2	Neptune, Sun, Mars	0

3.2 Use of chemical symbols / formulae

If a student writes a chemical symbol / formula instead of a required chemical name, full credit can be given if the symbol / formula is correct and if, in the context of the question, such action is appropriate.

3.3 Marking procedure for calculations

Marks should be awarded for each stage of the calculation completed correctly, as students are instructed to show their working. Full marks can, however, be given for a correct numerical answer, without any working shown.

3.4 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.5 Errors carried forward

Any error in the answers to a structured question should be penalised once only.

Papers should be constructed in such a way that the number of times errors can be carried forward is kept to a minimum. Allowances for errors carried forward are most likely to be restricted to calculation questions and should be shown by the abbreviation ecf in the marking scheme.

3.6 Phonetic spelling

The phonetic spelling of correct scientific terminology should be credited **unless** there is a possible confusion with another technical term.

3.7 Brackets

(.....) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.8 Allow

In the mark scheme additional information, 'allow' is used to indicate creditworthy alternative answers.

3.9 Ignore

Ignore is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

3.10 Do not accept

Do **not** accept means that this is a wrong answer which, even if the correct answer is given as well, will still mean that the mark is not awarded.

4. Level of response marking instructions

Extended response questions are marked on level of response mark schemes.

- Level of response mark schemes are broken down into levels, each of which has a descriptor.
- The descriptor for the level shows the average performance for the level.
- There are two marks in each level.

Before you apply the mark scheme to a student's answer, read through the answer and annotate it (as instructed) to show the qualities that are being looked for. You can then apply the mark scheme.

Step 1: Determine a level

Start at the lowest level of the mark scheme and use it as a ladder to see whether the answer meets the descriptor for that level. The descriptor for the level indicates the different qualities that might be seen in the student's answer for that level. If it meets the lowest level then go to the next one and decide if it meets this level, and so on, until you have a match between the level descriptor and the answer.

When assigning a level you should look at the overall quality of the answer. Do **not** look to penalise small and specific parts of the answer where the student has not performed quite as well as the rest. If the answer covers different aspects of different levels of the mark scheme you should use a best fit approach for defining the level.

Use the variability of the response to help decide the mark within the level, i.e. if the response is predominantly level 2 with a small amount of level 3 material it would be placed in level 2 but be awarded a mark near the top of the level because of the level 3 content.

Step 2: Determine a mark

Once you have assigned a level you need to decide on the mark. The descriptors on how to allocate marks can help with this.

The exemplar materials used during standardisation will help. There will be an answer in the standardising materials which will correspond with each level of the mark scheme. This answer will have been awarded a mark by the Lead Examiner. You can compare the student's answer with the example to determine if it is the same standard, better or worse than the example. You can then use this to allocate a mark for the answer based on the Lead Examiner's mark on the example.

You may well need to read back through the answer as you apply the mark scheme to clarify points and assure yourself that the level and the mark are appropriate.

Indicative content in the mark scheme is provided as a guide for examiners. It is not intended to be exhaustive and you must credit other valid points. Students do **not** have to cover all of the points mentioned in the indicative content to reach the highest level of the mark scheme.

You should ignore any irrelevant points made. However, full marks can be awarded only if there are no incorrect statements that contradict a correct response.

An answer which contains nothing of relevance to the question must be awarded no marks.

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID
01.1	LED		1	AO1.1 AO1 in isolation 6.2.1.1	A
01.2	the same as		1	AO1.1 6.2.1.2	G
01.3	1500 – 900 600 (thousand)	an answer of 600 (thousand) or 600 000 scores 2 marks two correct readings from the graph scores 1 mark allow a range of 1480 to 1520 and a range of 880 to 920 allow an answer in the range of 560 (thousand) to 640 (thousand) consistent with their allowed readings	1 1	AO2.2 6.2.1.2 WS 3.2	G
01.4	repeat the experiment using exactly the same method		1	AO3.3a 6.2.1.2	A
01.5	power = 0.80×0.020 power = 0.016 (W)	an answer of 0.016 (W) scores 2 marks	1 1	AO2.1 6.2.4.1 WS 3.3	E
01.6	power = (current) ² × resistance		1	AO1.1 AO1 in isolation 6.2.4.1	A
01.7	temperature increases		1	AO1.1 6.1.1.1	E

<p>01.8</p>	<p>Q = 0.020×180 Q = 3.6 (C)</p>	<p>an answer of 3.6 (C) scores 2 marks</p>	<p>1 1</p>	<p>AO2.1 6.2.1.2 WS 3.3</p>	<p>E</p>
<p>Total</p>			<p>11</p>		

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID
02.1	the brightness of the lamp		1	AO3/3a 6.1.3c WS 2.2	A
02.2	zero error		1	AO3/3b 6.1.3c WS 3.7	A
02.3	C		1	AO3/1b 6.1.3c WS 3.7	A
02.4	10.0	allow 10	1	AO3/1a 6.1.3c WS 3.5	G
02.5	$\frac{0.96}{8.0}$ = 0.12	an answer of 0.12 or 12% scores 2 marks allow 12%	1 1	AO2.1 6.1.2.2	E
02.6	replenished		1	AO1.1 in isolation 6.1.3b	G
02.7	E = 490 × 31 E = 15 190 E = 15 000 (J)	an answer of 15 000 (J) scores 3 marks allow 15 200 if correct substitution is seen allow an answer to 2 s.f. consistent with their calculated value of E using E=QV	1 1 1	AO2.1 6.2.4.2	E

02.8	less fossil fuel is burned		1	AO3.2a 6.1.3e	A
Total			11		

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID
03.1	they changed direction	allow deflected/reflected/repelled	1	AO 1/1 6.4.1.3	E
03.2	diameter = $\frac{0.18}{6000}$ = 0.000 030 (nm)	an answer of 0.000 03 (nm) or 3.0×10^{-5} (nm) scores 2 marks allow 3.0×10^{-5} (nm)	1 1	AO2/2 6.4.1.1	E
03.3	A		1	AO 1/1 6.4.1.1	A
03.4	1100 (°C)		1	AO3/2b 6.3.2.3	G
03.5	8 (minutes)	allow 12 (minutes)	1	AO3/2b 6.3.2.3	G
03.6	the rate of change of temperature of the gold		1	AO3/1a 6.1.1.3, 6.3.2.2	A
Total			7		

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID
04.1	${}_{91}^{234}\text{Pa}$		1	AO1/1 6.4.1.2	A
04.2	points correctly plotted to within 1 mm a curved line of best fit passing within 1 mm of all 5 points	ignore any line beyond 200 seconds	1 1	AO2.2 6.4.2.3 WS 3.2	E
04.3	70 (s)	allow an answer between 65 and 75 (s) allow an answer consistent with their drawn line	1	AO2/2 6.4.2.3 WS 3.5	E
04.4	70 (s)	allow an answer between 65 and 75 (s) allow their answer to question 04.3	1	AO3/2b 6.4.2.3	E
04.5	beta		1	AO1.1 6.4.2.1	A
04.6	articles in scientific journals are peer reviewed	allow articles in scientific journals are based on evidence/data allow newspaper articles may be oversimplified/inaccurate/biased	1	AO1.1 6.4.2.4 WS 1.6	E
Total			7		

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID
05.1	to stop the metal case of the toaster becoming live if a fault occurs		1	AO1.1 6.2.3.2	A
05.2	yellow brown blue		1 1 1	AO1.1 AO1 in isolation 6.2.3.2	G
05.3	E = 850×120 E = 102 000 (J)	an answer of 102 000 (J) scores 2 marks	1 1	AO2.1 6.2.4.2 6.1.1.4 WS 3.3	E
05.4	elastic potential kinetic		1 1	AO1.1 6.1.1.1	G
05.5	gravitational potential energy = mass \times gravitational field strength \times height or $E_p = m g h$	allow gpe allow any correct re-arrangement	1	AO1.1 AO1 in isolation 6.1.1.2	E
05.6	$0.049 = 0.050 \times 9.8 \times h$ $h = \frac{0.049}{0.050 \times 9.8}$ h = 0.10 (m)	an answer of 0.10 (m) scores 3 marks	1 1 1	AO2.1 6.1.1.2 WS 3.3	E
Total			12		

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID
06.1	ammeter in series with the resistor, voltmeter in parallel with the resistor		1	AO1/1 6.2.1.4 RP 16 WS 2.4	A
06.2	current decreased	ignore slows down	1	AO1/1 6.2.1.3 RP 16 WS 3.6	E
06.3	reverse the connections to the cell	allow battery for cell allow reverse the cell	1	AO1/2 6.2.1.3 RP 16 WS 2.2	E
06.4	(directly) proportional	do not allow inversely proportional do not allow indirectly proportional	1	AO1/2 6.2.1.3 RP 16 WS 3.5	G
06.5	potential difference = current × resistance or $V=IR$	allow voltage for potential difference allow any correct re-arrangement	1	AO1/1 6.2.1.3 RP 16 WS 3.3	E
06.6	$3.0 = 0.12 \times R$ $R = \frac{3.0}{0.12}$ $R = 25 (\Omega)$	an answer of 25 (Ω) scores 3 marks	1 1 1	AO2/1 6.2.1.3 RP 16 WS 3.3	E
Total			8		

Question	Answers	Extra information	Mark	AO / Spec. Ref.	ID																		
07.1	pressure decreased		1	AO2.1	E																		
	because molecules have less (kinetic) energy	allow less speed/velocity	1	6.3.3.1																			
	so fewer collisions (with the wall/container each second)	allow collide with less force allow less force on the walls	1																				
07.2	0.70 = m × 330 or 700 = m × 330 000	an answer of 0.0021(212121...) scores 3 marks	1	AO2.1 6.3.2.2 6.1.1.3	E																		
	$m = \frac{0.70}{330}$ or $m = \frac{700}{330\ 000}$	allow correct rearrangement using converted value(s) of E to J and/or L to J/kg	1																				
	m = 0.0021 (kg)	allow 0.0021(212121...) allow correct calculation using converted value(s) of E and/or L 3 marks can only be awarded for m = 0.0021(212121...) (kg)	1																				
07.3	<table border="1"> <thead> <tr> <th>Substance</th> <th>Solid</th> <th>Liquid</th> <th>Gas</th> </tr> </thead> <tbody> <tr> <td>Oxygen</td> <td></td> <td>✓</td> <td></td> </tr> <tr> <td>Nitrogen</td> <td></td> <td></td> <td>✓</td> </tr> <tr> <td>Carbon dioxide</td> <td>✓</td> <td></td> <td></td> </tr> </tbody> </table>				Substance	Solid	Liquid	Gas	Oxygen		✓		Nitrogen			✓	Carbon dioxide	✓			2	AO3/2b 6.3.1.1	E
	Substance	Solid	Liquid	Gas																			
	Oxygen		✓																				
	Nitrogen			✓																			
	Carbon dioxide	✓																					
2 correct answers scores 1 mark. if more than one tick in a row, neither can score a mark																							

07.4	Level 3: Relevant points (reasons/causes) are identified, given in detail and logically linked to form a clear account.	5–6	AO1.1 6.3.1.2	E
	Level 2: Relevant points (reasons/causes) are identified, and there are attempts at logical linking. The resulting account is not fully clear.	3–4		
	Level 1: Points are identified and stated simply, but their relevance is not clear and there is no attempt at logical linking.	1–2		
	No relevant content	0		
	Indicative content cooling <ul style="list-style-type: none"> • as the argon cools the particles slow down • particles in a liquid move slower than particles in a gas • particles in a solid move slower than particles in a liquid • as the liquid/solid cools the particles get closer together • as the liquid/solid cools the density increases gas to liquid <ul style="list-style-type: none"> • particles change from being spread apart to touching each other • particles will (collide with other particles more often and) change direction more often liquid to solid <ul style="list-style-type: none"> • particles change from a random arrangement to a regular pattern • particles change from moving freely to fixed positions • particles change from moving freely/randomly to vibrating explanation <ul style="list-style-type: none"> • (internal) energy (of the argon) decreases • (kinetic) energy (of the particles) decreases with temperature • (potential) energy (of the particles) changes with change of state (of the argon) • forces between particles in a gas are negligible/zero • attractive forces act between atoms when they are close to each other • attractive forces between particles are stronger in a solid than in a liquid to access level 3 there must be an explanation of changes to arrangement and movement of particles during either cooling or a change of state			
Total		14		