

GCSE MATHEMATICS

New Specimen Papers published June 2015 Paper 2 Higher - Mark Scheme

8300/2H

Version 1.0



Principal Examiners have prepared these mark schemes for specimen papers. These mark schemes have not, therefore, been through the normal process of standardising that would take place for live papers.

Further copies of this Mark Scheme are available from aga.org.uk

Glossary for Mark Schemes

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

M Method marks are awarded for a correct method which could

lead to a correct answer.

A Accuracy marks are awarded when following on from a correct

method. It is not necessary to always see the method. This can

be implied.

B Marks awarded independent of method.

ft Follow through marks. Marks awarded for correct working

following a mistake in an earlier step.

SC Special case. Marks awarded within the scheme for a common

misinterpretation which has some mathematical worth.

M dep A method mark dependent on a previous method mark being

awarded.

B dep A mark that can only be awarded if a previous independent mark

has been awarded.

oe Or equivalent. Accept answers that are equivalent.

eg accept 0.5 as well as $\frac{1}{2}$

[a, b] Accept values between a and b inclusive.

3.14... Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416

Use of brackets It is not necessary to see the bracketed work to award the marks.

Page 2 Version 1.0

Examiners should consistently apply the following principles

Diagrams

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

Responses which appear to come from incorrect methods

Whenever there is doubt as to whether a student has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the student. In cases where there is no doubt that the answer has come from incorrect working then the student should be penalised.

Questions which ask students to show working

Instructions on marking will be given but usually marks are not awarded to students who show no working.

Questions which do not ask students to show working

As a general principle, a correct response is awarded full marks.

Misread or miscopy

Students often copy values from a question incorrectly. If the examiner thinks that the student has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

Further work

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

Choice

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

Work not replaced

Erased or crossed out work that is still legible should be marked.

Work replaced

Erased or crossed out work that has been replaced is not awarded marks.

Premature approximation

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.



	T	T	
Q	Answer	Mark	Comments
1	1 2 4 8	B1	
2	AAA	B1	
3	$a + 20a^2$	B1	
4	y = 5x + 2	B1	
	$\frac{4}{5}$ or 80% seen or used	M1	oe May be implied
5	$29.4(0) \times 5 \div 4$ or $147 \div 4$ or $29.4(0) \div 4 (\times 5)$ or $7.35 (\times 5)$ or $29.4(0) \div 0.8$	M1	oe
	36.75	A1	
6(a)	3 5 7 11 S E E E B E E B B B B B B B B B B B B B	В3	B2 Any 2 or 3 of the 4 sections correct B1 Any 1 of the 4 sections correct
6(b)	1	B1ft	oe ft their Venn diagram

Page 4 Version 1.0

ft their Venn diagram

Q	Answer	Mark	Comments
7(a)	Two of $\frac{6}{50}$ $\frac{28}{100}$ $\frac{34}{150}$	B2	oe fraction, decimal, percentage B1 One of $\frac{6}{50}$ $\frac{28}{100}$ $\frac{34}{150}$ with at most one incorrect answer
7(b)	Chooses their probability from the larger number of trials and reason given that more trials are involved	B1ft	Must have two probabilities in (a)
8	Draws $3x + 2y = 6$ x = 2.5 and $y = -0.7$	B2 B1ft	B1 Works out or plots at least two points satisfying $3x + 2y = 6$ eg (2, 0) and (0, 3) ft their graph $\pm \frac{1}{2}$ square
9(a)	Correct product using at least one prime factor $2 \times 2 \times 3 \times 3 \times 7 \text{ or } 2$	M1	For example 2 (×) 126 or 3 (×) 84 or 7 (×) 36 or 2 (×) 2 (×) 63 or 2 (×) 3 (×) 42 May be implied eg in a factor tree or repeated division
9(b)	$2^2 \times 3^2 \times 7$ 84	B1	



Q	Answer	Mark	Comments		
	Alternative method 1				
	2 parts → 116	M1	oe		
	116 ÷ 2 × 16	M1	oe		
	928	A1			
10	Alternative method 2				
	Writes at least 3 ratios or numbers of boys and girls equivalent to 9:7	M1	eg 18:14 and 180:140 and 360:280		
	522 and 406	M1			
	928	A1			
11	(4)(+ 0) O	B1			
- 11	(x-4)(x+8)=0	DI			
12	$1.7 \times 10^6 \text{ or } 2 \times 10^6$	В3	B2 $1.72(8) \times 10^6$ or 1.73×10^6 or 1.700000 or 2000000		
		_	B1 1728 000 or 1730 000		
13	125 : 27	B1			

Page 6 Version 1.0

Q	Answer	Mark	Comments		
	Alternative method 1				
	10 ÷ 4 or 2.5 or 4 ÷ 10 or 0.4 or $\frac{1}{2}$ × (18 + 10) × 25 or 350	M1	oe		
	18 ÷ their 2.5 or 18 × their 0.4 or 7.2 or 25 ÷ their 2.5 or 25 × their 0.4 or 10	M1dep	oe		
14(a)	$\frac{1}{2} \times (18 + 10) \times 25 \text{ or } 350$ and $\frac{1}{2} \times (\text{their } 7.2 + 4) \times \text{their } 10 \text{ or } 56$	M1dep	Must see working		
	350 – 56 = 294	A1	Do not award without working seen		
	Alternative method 2				
	10 ÷ 4 or 2.5 or 4 ÷ 10 or 0.4 or $\frac{1}{2}$ × (18 + 10) × 25 or 350	M1	oe		
	(Area scale factor =) $(\text{their } 2.5)^2$ or $(\text{their } 0.4)^2$	M1dep			
	their 350 ÷ (their 2.5) ² or their 350 × (their 0.4) ² or 56	M1dep	Must see working		
	350 – 56 = 294	A1	Do not award without working seen		
	18 – 10 2 or 4	B1			
14(b)	$\tan x = \frac{25}{\text{their 4}}$	M1			
	[80.9, 81]	A1			



Q	Answer	Mark	Comments		
	Alternative method 1				
	1800		B2 $a \times b \times c \times d$ with at least 3 correct from 9, 10, 10 and 2		
			B1 $a \times b \times c \times d$ with at least 2 correct from 9, 10, 10 and 2		
		B3	or		
15			identifies 9 possibilities for first digit		
15			or identifies 2 possibilities for final digit		
	Alternative method 2				
	9000	M1	The number of digits between 1000 and 9999 inclusive		
	their 9000 ÷ 5	M1dep			
	1800	A1			
		1	I		
	$6c(c^2+5)$ or $3(c^2+5)$	M1			
16	$\frac{6c(c^2+5)}{3(c^2+5)}$	M1	This mark implies first M1		
	2c and multiple of 2 so even	A1	oe statement Must see method		

Page 8 Version 1.0

Q	Answer	Mark	Comments		
	Alternative method 1				
	$93\ 000\ 000 \times 2\pi$ or $186\ 000\ 000\pi$ or $[584\ 000\ 000\ ,\ 584\ 412\ 000]$	M1	oe Allow working in millions		
	365 × 24 or 8760	M1			
	their 186 000 000 π ÷ their 8760	M1	oe Allow working in millions Only allow if first M1 gained or if their circumference is 93 000 000 \times π		
	$[6.6 \times 10^4, 6.7 \times 10^4]$	A1	oe		
17(a)	Alternative method 2				
	$93\ 000\ 000 \times 2\pi$ or $186\ 000\ 000\pi$ or $[584\ 000\ 000,\ 584\ 412\ 000]$	M1	oe Allow working in millions		
	their 186 000 000π ÷ 365 or [1 598 904, 1 600 033]	M1	oe Allow working in millions Only allow if M1 gained or if their circumference is 93 000 000 \times π their 365.25 can be 365.25 \times 24 or 365.25 \times 60		
	their $[1.6 \times 10^6, 1.602 \times 10^6] \div 24$	M1			
	$[6.6 \times 10^4, 6.7 \times 10^4]$	A1	ое		
17(b)	The average speed would be (slightly) lower	B1	ое		



Q	Answer	Mark	Comments
18(a)	$(n-6)^2$ could be zero (so she is wrong) or The sixth term is 1	B1	oe
8(b)	1	B1	
19	<u>x</u> 3	B1	
	Alternative method 1		
	$2 = k\sqrt{36}$ or $\sqrt{36} = 6$	M1	
	$(k =) 2 \div \text{their 6 or } \frac{1}{3}$	M1dep	
	5 ÷ their $\frac{1}{3}$ or 15 (\sqrt{a} =)	M1	oe
	225	A1	
	Alternative method 2	1	

	$2k = \sqrt{36}$ or $\sqrt{36} = 6$	M1	
20	$(k =)$ their $6 \div 2$ or 3	M1dep	
	$5 \times \text{their 3 or 15 (} \sqrt{a} = \text{)}$	M1	oe
	225	A1	
•	Alternative method 2		

Alternative method 3

$2k = \sqrt{36}$ or $\sqrt{36} = 6$	M1	
5 ÷ 2 or 2.5	M1	
their 6 × their 2.5 or 15 (\sqrt{a} =)	M1dep	dep on M1 M1
225	A1	

Page 10 Version 1.0

Q	Answer	Mark	Comments		
	Alternative method 1				
	1.2 or 0.85	M1			
	1 ÷ 0.85 or 1.1(7) or 1.18	M1			
	1.1(7) or 1.18 and 1.2 and (Option) A	A1			
	Alternative method 2				
	1.2 or 0.85	M1			
	1 ÷ 1.2 or 0.83()	M1			
	0.83() and 0.85 and (Option) A	A1			
	Alternative method 3				
21	450×1.2 or 540 or $x \times 0.85$ or $0.85x$	M1	x is the usual cost of the box and may be a numerical value		
	$x \div \text{their } 540 \text{ or their } 0.85x \div 450$	M1dep			
	0.00185()x and 0.00188()x and (Option) A	A1	oe		
	Alternative method 4				
	450×1.2 or 540 or $x \times 0.85$ or $0.85x$	M1	x is the usual cost of the box and may be a numerical value		
	their 540 ÷ x or 450 ÷ their 0.85 x	M1dep			
	$\frac{540}{x}$ and $\frac{529.()}{x}$ and (Option) A	A1	oe		

Alternative method 5 on next page



Q	Answer	Mark	Comments		
	Alternative method 5				
	1/6 (free for A)	M1	oe fraction or decimal or percentage		
21	$\frac{3}{18}$ (free for A) and $\frac{3}{20}$ (free for B)	M1	oe pairs of fractions or pairs of decimal or pairs of percentages		
	$\frac{3}{18}$ (free for A) and $\frac{3}{20}$ (free for B) and (Option) A	A1			
	Alternative method 1				
	25 × 11 or 275	M1			
	their 275 ÷ 22 or 12.5	M1dep			
	15 × 30 ÷ their 12.5	M1			
	36	A1			
	Alternative method 2				
	25 × 11 or 275	M1			
	15 × 30 ÷ their 275 or [1.6, 1.64]	M1dep			
	their [1.6, 1.64] × 22	M1			
22	36	A1			
	Alternative method 3				
	11 squares				
	or	M1			
	275 squares 22 ÷ 11 or 2				
	or	M1dep			
	22 ÷ 275 or 0.08				
	their 2 × 18	N A A			
	or	M1			
		A1			
	22 ÷ 275 or 0.08 their 2 × 18	M1			

Page 12 Version 1.0

Q	Answer	Mark	Comments
	Alternative method 4		
	15 or 30 25 11	M1	
	$\frac{15}{25} \times \frac{30}{11}$ or $\frac{18}{11}$	M1dep	oe fraction
	their $\frac{18}{11} \times 22$	M1	
22	36	A1	
	Alternative method 5	_	
	$25 \times h = 22$ or $\frac{22}{25}$ or 0.88	M1	oe
	0.88 ÷ 11 or 0.08	M1dep	oe eg frequency density axis labelled with correct scale
	their $0.08 \times 30 \times 15$	M1	
	36	A1	
	$\frac{1}{1} \times \frac{4}{1} \times \pi \times 6_3$ or 144π		00 07 [450 450 45]
	2 3	M1	oe eg [452, 452.45]
	$\frac{2}{5} \times \text{their } 144\pi = \frac{1}{3} \times \pi \times x^2 \times 12$		oe eg [180.8, 181] = [12.5, 12.6] x^2
	5 3 or	M1	Must equate two volumes in terms of π
23	$57.6\pi = 4\pi x^2$		
	$3 imes \frac{2}{5} imes$ their $144\pi \div 12\pi$ or	M1dep	oe eg their [180.8, 181] ÷ their [12.5, 12.6] dep on 2nd M1
	14.4	,	Correct working to isolate x^2
	[3.79, 3.8]	A1	



Q	Answer	Mark	Comments			
24(a)	$\cos x = \frac{OA}{15}$ or $OA = 15 \cos x$	M1				
	$OA = 15 \cos x$					
	and					
	OB = 15 + 2	A1				
	and					
	$h = OB - OA = 17 - 15 \cos x$					
24(b)	17 – 15 cos 120					
	or	M1				
	15 sin 30 or 7.5					
	24.5	A1	oe			
24(c)	(180, 32)	B2	B1 one correct coordinate			
			SC1 (32, 180)			

	Alternative method 1			
25(a)	$a = 2$ or $2(x^2 - 3x + 2.5)$ or $2(x^2 - 3x) + 5$	M1		
	$x^{2} - 3x = (x - 1.5)^{2} - 1.5^{2}$	M1dep	oe ft their $x^2 - 3x$	
	a = 2 and $b = 1.5$ and $c = 0.5$	A1	oe eg $2(x-1.5)^2+0.5$	
	Alternative method 2			
	<i>a</i> = 2	B1		
	$x^{2}-bx-bx+b^{2} or$ $x^{2}-2bx+b^{2} or$ $-2ab = -6 or$ $-ab = -3 or$ $b = 1.5$	M1	oe	
	a = 2 and $b = 1.5$ and $c = 0.5$	A1	oe eg $2(x-1.5)^2+0.5$	

Page 14 Version 1.0

Q	Answer	Mark	Comments			
	Alternative method 1					
	their $2(x-1.5)^2 = 8.5$ - their 0.5	M1				
	their $(x - 1.5) = \pm \sqrt{\frac{8.5 - \text{their } 0.5}{2}}$	M1dep	oe			
	3.5 and -0.5	A1	oe			
	Alternative method 2					
25(b)	$2x^{2} - 6x - 3.5 (= 0)$ or $4x^{2} - 12x - 7 (= 0)$	M1	oe 3-term quadratic equation or expression			
	Correct use of quadratic formula $eg = \frac{12 \pm \sqrt{(-12)^2 - 4 \times 4 \times -7}}{2 \times 4}$ or correct factorisation $eg = (2x - 7)(2x + 1) = 0$	M1dep	oe			
	3.5 and -0.5	A1	oe			
26	144% or 1.44 seen	B1				
	$\sqrt{1.44}$ or 1.2	M1	oe			
	their 1.2 × 32	M1dep				
	38.4	A1				



Copyright © 2015 AQA and its licensors. All rights reserved.

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.