

# GCSE MATHEMATICS 8300/3H

Higher Tier Paper 3 Calculator

Mark scheme

June 2023

Version: 1.0 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aga.org.uk

#### Copyright information

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2023 AQA and its licensors. All rights reserved.

# **Glossary for Mark Schemes**

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

М	Method marks are awarded for a correct method which could lead to a correct answer.
Α	Accuracy marks are awarded when following on from a correct method. It is not necessary to always see the method. This can be implied.
В	Marks awarded independent of method.
ft	Follow through marks. Marks awarded for correct working following a mistake in an earlier step.
SC	Special case. Marks awarded for a common misinterpretation which has some mathematical worth.
M dep	A method mark dependent on a previous method mark being awarded.
B dep	A mark that can only be awarded if a previous independent mark has been awarded.
oe	Or equivalent. Accept answers that are equivalent. eg accept 0.5 as well as $\frac{1}{2}$
[a, b]	Accept values between a and b inclusive.
[a, b)	Accept values a
3.14	Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416
Use of brackets	It is not necessary to see the bracketed work to award the marks.

Examiners should consistently apply the following principles.

# **Diagrams**

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

# Responses which appear to come from incorrect methods

Whenever there is doubt as to whether a student has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the student. In cases where there is no doubt that the answer has come from incorrect working then the student should be penalised.

#### Questions which ask students to show working

Instructions on marking will be given but usually marks are not awarded to students who show no working.

#### Questions which do not ask students to show working

As a general principle, a correct response is awarded full marks.

#### Misread or miscopy

Students often copy values from a question incorrectly. If the examiner thinks that the student has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

#### **Further work**

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

#### Choice

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

## Work not replaced

Erased or crossed out work that is still legible should be marked.

# Work replaced

Erased or crossed out work that has been replaced is not awarded marks.

#### **Premature approximation**

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.

# **Continental notation**

Accept a comma used instead of a decimal point (for example, in measurements or currency), provided that it is clear to the examiner that the student intended it to be a decimal point.

Q	Answer	Mark	Comments
1	7	B1	

Q	Answer	Mark	Commen	its		
	$\frac{15}{8}$ or $1\frac{7}{8}$	B1	oe fraction eg $\frac{1875}{1000}$			
	Additional Guidance					
2	Ignore attempts to simplify after correct answer seen					
	Do not allow fractions with decimal numerators or denominators					
	eg $\frac{18.75}{10}$					

Q	Answer	Mark	Commen	its
3	19 – 11 or 8 or 11 – 19 or –8	M1		
	4	A1		
	Add	ditional G	Guidance	
	Answer 4 with no working or no inco	M1A1		
	Embedded answer eg $5 \times 4 + 11 = 3$	M1A0		

Q	Answer	Mark	Comments
4	4.5 × 5000 or 22500 or 5000 ÷ 100 or 50 or 4.5 ÷ 100 or 0.045	M1	
	225	A1	

Q	Answer	Mark	Comments		
	1 – 0.04 or 0.96 or 0.04 × 1000000 or 40000 or 960000	M1	oe eg 1 – $\frac{4}{100}$ 1 040 000 implies M1		
	Full method for exactly 5 compounded percentage calculations with their multiplier	M1	oe eg 1000000 × their 0.96	<b>S</b> <sup>5</sup>	
5	[800 000, 820 000] with M2 awarded	A1			
	Additional Guidance				
	815372.() or 815373 with M2 awarded				
	Answer 800 000 from 40 000 × 5				
	Answer 800 000 without either 40 000	r M2 awarded	M0M0A0		
	Intermediate values for separate calculations are 960 000, 921 600, 884 736, 849 346.()				

Q	Answer	Mark	Comments			
	No ticked		eg 2 faces are hidden			
	and		B1 No ticked			
	correct reason					
	or	<b>D</b> 0				
	correct evaluation of the surface areas for any numerical or algebraic values	B2				
	or					
	correct ratio of the surface areas					
	Ade	ditional G	Guidance			
	Ignore irrelevant reasons or evaluation evaluation, unless contradictory	ons alongs	side a correct reason or			
	"No" may be implied by a correct reas					
6	Accept reasoning that uses A as a cu					
	No ticked and					
	A has 6, B has 10 (condone sides fo	r faces)		B2		
	A has 3, B has 5			B2		
	A has 6 sides, on B each cube only h	as 5		B2		
	Ratio is 3:5 (accept equivalent ratios	)		B2		
	The bottom and the top are missing	(or covere	ed)	B2		
	When they are put together you lose	two faces	•	B2		
	You wouldn't count two sides (condo	ne sides	for faces)	B2		
	Some of the faces are covered					
	You cannot see one side because they are stacked together					
	One face covered					
	Part of the area of A is covered where it joins B					
	Both touching sides			B2		
	Yes ticked or Cannot tell ticked			В0		

Q	Answer			Mar	k		Commer	nts	
	0 and 3 in the correct positions				B1	0 or 3 in	the correc	ct po	sition
		Additional Guidance							
7(a)								]	
(4.)		x	-3	-2	-1	0	1		B2
		у	3	0	-1	0	3		DZ.

Q	Answer	Mark	Comments		
	Plots at least three points correctly	M1	correct or ft their table in (a) $\pm \frac{1}{2}$ small square points may be implied by grathrough them	ph passing	
7(b)	Correct graph drawn through the five correct points	A1	$\pm \frac{1}{2}$ small square smooth quadratic curve		
		ditional G			
	Correct graph drawn without plotting	the correc	ct points	M1A1	
	Ignore any extra points plotted				
	Ignore any part of graph drawn for $x < -3$ or $x > 1$				
	Ruled straight lines			A0	

Q	Answer	Mark	Comments			
	Alternative method 1					
	2450 ÷ (2 + 5) or 2450 ÷ 7 or 350	M1	oe			
	their $350 \times 5$ or $1750$ or their $350 \times 2$ or $700$ or their $350 \div 4$ or $87.5(0)$	M1dep	oe $2450 \times \frac{5}{7}$ is M2 $2450 \times \frac{2}{7}$ is M2 $2450 \div 28$ is M2			
8	their 1750 $\div$ 4 or $(2450 - \text{their } 700) \div$ 4 or their 87.5(0) $\times$ 5 or 437.5(0)	M1dep	oe dep on M2 $350 \times \frac{5}{4} \text{ is M3}$			
	437.5(0) and Yes	A1	accept 437.5(0) > 430			
	Alternative method 2					
	2450 ÷ 4 or 612.5(0)	M1	oe			
	their $612.5(0) \div (2+5)$ or their $612.5(0) \div 7$ or $87.5(0)$	M1dep	oe 2450 ÷ 28 is M2			
	their $87.5(0) \times 5$ or their $612.5(0)$ – their $87.5(0) \times 2$ or 437.5(0)	M1dep	oe dep on M2 $612.5(0) \times \frac{5}{7}$ is M3			
	437.5(0) and Yes	A1	accept 437.5(0) > 430			

Mark scheme and Additional Guidance continue on the next page

	Alternative method 3					
	430 × 4 or 1720	M1				
	2450 ÷ (2 + 5) or 2450 ÷ 7 or 350	M1	oe			
	their 350 × 5 or 1750 or their 350 × 2 or 700	M1dep	oe dep on 2nd M $2450 \times \frac{5}{7} \text{ is M2}$ $2450 \times \frac{2}{7} \text{ is M2}$			
	1720 and 1750 and Yes	A1	2450 - 1720 = 730 and 700	and Yes		
	Alternative method 4					
8	430 × 4 or 1720	M1				
cont	their 1720 ÷ 5 or 344 or their 1720 × 2 or 3440	M1dep	oe			
	their 344 × 2 or their 3440 ÷ 5 or 688	M1dep	oe dep on M2 $1720 \times \frac{2}{5} \text{ is M3}$			
	2408 and Yes	A1				
	Additional Guidance					
	Up to M3 may be awarded for correct answer, even if this is seen amongst					
	2450 ÷ 7 × 1.25 or 350 × 1.25	M1M1M1				
	Yes may be implied eg They receive 7.50 more than 430	M3A1				
	Condone £437.50p and Yes			M3A1		

Q	Answer	Mark	Comments		
	80 – 25 or 55 or 360 – 80 – 25 or 255	M1	oe implied by 1 degree = 2.4 people or 5 degrees = 12 people		
9	$\frac{132}{\text{their } 55} \times 360 \text{ or } 864$ or $\frac{132}{\text{their } 55} \times 80 \text{ or } 192$ or $\frac{132}{\text{their } 55} \times 25 \text{ or } 60$ or $\frac{132}{\text{their } 55} \times \text{their } 255$ or $\frac{132}{\text{their } 55} \times (80 + 25) \text{ or } 252$ or their $255 \div \frac{\text{their } 55}{132}$	M1dep	oe 2.4 × their 255 is M2 12 × 51 is M2 2.4 × 105 is M2		
	612	A1			
	Additional Guidance				
	Up to M2 may be awarded for correct answer, even if this is seen amongst				

Q	Answer	Mark	Commen	ts
	Alternative method 1 – using tange	ent of an	angle	
	tan chosen or used	M1		
	$\tan 58 = \frac{x}{46}$ or $46 \times \tan 58$	M1dep	oe	
	or $\tan 32 = \frac{46}{x}$ or $\frac{46}{\tan 32}$			
	[73.6, 74]	A1		
	Alternative method 2 – finding hyp	otenuse	first	
10	$\frac{46}{\cos 58}$ or $\frac{46}{\sin 32}$ or 86.8() or 87	M1	oe	
10	$\sqrt{(\text{their }86.8())^2 - 46^2}$ or $\sqrt{5418.()}$ or their $86.8() \times \sin 58$ or their $86.8() \times \cos 32$	M1dep	oe	
	[73.6, 74]	A1		
	Ad	ditional G	Guidance	
	Do not accept scale drawing			
	Answer 73 after answer in range see	n		M1M1A1
	$\frac{\sin 32}{46} = \frac{\sin 58}{x}$			M1

Q	Answer	Mark	Comments	
	8 or 10 M1 8 may be implied by 2 <sup>2</sup> or 4		1	
	8 and 10		8 may be implied by 2 <sup>2</sup> or <sup>2</sup>	1
	and	A1	accept 0.03 with $\frac{1}{40}$ or 0.0	25 seen
	Additional Guidance			
11(a)	Do not allow exact calculations for M1A1			
	eg 4.113 = 4 and 10.21 = 10 and $\frac{1}{40}$			
	1 or 0.025 without 9 or 10 open (9 may be implied by 2 <sup>2</sup> or 4)			M1A0
				M0A0

Q	Answer	Mark	Comments		
	Valid explanation	rounded			
	Additional Guidance				
	Ignore irrelevant reasons alongside a correct reason, unless contradictory				
	Ignore a calculation using exact value	es alongsi	de a correct reason		
	eg 0.025 is greater than 0.0238 a	nd both r	numbers rounded down	B1	
	0.025 is greater than 0.0238			В0	
	The denominator is smaller	B1			
	The denominator using the exact values is bigger			B1	
11(b)	(Decimals) rounded down				
11(2)	Because 8.34 is more than 8 and 10.21 is more than 10  One is divided by less (with answer more)  Estimating rounds the numbers down which makes the denominator less  B				
	Estimating rounds the numbers down	n which m	akes it less	В0	
	Because it rounds up			В0	
	Because she rounded each number to one significant figure				
	The numbers get rounded up so mor	e than the	exact value	В0	
	Rounded up when estimating			В0	
	Removing the decimals makes the ne	umber big	ger	В0	

Q	Answer	Mark	Comments	
	Ben and valid reason	B1	eg spun the most times	
12(a)	Additional Guidance			
	Do not accept an incorrect reason alongside a correct response			
	Do not accept reasons which refer to the probability increasing			
	Ignore reasons that refer to results be	Ignore reasons that refer to results being more accurate		

Q	Answer	Mark	Comments	
	Valid reason	B1	eg 14.8 is not a whole numb	er
	Ade	ditional G	Guidance	
	Do not accept an incorrect reason ald	ongside a	correct response	
	0.185 × 80 is not a whole number			B1
	Number of spins would be a decimal			
40/5)	Number of spins must be a whole number			
12(b)	Cannot land on the spinner 14.8 times B1			
	Have to spin 14.8 times B0			
	$0.185 \times 80 = 14.8$			
	14.8 B0			
	It is a decimal B0			
	Must be a whole number			В0

Q	Answer	Mark	Comments
12(c)	125 × 0.32 or 40 or	M1	oe
12(0)	1 – 0.32 or 0.68		
	85	A1	

Q	Answer	Mark	Comments	
	176 ÷ 48 or 3.66 or 3.67 or $\frac{11}{3}$ or 3h 40 mins	M1	oe eg 220 mins implied by 12 40 pm	
	$(293 - 176) \div 65$ or $117 \div 65$ or $1.8$ or $\frac{9}{5}$ or $1 \text{ h } 48 \text{ mins}$	M1	oe eg 108 mins	
	their 3.66 + their 1.8 or $\frac{82}{15}$ or [5.46, 5.47] or 5 h 28 mins or [2 27 (pm), 2 28.2 (pm)]	M1dep	oe eg 328 mins dep on M2  implied by adding times eg 9 + 3 40 + 1 48  oe arrival time	
13	or 5h 30 mins and 5h 28 mins and Yes or 330 mins and 328 mins and Yes or [2 27 (pm), 2 28.2 (pm)] and Yes	A1	must be in a comparable tim	ne format
	Ad	ditional G	Guidance	
	Up to M3 may be awarded for correct not subsequently used	t work see	en in multiple attempts even if	
	Accept use of 24 hour clock throughout	out		
	Do not accept 2 28 am as a correct a	rrival time		
	$\frac{176}{48} = 3.6$ , $\frac{117}{65} = 1.8$ , $3.6 + 1.8 = 5$	.4, 2 24 p	m and Yes	M1M1M1A0
	$\frac{176}{48} = 3.7$ , $\frac{117}{65} = 1.8$ , $3.7 + 1.8 = 5$	.5, 2 30 p	m and Arrives on time	M1M1M1A0
	3.6 + 1.8 = 5.4, 2 24 pm and Yes			M0M1M0A0
	3.7 + 1.8 = 5.5, 2 30 pm and Arrives	on time		M0M1M0A0

Q	Answer	Mark	Comments	
	5186 ÷ 0.2 or 5186 × 5 or 25930	M1	oe	
	38 500	A1		
	(their 38 500 – 9880) × 0.1325		their 38 500 must be > 988	0
	or 28620 × 0.1325	M1	full method to calculate Nat Insurance	tional
	3792(.15) A1ft ft their 38 500, which must be			
14	Additional Guidance  Accept final answer rounded or truncated to the nearest pound if a more accurate value is seen in working  Do not accept '13.25% of 28 620' or 13.25% × 28 620 for M mark unless accompanied by a correct method or value  (25 930 – 9880) × 0.1325 = 2126.62 or 2126.63  M1A0M1A1			
				M1A0M1A1ft
	25 930 × 0.1325 or 3435.72 or 343	35.73		M1A0M0A0ft

Q	Answer	Mark	Comments
	20 × 0.8 or 16 or 20 × 1.8 or 36 or 40 × 1.2 or 48 or 40 × 0.7 or 28 or 60 × 0.4 or 24	M1	one correct area calculation or frequency value may be on diagram
15(a)	$20 \times 0.8 + 20 \times 1.8 + 40 \times 1.2 +$ $40 \times 0.7 + 60 \times 0.4$ or $16 + 36 + 48 + 28 + 24$ or $152$	M1dep	allow 1 error or 1 omission or 1 misread of a frequency density value
	28	A1	

Q	Answer	Mark	Comments	
	Rectangular box plot with whiskers to 5 and 23	B1		
	Lower quartile drawn at 11 and median drawn at 18	B1		
	Upper quartile drawn at 20		correct or ft their lower quart	ile + 9
		B1ft	must be the vertical line at ri their box	ght side of
	Add	ditional G	Guidance	
	Mark intention eg any height and allo	w horizon	tal line through centre of box	
	Allow ends of whiskers to be vertical lines of any length, dots, crosses or stops			
15(b)	b) $\pm \frac{1}{2}$ small square tolerance			
	Median must be the second vertical line of a box with three vertical lines			
	Only vertical lines or points plotted			В0
	0 5 10 Distance	15 run (miles	20 25 s)	В3

Q	Answer	Mark	Comments	3	
	Alternative method 1 – using Pytha	agoras' th	neorem or 3, 4, 5 triangle		
	$16 \div 4 \times 5$ or $20$ (cm) or identifies triangle as 3, 4, 5	M1	oe length of $c$ may be on diagram		
	$\sqrt{(\text{their } 20)^2 - 16^2}$ or $\sqrt{400 - 256}$ or $\sqrt{144}$ or $4 \times 3$	M1dep			
	12 (cm)	A1	length of $b$ may be on diagram		
	96	A1ft	ft $\frac{1}{2} \times 16 \times \text{their } 12 \text{ with}$	M2 awarded	
16	Alternative method 2 – using trigonometry and ½ab sin C formula				
	16 ÷ 4 × 5 or 20 (cm)	M1	oe length of $\emph{c}$ may be on diagram		
	$\cos^{-1}\left(\frac{16}{20}\right)$ or 36.8() or 36.9	M1dep	angle between sides $a$ and	d c	
	$\frac{1}{2} \times 16 \times 20 \times \sin (\text{their } 36.8())$	M1dep	dep on M2		
	96	A1			
	Ac	ditional (	Guidance		
	$\frac{1}{2} \times 16 \times 12 \times \sin 90$			M1M1M1	

Q	Answer	Mark	Comments	
	Alternative method 1 – multiplies t	hrough b	y 10 or common denominator of 10	
	5(x+8) + 2(9-x) or 5x + 40 + 18 - 2x	M1	oe numerator on the left-hand side if written as a fraction allow one error or omission in the expansion if brackets not seen eg $5x + 18 - 2x$	
	3x + 58	A1	may be implied by eg $3x + 18 = 0$ or $3x = -18$	
17	their $(3x + 58) = 4 \times$ (their 10) or their $(3x + 58) = 40$ or $3x + 18 = 0$ or $3x = -18$	M1	oe allow an unsimplified expression for their $(3x + 58)$ equation may be implied by answer	
17	-6	A1ft	ft M1A0M1	
	Alternative method 2 – collects terms with fractions			
	$\frac{x}{2} + 4 + \frac{9}{5} - \frac{x}{5}$	M1	oe eg $0.5x + 4 + 1.8 - 0.2x$ allow one error	
	$\frac{3}{10}x + \frac{29}{5}$	A1	oe eg 0.3x + 5.8	
	$\frac{3}{10}x = \frac{20}{5} - \frac{29}{5}$ or $\frac{3}{10}x = -\frac{9}{5}$	M1	oe eg $0.3x = -1.8$ terms must be collected	
	-6	A1ft	ft M1A0M1	

# Additional Guidance is on the next page

	Additional Guidance	
	Accept decimal answers for follow through correct to 1 dp or better	
	Apply the principles of alt 1 for any use of other common denominators eg common denominator of 20 (or multiplication through by 20) $10(x+8) + 4(9-x) = 6x + 116$ $6x + 116 = 80 \qquad x = -6$	M1A1 M1A1
	An incorrect simplification of $5x + 40 + 18 - 2x$ may still gain the third and fourth marks	
	eg $5x + 40 + 18 - 2x = 3x + 68$ followed by $3x + 68 = 40$ and $x = -\frac{28}{3}$	M1A0M1 A1ft
17	eg $5x + 40 + 18 - 2x = 2x + 68$ followed by $2x + 68 = 40$ and $x = -14$	M1A0M1 A1ft
cont	An incorrect denominator may still gain the third and fourth marks $\frac{5x+40+18-2x}{7}$ followed by $5x+40+18-2x=28$ and $x=-10$	M1A0M1 A1ft
	Denominator not processed	
	3x + 58 = 4 followed by $3x = -54$ and $x = -18$	M1A1M0A0
	(x+8)+(9-x)=40	M0A0M1A0
	Two errors in the expansion but with brackets seen may go on to get the third and fourth marks	
	5(x+8)+2(9-x)=5x+8+18-x	1st M1A0
	Two errors in the expansion and no brackets seen, no follow through allowed	
	$5x + 8 + 18 - x$ followed by $4x + 26 = 40$ and $x = \frac{14}{4}$	M0A0M1A0

Q	Answer	Mark	Comments
	$(2x+4)^2+6(2x+4)$	M1	may be seen in a grid
	$4x^2 + 8x + 8x + 16 + 12x + 24$ or $4x^2 + 16x + 16 + 12x + 24$	M1don	fully expanded expression with terms summed
	014x + 10x + 10 + 12x + 24	M1dep	allow one omission or one arithmetic error
18(a)	$4x^2 + 8x + 8x + 16 + 12x + 24$ or $4x^2 + 16x + 16 + 12x + 24$ and	A1	
	$4x^2 + 28x + 40$ Additional Guidance		
	$4x^2 + 16 + 12x + 24$ is two errors		

Q	Answer	Mark	Comments	
	$4x^2 + 28x + 45 (= 0)$	M1	must be correct	
18(b)	$(2x+5)(2x+9) (= 0)$ or $(2x+7)^2 - 49 + 45 (= 0)$ or $\frac{-28 \pm \sqrt{28^2 - 4 \times 4 \times 45}}{2 \times 4}$ or $\frac{-28 \pm \sqrt{64}}{8}$ or $\frac{-28 \pm 8}{8}$ or $\frac{-7 \pm \sqrt{4}}{2}$	M1dep	oe implies first M1	
	(x =) -2.5 and $(x =) -4.5$	A1	oe fraction or decimal SC2 (x =) [-1.63, -1.629] a (x =) [-5.371, -5.37]	and
	Additional Guidance			
	SC2 from using $4x^2 + 28x + 35 (= 0)$			
	Trial and improvement with both answers correct and chosen from any list			M1M1A1
	Trial and improvement with one answer correct			M0M0A0

Q	Answer	Mark	Comments	
	Creates an algebraic product in the form $(x + a)(x + b)$ where there is a difference of 6 between $a$ and $b$	M1	accept any letter for $x$ eg $x(x+6)$ or $x^2+6x$ or $x(x-6)$ or $x^2-6x$	
	Correctly expands their product, adds 9 and simplifies to a quadratic expression	M1dep	eg $x^2 + 6x + 9$ or $x^2 - 6x + 9$	)
	Correctly factorises their quadratic expression to the form $(x + c)^2$ with M2 awarded	A1	eg $(x+3)^2$ or $(x-3)^2$	
19	Additional Guidance			
	Trialling integers scores no marks, but alongside correct algebra	any testing of values		
	Ignore any further work or attempts to	o solve aft	ter correct answer seen	
	Missing brackets may be recovered e	$eg x \times x +$	6 followed by $x^2 + 6x + 9$	M1M1
$(x+3)(x+3)$ without $(x+3)^2$ seen does not score the A mark			score the A mark	
	(x-2)(x-8)			M1
	$x^2 - 2x - 8x + 16 + 9 = x^2 - 10x + 25$			M1
	$(x-5)^2$			A1

Q	Answer	Mark	Comments	
	Substitutes a correct pair of coordinates and states that the equation is correct	B1	eg $18 = \frac{36}{2}$ so he is right	
	Ade	ditional G	Guidance	
	Accept 'Yes' or a tick or any clear ind	ication th	at he is correct	
	Do not accept pairs of values not on the graph  Do not accept a correct answer alongside an incorrect response unless clearly chosen			
20(a)	Do not accept a coordinate with no si	ubstitution	seen	
	Pairs with integer $x$ or $y$ include			
	$18 = \frac{36}{2}, \ 15 = \frac{36}{2.4}, \ 12 = \frac{36}{3}, \ 10 = \frac{36}{3.6}$ $9 = \frac{36}{4},  8 = \frac{36}{4.5}, \ 7.2 = \frac{36}{5},  6 = \frac{36}{6}$			
	Substituting values incorrectly			
	eg $2 = \frac{36}{18}$ or $4 = \frac{36}{9}$			В0

Q	Answer	Mark	Comments	
	Alternative method 1			
	$G \propto \sqrt{H}$ or $G = k\sqrt{H}$ or $16 \div 2 \times 3 = k\sqrt{16}$ or $24 = k\sqrt{16}$ $k = \frac{\text{their } 24}{\sqrt{16}} \text{ or } k = 6$	M1	oe equation $k$ may be any letter their 24 must be the result	of 16 ÷ 2 × 3
	or $G = \text{their } 6\sqrt{H}$			
	their 6 × their $\sqrt{100}$ or 60	M1dep	dep on M2	
	60:100 or 3:5	A1	oe ratio	
	Alternative method 2			
20(b)	100 ÷ 16 or 6.25	M1		
	$\sqrt{\text{their 6.25}}$ or 2.5	M1dep		
	2 × their 2.5 or 5 or 24 × their 2.5 or 60	M1dep	dep on M2	
	60:100 or 3:5	A1	oe ratio	
	Additional Guidance			
	Ignore an incorrect attempt to simplify a correct ratio eg 60 : 100 followed by 3 : 4			M1M1M1A1
	k = 6 implies M2 unless from incorrect working			
	$G lpha k \sqrt{H}$ is M0 unless recovered			
	$G = k\sqrt{H}$ $\sqrt{16} = 4$ $G: H = 6: 4$ $6 = k \times 4$ $k = \frac{6}{4}$ followed by $G = 1.5 \times 10$ 150:100			M1M0M0A0
	G = 24 with no correct further work			МО

Q	Answer	Mark	Comments
21	72 (-) 6 or 66 or 63 (-) 6 or 57 or 45 (+) 21 or 66 or 36 (+) 21 or 57 or 56 (+) 10 or 66 or 49 (+) 8 or 57	M1	large rectangle subtract missing rectangle, implied by volumes of 864 and 72  splits side elevation vertically, implied by volumes of 540 and 252  splits side elevation horizontally, implied by volumes of 672 and 120  oe
	792 or 165 A1	A1	may be on diagram
	Maximum 792 and Minimum 165	A1	

Q	Answer	Mark	Commen	ts
	Enlargement	B1	accept Enlarge	
	(Scale factor) $-\frac{1}{2}$	B1	oe	
	(Centre) (7, 4)	B1	oe	
	Ado	ditional G	Guidance	
	Do not accept reduces, gets smaller,			
22	Do not accept $\div -\frac{1}{2}$ or $\div -2$ for sca			
	Ignore missing brackets on 7, 4			
	Do not accept $\binom{7}{4}$ for centre of enlargement, however this does not imply a combined transformation			
	Enlarge, $-\frac{1}{2}$ , (7, 4)			B1B1B1
	Combined transformation			B0B0B0

Q	Answer	Mark	Comments	
	$35^2 + 65^2 - 2 \times 35 \times 65 \times \cos 100$	M1	oe valid trigonometric meth	od used
23(a)	$\sqrt{35^2 + 65^2 - 2 \times 35 \times 65 \times \cos 100}$ = 78.9() or $\sqrt{6240.(0992)} = 78.9()$	A1	CA = 78.99429858	
	Additional Guidance			
	Using sine rule with $CA = 79$ to obtain $AB$ or $BC$		M0A0	

Q	Answer	Mark	Comments	
	Alternative method 1 – sine rule to find ACB			
	$\frac{\sin ACB}{35} = \frac{\sin 100}{79}$	M1	oe 79 may be 78.9()	
	$\sin ACB = 35 \times \frac{\sin 100}{79}$ or $\sin ACB = 35 \times 0.0124$ or $\sin ACB = 0.436$	M1dep	oe	
	<i>ACB</i> = [25.8, 26]	A1		
	234.()	A1ft	ft 360 - 100 - their ACB with M2 scored	
23(b)	Alternative method 2 – cosine rule to find ACB			
	$35^2 = 79^2 + 65^2 - 2 \times 79 \times 65 \times \cos$ <i>ACB</i>	M1	oe 79 may be 78.9()	
	$\cos ACB = \frac{79^2 + 65^2 - 35^2}{2 \times 79 \times 65}$			
	or $\cos ACB = \frac{9241}{10270}$	M1dep		
	or cos <i>ACB</i> = 0.899			
	ACB = [25.8, 26]	A1		
	234.()	A1ft	ft 360 – 100 – their ACB with M2 scored	

Mark scheme and Additional Guidance continue on the next page

	Alternative method 3 – sine rule to	find BAC	;
	$\frac{\sin BAC}{65} = \frac{\sin 100}{79}$	M1	oe 79 may be 78.9()
	$\sin BAC = 65 \times \frac{\sin 100}{79}$ or $\sin BAC = 65 \times 0.0124$ or $\sin BAC = 0.81(0)$	M1dep	oe
	BAC = [54.1, 54.3]	A1	
	234.()	A1ft	ft their BAC + 180 with M2 scored
	Alternative method 4 – cosine rule	to find B	AC
23(b)	$65^2 = 79^2 + 35^2 - 2 \times 79 \times 35 \times \cos$ <i>BAC</i>	M1	oe 79 may be 78.9()
cont	$\cos BAC = \frac{79^2 + 35^2 - 65^2}{2 \times 79 \times 35}$ or $\cos BAC = \frac{3241}{5530}$ or $\cos BAC = 0.586$	M1dep	
	BAC = [54.1, 54.3]	A1	
	234.()	A1ft	ft their BAC + 180 with M2 scored
	Ade	ditional G	duidance
	CA = 79 is given in part (a) or 78.9() can be used. There is no follow through from part (a).		
	Accept any notation for the angle eg $\sin x$ or $\sin C$ for angle $ACB$		
	Correct work for part (b) seen in part (a) may be awarded method marks in part (b)		