

Write your name here

Surname

Other names

**Edexcel**

**International GCSE**

Centre Number

--	--	--	--	--	--

Candidate Number

--	--	--	--	--	--

# Chemistry

**Unit: 4CH0**

**Paper: 2C**

Friday 20 January 2012 – Morning

**Time: 1 hour**

Paper Reference

**4CH0/2C**

**You must have:**

Ruler

Calculator.

Total Marks

## Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- Show all the steps in any calculations and state the units.

## Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

## Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P40127A

©2012 Pearson Education Ltd.

1/1/1/



P 4 0 1 2 7 A 0 1 1 6

**PEARSON**

# THE PERIODIC TABLE

0

7

6

5

4

3

Group

2

1

Period

4	He	Helium	2
---	----	--------	---

1	H	Hydrogen	1
---	---	----------	---

20	Ne	Neon	10
19	F	Fluorine	9
16	O	Oxygen	8
14	N	Nitrogen	7
12	C	Carbon	6
11	B	Boron	5
40	Ca	Calcium	20
39	K	Potassium	19
38	Sr	Strontium	38
37	Rb	Rubidium	37
36	Kr	Krypton	36
35	Br	Bromine	35
34	Se	Selenium	34
33	As	Arsenic	33
32	Ge	Germanium	32
31	Ga	Gallium	31
30	Zn	Zinc	30
29	Cu	Copper	29
28	Ni	Nickel	28
27	Co	Cobalt	27
26	Fe	Iron	26
25	Mn	Manganese	25
24	Cr	Chromium	24
23	V	Vanadium	23
22	Ti	Titanium	22
21	Sc	Scandium	21
18	Ar	Argon	18
17	Cl	Chlorine	17
16	S	Sulfur	16
15	P	Phosphorus	15
14	N	Nitrogen	14
13	Al	Aluminium	13
12	Si	Silicon	12
11	B	Boron	11
84	Kr	Krypton	84
83	Br	Bromine	83
82	Se	Selenium	82
81	As	Arsenic	81
80	Ge	Germanium	80
79	Ga	Gallium	79
78	Zn	Zinc	78
77	Cu	Copper	77
76	Ni	Nickel	76
75	Co	Cobalt	75
74	Fe	Iron	74
73	Mn	Manganese	73
72	Cr	Chromium	72
71	V	Vanadium	71
70	Ti	Titanium	70
69	Sc	Scandium	69
54	Xe	Xenon	54
53	I	Iodine	53
52	Te	Tellurium	52
51	Sb	Antimony	51
50	Sn	Tin	50
49	In	Indium	49
48	Cd	Cadmium	48
47	Ag	Silver	47
46	Pd	Palladium	46
45	Rh	Rhodium	45
44	Ru	Ruthenium	44
43	Tc	Technetium	43
42	Mo	Molybdenum	42
41	Nb	Niobium	41
40	Zr	Zirconium	40
39	Y	Yttrium	39
38	Sr	Strontium	38
37	Rb	Rubidium	37
36	Kr	Krypton	36
35	Br	Bromine	35
34	Se	Selenium	34
33	As	Arsenic	33
32	Ge	Germanium	32
31	Ga	Gallium	31
30	Zn	Zinc	30
29	Cu	Copper	29
28	Ni	Nickel	28
27	Co	Cobalt	27
26	Fe	Iron	26
25	Mn	Manganese	25
24	Cr	Chromium	24
23	V	Vanadium	23
22	Ti	Titanium	22
21	Sc	Scandium	21
20	Ca	Calcium	20
19	K	Potassium	19
18	Ar	Argon	18
17	Cl	Chlorine	17
16	S	Sulfur	16
15	P	Phosphorus	15
14	N	Nitrogen	14
13	Al	Aluminium	13
12	Si	Silicon	12
11	B	Boron	11
10	Ne	Neon	10
9	F	Fluorine	9
8	O	Oxygen	8
7	N	Nitrogen	7
6	C	Carbon	6
5	B	Boron	5
4	Be	Beryllium	4
3	Li	Lithium	3
2	He	Helium	2
88	Ra	Radium	88
87	Fr	Francium	87
86	Rn	Radon	86
85	At	Astatine	85
84	Po	Polonium	84
83	Bi	Bismuth	83
82	Pb	Lead	82
81	Tl	Thallium	81
80	Hg	Mercury	80
79	Au	Gold	79
78	Pt	Platinum	78
77	Ir	Iridium	77
76	Os	Osmium	76
75	Rh	Rhenium	75
74	W	Tungsten	74
73	Ta	Tantalum	73
72	Hf	Hafnium	72
71	Ta	Tantalum	71
70	Nb	Niobium	70
69	Zr	Zirconium	69
68	Y	Yttrium	68
67	La	Lanthanum	67
66	Ba	Barium	66
65	Cs	Caesium	65
64	La	Lanthanum	64
63	Pr	Praseodymium	63
62	Nd	Niodymium	62
61	Pm	Promethium	61
60	Sm	Samarium	60
59	Eu	Europium	59
58	Gd	Gadolinium	58
57	Tm	Thulium	57
56	Dy	Dysprosium	56
55	Ho	Holmium	55
54	Er	Erbium	54
53	Tm	Thulium	53
52	Yb	Ytterbium	52
51	Lu	Lutetium	51
50	Hf	Hafnium	50
49	Ta	Tantalum	49
48	W	Tungsten	48
47	Re	Rhenium	47
46	Os	Osmium	46
45	Ir	Iridium	45
44	Pt	Platinum	44
43	Au	Gold	43
42	Hg	Mercury	42
41	Tl	Thallium	41
40	Pb	Lead	40
39	Bi	Bismuth	39
38	Po	Polonium	38
37	At	Astatine	37
36	Rn	Radon	36
35	Fr	Francium	35
34	Ra	Radium	34
33	Ac	Actinium	33
32	Th	Thorium	32
31	Pa	Protactinium	31
30	U	Uranium	30
29	Np	Neptunium	29
28	Pu	Plutonium	28
27	Am	Americium	27
26	Cm	Curium	26
25	Bk	Berkelium	25
24	Cf	Californium	24
23	Es	Einsteinium	23
22	Fm	Fermium	22
21	Mn	Mendelevium	21
20	Nv	Nobelium	20
19	Lr	Lutetium	19
18	Lu	Lutetium	18
17	Yb	Ytterbium	17
16	Tm	Thulium	16
15	Er	Erbium	15
14	Dy	Dysprosium	14
13	Ho	Holmium	13
12	Er	Erbium	12
11	Tm	Thulium	11
10	Yb	Ytterbium	10
9	Lu	Lutetium	9
8	Hf	Hafnium	8
7	Ta	Tantalum	7
6	W	Tungsten	6
5	Re	Rhenium	5
4	Os	Osmium	4
3	Ir	Iridium	3
2	Pt	Platinum	2
1	Au	Gold	1
0	Hg	Mercury	0

Key

Relative atomic mass
Symbol
Name
Atomic number



**Answer ALL questions.**

- 1 (a) Complete the table to show the relative mass and relative charge of a proton, a neutron and an electron.

(4)

	Proton	Neutron	Electron
Relative mass			1/1840
Relative charge	+ 1		

- (b) The symbol for an atom of one isotope of hydrogen is  ${}^3_1\text{H}$

- (i) State the number of protons, neutrons and electrons present in one atom of this isotope.

(2)

Number of protons .....

Number of neutrons .....

Number of electrons .....

- (ii) What is meant by the term **isotopes**?

(2)

.....

.....

.....

- (c) Bromine has two naturally-occurring isotopes with mass numbers 79 and 81.  
A sample of bromine contained the two isotopes in the following proportions:

$$\text{bromine-79} = 50.7\% \quad \text{and} \quad \text{bromine-81} = 49.3\%$$

Use this information to calculate the relative atomic mass of bromine.  
Give your answer to **two** decimal places.

(2)

**(Total for Question 1 = 10 marks)**



2 Use the Periodic Table on page 2 to help you answer this question.

(a) Part of the Periodic Table is shown.

																	<b>A</b>	
	<b>E</b>												<b>D</b>					
<b>B</b>																	<b>C</b>	

In each part of this question, place a cross (☒) in **one** box to identify the letter, **A** to **E**, that represents

(i) a metal that reacts violently with water (1)

**A**      **B**      **C**      **D**      **E**  
               

(ii) a noble gas (1)

**A**      **B**      **C**      **D**      **E**  
               

(iii) a Group 2 metal (1)

**A**      **B**      **C**      **D**      **E**  
               

(iv) a halogen (1)

**A**      **B**      **C**      **D**      **E**  
               



(b) Complete these sentences by placing a cross (☒) in **one** box next to the correct answer.

(i) The elements in the Periodic Table are arranged in order of increasing (1)

- number of neutrons
- atomic number
- relative atomic mass
- mass number

(ii) Elements in the same group in the Periodic Table have the same number of (1)

- electrons in the outer shell
- protons in the nucleus
- neutrons in the nucleus
- atoms

**(Total for Question 2 = 6 marks)**

---



3 Lead(II) sulfate,  $\text{PbSO}_4$ , is an insoluble salt.

It can be made as a precipitate from a solution of lead(II) nitrate,  $\text{Pb}(\text{NO}_3)_2$

(a) (i) Identify a substance that could be added to lead(II) nitrate solution to form a precipitate of lead(II) sulfate.

(1)

(ii) Write a chemical equation for the reaction between lead(II) nitrate and the substance you identified in (a)(i).


(2)

(iii) Outline how you would produce a pure, dry sample of lead(II) sulfate from the reaction mixture in (a)(ii).

(3)

(b) A solution of lead(II) nitrate can be made by reacting solid lead(II) carbonate with dilute nitric acid.

The equation for this reaction is:



State **two** observations you would make when dilute nitric acid is added to solid lead(II) carbonate.

(2)

1

2

(Total for Question 3 = 8 marks)




4 When magnesium is burned in air, it reacts with oxygen,  $O_2$ , to form magnesium oxide,  $MgO$

A class of students investigated the relationship between the mass of magnesium burned and the mass of magnesium oxide formed.

Each student was given a different mass of clean magnesium to heat.

The students used the following method.

- Weigh a crucible and lid
- Place the magnesium ribbon in the crucible, replace the lid, and reweigh
- Heat the crucible as shown in the diagram until the magnesium burns



- Lift the lid from time to time until there is no sign of further reaction
- Allow the crucible and lid to cool and reweigh
- Repeat the heating, cooling and reweighing until two consecutive masses are the same
- Calculate the mass of magnesium oxide formed

(a) (i) Why is it necessary to lift the lid from time to time while heating?

(1)

(ii) Why is it necessary to repeat the heating until two consecutive masses are the same?

(1)



(b) Show how the mass of magnesium oxide formed can be calculated from the readings obtained. (1)

.....

.....

.....

(c) The results of each experiment are given in the table.


Mass of magnesium in g	Mass of magnesium oxide in g
0.24	0.40
0.26	0.64
0.42	0.70
0.62	1.04
0.70	1.20
0.80	1.33





(i) Plot the results on the grid and draw a straight line of best fit.

(3)



(ii) Draw a circle around the anomalous result.

(1)


(iii) Use your graph to find the mass of magnesium oxide formed when 0.48 g of magnesium is burned.

(1)

(Total for Question 4 = 8 marks)



5 The apparatus shown can be used to electrolyse aqueous sodium chloride in the laboratory.



(a) Gases are evolved at both electrodes.

(i) Describe a chemical test to show that the gas evolved at the positive electrode is chlorine. (2)

.....

.....

.....

.....

(ii) Identify gas A. (1)

.....




(b) Some of the solution formed after the electrolysis was tested with the indicator phenolphthalein. The indicator turned pink

Explain this result.

(1)

(c) The equation for the reaction taking place at the positive electrode is:



Ten faradays (10 F) of electricity were passed through an aqueous solution of sodium chloride.

(i) Calculate the amount, in moles, of chlorine formed.

(1)

(ii) Calculate the volume of chlorine formed.

(One mole of a gas occupies 24 dm<sup>3</sup> at this temperature and pressure)

(2)

**(Total for Question 5 = 7 marks)**



6 Compound **X** is a blue, crystalline solid. It contains copper(II) ions ( $\text{Cu}^{2+}$ ), sulfate ions ( $\text{SO}_4^{2-}$ ) and water of crystallisation.

(a) A student dissolved some of compound **X** in water and then added aqueous sodium hydroxide solution. She obtained a blue precipitate.

Give the formula of the blue precipitate formed in the reaction.

(1)

(b) Another student tested a solution of compound **X** for sulfate ions using dilute hydrochloric acid, followed by a few drops of barium chloride solution. She obtained a white precipitate.

Why is the dilute hydrochloric acid necessary in this test?

(1)

(c) The empirical formula of compound **X** is  $\text{CuSO}_9\text{H}_{10}$

Write the formula of compound **X** to show its water of crystallisation.

(1)

(d) Compound **X** gives a blue-green colour in a flame test.

Outline how you would carry out a flame test.

(2)

(Total for Question 6 = 5 marks)



**BLANK PAGE**



- 7 The table shows percentage by mass of the fractions obtained from a sample of crude oil and the percentage market demand for these fractions.

Fraction	Percentage by mass in crude oil	Market demand (%)
refinery gases	3	5
gasoline	12	28
kerosene	9	20
diesel	15	25
fuel oil	51	20
bitumen	10	2

- (a) Why is the market demand for the gasoline fraction greater than that for the fuel oil fraction? (1)

.....

.....

.....

- (b) Cracking is used to make long-chain hydrocarbon molecules into shorter-chain hydrocarbon molecules.

- (i) Complete the equation to show the other hydrocarbon molecule formed when  $C_{20}H_{42}$  is cracked.

(1)



- (ii) Give the name of a catalyst used in industry to crack long-chain hydrocarbons and state a temperature at which cracking is carried out.

(2)

Catalyst .....

Temperature .....



(c) Ethene ( $C_2H_4$ ) can be produced by cracking long-chain hydrocarbon molecules obtained from crude oil. The ethene produced can then be used to make ethanol.

Ethanol can also be made by the fermentation of sugars.

(i) Give **two** advantages of making ethanol from ethene, rather than by fermentation.

(2)

1 .....

2 .....

(ii) Suggest **two** reasons why ethanol is sometimes made by fermentation, rather than from ethene.

(2)

1 .....

2 .....

**(Total for Question 7 = 8 marks)**


**TURN OVER FOR QUESTION 8**



8 Sulfur dioxide,  $\text{SO}_2$ , is used as a preservative in wine.

The sulfur dioxide content of a wine can be found by titration. A chemist found that  $25.0 \text{ cm}^3$  of a sample of wine reacted with exactly  $15.00 \text{ cm}^3$  of  $0.0010 \text{ mol/dm}^3$  aqueous iodine,  $\text{I}_2(\text{aq})$ .

The equation for the reaction is



(a) Calculate the amount, in moles, of iodine in  $15.00 \text{ cm}^3$  of a  $0.0010 \text{ mol/dm}^3$  solution.

(2)

(b) Deduce the amount, in moles, of sulfur dioxide in  $25.0 \text{ cm}^3$  of the wine.

(1)

(c) Calculate the concentration, in  $\text{mol/dm}^3$ , of sulfur dioxide in the wine.

(2)

(d) Calculate the concentration, in  $\text{g/dm}^3$ , of sulfur dioxide in the wine.

(2)

(e) A concentration of sulfur dioxide that is greater than  $0.16 \text{ g/dm}^3$  makes wine unpleasant to drink.

Use the value you have calculated in (d) to state whether the wine is drinkable.

(1)

---

(Total for Question 8 = 8 marks)

---

TOTAL FOR PAPER = 60 MARKS

