

Mark Scheme (Results)

January 2015

Pearson Edexcel International GCSE in
Chemistry (4CH0) Paper 2C

Pearson Edexcel Certificate in
Chemistry (4CH0) Paper 2C

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of **learning, for all kinds of people, wherever they are in the world. We've been involved** in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2015

Publications Code UG040458

All the material in this publication is copyright

© Pearson Education Ltd 2015

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be **prepared to award zero marks if the candidate's response is not worthy of credit** according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark **scheme to a candidate's response, the** team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Accept	Reject	Marks
1 (a)	D (a molecule)			1
(b)	A (covalent)			1
(c)	NH ₃	H ₃ N		1

Total 3 marks

Question number	Answer	Accept	Reject	Marks
2 (a) (i)	(solubility/it) increases as temperature increases	positive correlation	references to proportionality	1
(ii)	(solid) B			1
(b)	<p>M1 – solid/crystals would form</p> <p>M2 – (solid A) becomes less soluble (as the solution cools) / solubility (of solid A) decreases (as temperature decreases)</p>	precipitate for solid goes cloudy		1
		reverse argument		1

Total 4 marks

Question number	Expected Answer	Accept	Reject	Marks										
3 (a)	<p>M1 P – iron ore / haematite ignore iron(III) oxide/Fe_2O_3</p> <p>M2 Q - calcium silicate</p>	slag / CaSiO_3		2										
(b)	<table border="1" data-bbox="302 467 976 775"> <thead> <tr> <th data-bbox="302 467 761 528">Type of reaction</th> <th data-bbox="761 467 976 528">Letter</th> </tr> </thead> <tbody> <tr> <td data-bbox="302 528 761 588">one that gives out heat</td> <td data-bbox="761 528 976 588">A</td> </tr> <tr> <td data-bbox="302 588 761 649">one that is a thermal decomposition</td> <td data-bbox="761 588 976 649">D ;</td> </tr> <tr> <td data-bbox="302 649 761 710">one that is a neutralisation</td> <td data-bbox="761 649 976 710">E ;</td> </tr> <tr> <td data-bbox="302 710 761 770">one that forms a poisonous gas</td> <td data-bbox="761 710 976 770">B ;</td> </tr> </tbody> </table>	Type of reaction	Letter	one that gives out heat	A	one that is a thermal decomposition	D ;	one that is a neutralisation	E ;	one that forms a poisonous gas	B ;			3
Type of reaction	Letter													
one that gives out heat	A													
one that is a thermal decomposition	D ;													
one that is a neutralisation	E ;													
one that forms a poisonous gas	B ;													
(c)	<p>M1- oxygen</p> <p>IGNORE O</p> <p>M2 – water</p>	air O_2 moisture/ H_2O		2										

(d)	<p>M1 zinc corrodes/reacts instead of iron / faster than iron</p> <p>M2 iron corrodes/reacts instead of tin / faster than tin</p> <p>lack of comparison with other metal max 1 from M1 and M2 ignore references to tin rusting</p> <p>M3 correct reference to order of reactivity of all three metals</p>	<p>zinc loses electrons/is oxidised instead of iron</p> <p>iron loses electrons/is oxidised instead of tin</p> <p>accept reverse arguments</p>	<p>zinc rusts (instead of iron)</p>	<p>3</p>
-----	--	--	-------------------------------------	----------

Total 10 marks

Question number	Answer	Accept	Reject	Marks
4(a)(i)	fermentation			1
(ii)	(to provide the) catalyst/enzyme/zymase	to increase the rate of the reaction		1
(b)(i)	M1 (test) – flame test	suitable description of flame test		2
	M2 (observation) – brick red / orange-red	red		
(ii)	copper(II) ions:	accept other suitable alkalis		5
	M1 (test) – (aqueous) sodium hydroxide / NaOH	suitable alternatives to precipitate		
	M2 (observation) – blue precipitate ignore shades of blue		all other colours	
	M2 dep on M1 or near miss of formula, eg Na(OH) ₂	(dilute) nitric acid / HNO ₃		
	sulfate ions:	(aqueous) barium nitrate / Ba(NO ₃) ₂	Reject sulfuric acid for M1 only	
	M1 (test) – (dilute) hydrochloric acid / HCl			
	M2 (test) - (aqueous) barium chloride / BaCl ₂			
	M3 (observation) – white precipitate			
	M3 dep on M2 or near miss			

Question number	Answer	Accept	Reject	Marks
4 (c)	<p>M1 (pressure) – 60-70 atm</p> <p>M2 (catalyst) – phosphoric acid / H_3PO_4 ignore references to concentration</p>	<p>any pressure or range within this range</p> <p>phosphoric(V) acid</p>	any other oxidation state	2
(d)	<p>M1 (Σ bonds broken) $348 + 412 + 360 (= 1120)$</p> <p>M2 (Σ bonds made) $612 + 463 (= 1075)$</p> <p>M3 M1 – M2 / Σ bonds broken – Σ bonds made</p> <p>M4 (+)45 (kJ/mol)</p> <p>Correct answer with no working scores 4</p> <p>– 45 (kJ/mol) scores 3</p>	<p>3231</p> <p>3186</p>		4

Total 15 marks

Question number	Answer		Accept	Reject	Marks						
5 (a)	<table border="1"> <tr> <td data-bbox="297 282 678 316">M1 temperature after</td> <td data-bbox="678 282 831 316">27.1</td> </tr> <tr> <td data-bbox="297 316 678 349">M2 temperature before</td> <td data-bbox="678 316 831 349">18.8</td> </tr> <tr> <td data-bbox="297 349 678 424">M3 temperature change</td> <td data-bbox="678 349 831 424">(+) 8.3</td> </tr> </table>	M1 temperature after	27.1	M2 temperature before	18.8	M3 temperature change	(+) 8.3	<p>Recorded temperatures correct but in wrong order scores 1 for M1 and M2 M3 csq on M1 and M2</p>	one trailing zero	more than one trailing zero	3
M1 temperature after	27.1										
M2 temperature before	18.8										
M3 temperature change	(+) 8.3										
(b)	<p>M1 heat (energy) /thermal energy lost (to the atmosphere) ignore just energy lost</p> <p>M2 potassium hydroxide dissolves (very/too) slowly</p>		<p>water evaporates</p> <p>potassium hydroxide does not completely dissolve</p> <p>potassium hydroxide is impure</p> <p>less than 3 g of potassium hydroxide is used</p> <p>more than 50 cm³ of water is used</p>		2						

Total 5 marks

Question number	Answer				Accept	Reject	Marks
6 (a)	Element	Arrangement of electrons in atom	Arrangement of electrons in ion	Charge on ion	K ⁽¹⁾⁺ / K ⁺¹ S ²⁻ / S ⁻² positive for potassium and negative for sulfide for 1 mark		3
			2.8.8	(1)+/+1			
			2.8.8	2-/-2			
M1 – <u>both</u> arrangements correct M2 – charge on potassium ion M3 – charge on sulfide ion							
(b) (i)	<u>ions</u> move/travel (to the electrodes)				<u>ions</u> are free to move / <u>ions</u> are mobile	electrons free to move	1
	(ii)	M1 (electrostatic) forces (of attraction) between (oppositely charged) <u>ions</u> M2 are (relatively) strong M3 large amount of energy required to overcome the forces / separate the ions from the lattice M2 dep on mention of forces (of attraction) or bonds Mention of covalent bonds or intermolecular forces no M1				<u>ionic</u> bonding / <u>ionic</u> bonds break the bonds	

Total 7 marks

Question number	Answer	Accept	Reject	Marks
7 (a)	$\text{H}_2\text{S}_2\text{O}_7 + \text{H}_2\text{O} \rightarrow 2\text{H}_2\text{SO}_4$	multiples and fractions		1
(b)	<p>M1 32 (of S) \rightarrow 80 (of SO_3) (tonnes or g)</p> <p>M2 mass of $\text{SO}_3 = \frac{80}{32} \times 80$</p> <p>M3 = 200 (tonnes)</p> <p>M2 csq on M1</p> <p>M3 csq on M2</p> <p>Correct answer with no working scores 3</p>	<p>M1 $n(\text{S}) = (n(\text{SO}_3)) = \frac{80 \times 10^6}{32}$ (mol) (= 2 500 000 (mol))</p> <p>M2 mass of $\text{SO}_3 = \mathbf{M1 \times 80 (= 200\ 000\ 000\ (g))}$</p> <p>M3 = M2 $\div 10^6 / 200$ (tonnes)</p>		3
(c)	<p>M1 64 (g) (of SO_2) reacts with 12 (dm^3) (of O_2)</p> <p>M2 (64 tonnes) reacts 12×10^6 (dm^3) OR 1.2×10^7 (dm^3)</p> <p>M2 csq on M1</p> <p>Correct answer with no working scores 2</p>	<p>M1 $n(\text{SO}_2) = \frac{64 \times 10^6}{64}$ (mol) (= 10^6 mol)</p> <p>M2 $\frac{\mathbf{M1}}{2} \times 24 / 1.2 \times 10^7$ (dm^3)</p> <p>OR</p> <p>M1 mass of oxygen</p> <p>accept 1.2×10^{10} <u>cm^3</u></p>		2

Total 6 marks

Question number	Answer	Accept	Reject	Marks
8	<p>M1 – add (aqueous) chlorine to (aqueous) KBr</p> <p>M2 – (solution) turns orange</p> <p>M3 – add (aqueous) bromine to (aqueous) KI</p> <p>M4 - (solution) turns brown</p> <p>M5 – $\text{Cl}_2 + 2\text{KBr} \rightarrow \text{Br}_2 + 2\text{KCl}$</p> <p>OR</p> <p>$\text{Br}_2 + 2\text{KI} \rightarrow \text{I}_2 + 2\text{KBr}$</p> <p>Ignore state symbols</p>	<p>yellow / brown</p> <p>red-brown / orange</p> <p>correct ionic equations</p> <p>accept $\text{Cl}_2 + 2\text{KI} \rightarrow \text{I}_2 + 2\text{KCl}$ if chlorine is added to potassium iodide</p>	<p>red</p> <p>yellow</p>	5

Total 5 marks

Question number	Answer	Accept	Reject	Marks
8	<p>M1 – add (aqueous) bromine to (aqueous) KCl</p> <p>M2 – no change</p> <p>M3 – add (aqueous) iodine to (aqueous) KBr</p> <p>M4 - no change / no change</p> <p>If this route is chosen then M5 cannot be scored</p>	<p>orange / yellow / brown solution/colour produced only if it is clear that no reaction has occurred</p> <p>brown / red-brown / orange solution/colour produced only if it is clear that no reaction has occurred</p>	<p>red</p> <p>yellow</p>	5

Total 5 marks

Question number	Answer	Accept	Reject	Marks
9 (a)(i)	shifts to left	moves in the endothermic direction		1
(ii)	shifts to the right	shifts to the side of the reactants OWTTE		1
(iii)	impossible to know which shift is greater / impossible to know which change has the greater effect	moves in the exothermic direction shifts to the side of the products OWTTE shifts to the side with fewer (gas) moles/molecules OWTTE the (two) effects are opposing one another		1
(b)	M1 – greater proportion of NO ₂ M2 – (increase of) temperature has a greater effect than (increase of) pressure	more NO ₂ present equilibrium shifts to left		2

Total 5 marks

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London WC2R 0RL