

Mark Scheme (Results)

Summer 2013

International GCSE
Chemistry (4CH0) Paper 1C
Science Double Award (4SC0)
Paper 1C

Edexcel Level 1/Level 2 Certificate
Chemistry (KCH0) Paper 1C
Science (Double Award) (KSC0)
Paper 1C

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson. Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2013

Publications Code UG037094

All the material in this publication is copyright

© Pearson Education Ltd 2013

Question number	Answer	Notes	Marks
1 (a) i	5		1
ii	11		1
iii	5		1
iv	6		1
v	5		1
1 (b) i	more		1
ii	more		1
iii	the same number of		1
1 (c)	cross in box D (2.8.3)		1
		Total	9

Question number	Answer	Notes	Marks
2 (a) i	cross in box A (zinc sulfate)		1
ii	cross in box B (iron) cross in box C (magnesium)	Apply list principle - 3 crosses = max 1 4 or 5 crosses = 0 marks	1 1
2 (b)	burns with a pop/squeak OR use burning/lit splint/flame to see if pop/squeak	Must be reference to test and result Reference to splint/match with no indication of flame is not enough Reject reference to glowing splint Ignore flame extinguished 'Squeaky pop test' on its own is not sufficient	1
2 (c)	2 (1) 2	Accept multiples and fractions	1
2 (d) i	cross in box 3		1
ii	reversible / can go in both directions / can go backwards and forwards	Ignore references to equilibrium Ignore references to other reaction types (e.g. hydration / oxidation / exothermic) Accept either equation with \rightleftharpoons	1
		Total	7

Question number	Answer	Notes	Marks
3 (a) i	gas / (g) / g	Accept equivalents such as gaseous / vapour Ignore colours	1
ii	darker / dark grey	Accept black Ignore references to states Ignore more intense Reject reference to any other colours	1

Question number	Answer	Notes	Marks
3 (b) i	no reaction (possible) / no displacement OR halogens do not react with their own halide ions	Accept no change Ignore references to lithium chloride containing chlorine / already reacted / OWTTE	1
ii	iodine/it is less reactive than bromine / bromine more reactive than iodine	Accept correct references to positions in (re)activity series Both halogens must be mentioned, except assume it refers to iodine Reject -ide endings Accept symbols and formulae Ignore references to only one element, e.g. iodine is unreactive Reject any comparison involving sodium	1
iii	iodine	Ignore references to states Ignore I and I ₂	1
iv	cross in box D (bromine displaces iodine)		1
v	2KCl + Br ₂	Either order Penalise incorrect symbols/numbers / unconventional formulae in this part, e.g. CL, br, Br ² , ClK	1
		Total	7

Question number	Answer	Notes	Marks
4 a	<p>4 electrons shared between 2 (carbon) atoms</p> <p>4 electron pairs between 2C and 4H atoms</p>	<p>Ignore inner electrons even if wrong Ignore number of hydrogen atoms</p> <p>Accept all permutations of dots and crosses Ignore intersecting circles Accept H atoms at all angles At least one C or one H atom must be labelled – max 1 if not Max 1 if more than 2 C atoms Max 1 if wrong number of electrons in outer shell of any atom</p>	<p>1</p> <p>1</p>

Question number	Answer	Notes	Marks
4 b i	setting out correct division of each % by A_r OR 3.2, 9.7 and 3.2	Award 0/3 if division by any atomic numbers / wrong way up / multiplication used Do not penalise roundings and minor misreads of % values, eg 38 or 39 for carbon If molecular masses used for H and/or O, no M1, but can award M2 and M3 but no CQ in ii Using 2 and 32 gives C_2H_3O Using 1 and 32 gives C_2H_6O Using 2 and 16 gives $C_2H_3O_2$ Working required for these answers	1
	division by smallest /ratio of 1 : 3 : 1	M2 subsumes M1	1
	CH_3O	Accept elements in any order Award 3 for correct final answer with no working No ECF from M2	1
		Accept use of 62 from ii, i.e. $62 \times 0.387 = 24$ etc scores M1 ratio scores M2, answer scores M3	
ii	$C_2H_6O_2$	Accept elements in any order No other answer acceptable	1
		Total	6

Question number	Answer	Notes	Marks
5 a i	haematite	Ignore iron oxide / formulae	1
	ii Al_2O_3 / $\text{Al}^{3+}_2\text{O}^{2-}_3$	Ignore reactants in equation	1
	iii carbon / C (positive/negative) electrode	Accept graphite Ignore coke / anthracite DEP on M1 (including coke or any metal) Accept anode / cathode Accept references to letting electrons/electricity/current flow Accept references to letting electrons/electricity/current in/out of the cell/electrolyte Ignore references to attracting ions Accept references to conducting electricity Ignore references to electrolysis Ignore references to burning or combining with oxygen Ignore references to reduction/oxidation	1 1

Question number	Answer	Notes	Marks
5 (b) i	redox		1
ii	(it/ Al ³⁺ /aluminium ions) gain of electron(s) OR Al ³⁺ combines/reacts with electrons	Reject gain of electrons by Al/aluminium Accept decrease in oxidation state Ignore/ references to gain of oxygen	1
iii	C / carbon gain of oxygen / O / O ₂ OR reacts/combines/joins/bonds with oxygen or O or O ₂ OR reduces water/H ₂ O	Accept graphite/coke DEP on M1 Ignore forms carbon monoxide Ignore references to loss of electrons Accept increase in oxidation state Ignore (acts as) reducing agent	1 1

Question number	Answer	Notes	Marks
5 d i	$\text{Fe}_2\text{O}_3 + 3\text{CO} \rightarrow 2\text{Fe} + 3\text{CO}_2$	M1 for correct reactants and products M2 for balancing M2 dep on M1	1 1
ii	neutralisation	Accept acid-base Ignore all other descriptions	1
iii	slag		1
		Total	17

Question number	Answer	Notes	Marks
6 (a)	C	Accept formula of C	1
6 (b) i	(compound/molecule/substance containing) carbon and hydrogen (atoms/elements)	Reject atom/element in place of compound/molecule Reject compound/molecule in place of atoms/elements Reject mixture	1
	Only	M2 dependent on mention of carbon and hydrogen even if M1 not awarded Accept other terms with same meaning, e.g. solely / exclusively / just	1
	A	M3 independent Accept name/formula of A	1
	ii contains a (C=C) double bond	Accept multiple bond Ignore references to type of compound, eg hydrocarbon Reject double bond between C and H Do not penalise incorrect terms such as atom or element Ignore not all bonds are single Accept can undergo addition reactions Accept does not contain the maximum number of hydrogens/hydrogen atoms	1
	B	M2 independent Accept name/formula of B	1

	Answer	Notes	Marks
6 b iii	(compounds / molecules / substances with) same molecular formula / same number of each type of atom	Ignore same (chemical) formula /same compound No penalty for reference to hydrocarbons Reject same empirical/general formula If atoms or elements instead of compounds or molecules, only 1 of M1 and M2 can be awarded	1
	different structures /different structural/displayed formulae OR atoms arranged differently	Ignore different molecular arrangement	1
	C and F	Accept in either order Accept formulae of C and F	1

Question number	Answer	Notes	Marks
6 c i	<p>same/similar chemical properties/reactions/behaviour/characteristics</p> <p>gradation / gradual change / trend / increase / decrease of physical properties</p> <p>same functional group</p> <p>same general formula</p>	<p>Ignore specific examples such as react with oxygen</p> <p>Ignore similar (type of) reactivity</p> <p>Do not penalise reference to trends</p> <p>Accept reference to specific property, eg boiling point</p> <p>Reject same / similar physical properties</p> <p>Ignore variable physical properties</p> <p>Ignore reference to specific group</p> <p>Accept alkanes have the (general) formula C_nH_{2n+2}</p> <p>Reject same empirical/molecular formula</p> <p>Any two for 1 each</p> <p>Accept two answers on one answer line</p> <p>Ignore any reference to properties not specified as physical or chemical</p>	2
ii	D AND E	<p>Reject any other combinations</p> <p>Accept correct formulae</p>	1

Question number	Answer	Notes	Marks
6 d i	<pre> H H H-C - C-H Br Br </pre>	Ignore bond angles and positioning of Br (as long as one on each C)	1
ii	<pre> H H-C-H H </pre>		1
Total			14

Question number	Answer	Notes	Marks
7 d i	product formulae or names / products (word) above reactants	Horizontal line not needed Ignore formula errors and one or two missing product(s) Ignore curves and intermediates	1
	ii (approximately) vertical line between reactants and products / between two levels AND labelled ΔH / energy change / heat change / enthalpy change	Ignore arrowheads on vertical line Ignore sign of ΔH Mark can be awarded for exothermic reaction Accept 2310 or any other number in place of ΔH	1

Question number	Answer	Notes	Marks
7 (e) i	temperature change = (-)5.5 (°C) heat change (= $100 \times 4.2 \times 5.5$) = 2310/2300 (J)	Award M1 for 5.5 anywhere CQ on candidate temperature change, provided other values correct Accept answer in kJ Ignore signs Correct final answer scores 2 2.31 (J) scores 1 mark if M1 not awarded	1 1
ii	concentration (of vinegar / (ethanoic) acid / CH_3COOH)	Ignore strength Ignore reference to M_r Accept concentration even if in an incorrect expression	1
Total			10

Question number	Answer	Notes	Marks
8 (a)	mass surface area / size / volume	Accept weight Ignore length / width / height / thickness / shape / type Ignore temperature / purity / density	1 1
8 (b)	(2) time / how long (to collect gas) (3) number of / how many (marble) chips (4) volume of gas / how much gas collected (5) percentage / concentration (of acid)	Accept length of reaction/experiment Ignore amount of marble Accept amount of (marble) <u>chips</u> Accept carbon dioxide/CO ₂ in place of gas Accept amount of gas Accept strength (of acid) Ignore volume of acid Reject amount (of acid)	1 1 1 1

Question number	Answer	Notes	Marks	
8 (c)	i	all six points plotted to nearest gridline	Deduct 1 mark for each error up to max 2, including extra points, but no penalty for point at 2.4 and 50%	2
		straight line of best fit	Line need not be extrapolated Must be drawn with a ruler CQ on candidate's plotted points	1
	ii	anomalous point circled	at 1.5, 20 or CQ on candidate's line of best fit	1
	iii	timer started late / stopped early	Ignore just wrong time	2
		concentration/% too high/more than 20%	Ignore just wrong concentration	
	marble chips bigger / more marble chips	Ignore just wrong mass/size		
	air in measuring cylinder before experiment started (room) temperature higher	Accept smaller volume of gas collected		
		Ignore references to calculation Any two for 1 mark each M1 to M5 CQ on position of anomalous point, so if anomalous point below line of best fit, then: M1 timer started too early/stopped too late M2 concentration/% too low/less than 20% M3 marble chips smaller / fewer marble chips M4 larger volume of gas collected M5 (room) temperature lower		
	iv	vertical line from 50% to drawn graph line	Accept short vertical line crossing graph line at 50% Accept point on graph line at 50%	1
		2.4 (cm ³ /s)	CQ on drawn graph line but can be awarded without vertical line Do not penalise incorrect units	1
			Total	14

Question number	Answer	Notes	Marks
9 (a) i	beaker	Ignore qualifiers such as measuring / graduated / 250 cm ³	1
	ii Pipette	Ignore qualifiers such as measuring / graduated / 25 cm ³	1
	iii colour change is gradual /not sharp/not defined OR end point not sharp/defined/accurate	Ignore reference to many colours	1
	methyl orange / phenolphthalein / litmus	Accept other correct indicators Ignore reference to pH meter Reject litmus paper	1
	iv (measures) only one volume / fixed volume / 25 cm ³ has only one graduation mark OR cannot deliver (measured) variable volume OR volume required is not known OR cannot be controlled / hard to control	Ignore named colours and colour changes Ignore reference to accuracy / size Accept reverse arguments based on suitability of burette, eg burette can deliver/measure any volume	1

Question number	Answer	Notes	Marks
9 (b)	(after) 22.60 (before) 2.75 (added) 19.85	CQ on before and after readings Award 1 for before and after values both correct but in wrong order All values must be to 2 dp Penalise answers to other than 2 dp once only	1 1 1

Question number	Answer	Notes	Marks
9 (d) i	divide/÷ by 1000 (not by 100) OR convert volume/cm ³ to dm ³ OR use 1000 instead of 100	Accept $\frac{0.0500 \times 23.60}{1000}$ (= 0.00118 mol) Accept divide (final) <u>answer</u> by 10	1
ii	multiply/× (amount of H ₂ SO ₄) by 2 (not divide by 2)	Accept 0.(0)0118 × 2 (= 0.0(0)236 mol) Accept any other number in place of 0.0(0)118 Accept multiply (final) answer by 4	1
iii	divide by 25.0 (not by 23.60) OR divide by volume of KOH (not by volume of H ₂ SO ₄) OR use 25.0/volume of KOH instead of 23.60/volume of H ₂ SO ₄	Accept $\frac{0.00590 \times 1000}{25.0}$ (= 0.236 mol) Accept any other number in place of 0.00590 Must be positive statement about correction needed, e.g. in iii, ignore statement about volume of H ₂ SO ₄ should not have been used	1
		Total	14

Question number	Answer	Notes	Marks
10 (a)	air / atmosphere		1
10 (b)	i (equilibrium / it) shifts to left/reactants	Accept favours reverse reaction Accept less product/ $\text{CO} + (3)\text{H}_2$ / lower yield Accept more reactant / $\text{CH}_4 + \text{H}_2\text{O}$	1
	fewer (gas) moles/molecules on left OR favours side with fewer moles/molecules	Accept opposite arguments, eg more moles on right Ignore references to volume and pressure Accept references to particles in place of molecules Reject atoms in place of molecules Ignore references to rate / collisions If wrong shift, M2 cannot be awarded If no shift, M2 can be awarded Marks can be awarded for both points in either prediction or reason	1
	ii (equilibrium / it) shifts to left/reactants	Accept favours reverse reaction/less product Accept less product/ $\text{CO}_2 + \text{H}_2$ / lower yield Accept more reactant / $\text{CO} + \text{H}_2\text{O}$	1
	(forward) reaction exothermic OR reverse reaction endothermic	Ignore references to rate If wrong shift, M2 cannot be awarded If no shift, M2 can be awarded Marks can be awarded for both points in either prediction or reason	1

Question number	Answer	Notes	Marks
10 (c)	Increased (rate) Increased (rate)	Accept alternatives such as faster reactions Accept alternatives such as faster reactions Ignore references to equilibrium/yield Ignore references to forward and reverse reactions	1 1

Question number	Answer	Notes	Marks
10 d i	$M_r(\text{NH}_4\text{NO}_3) = 80$ $n (= \frac{40000}{80}) = 500 \text{ (mol)}$	Award M1 for 80 anywhere ECF from incorrect M_r 500 scores M1 and M2 0.5 (mol) scores only M1	1 1
ii	Answer to di / 500 (mol)		1
iii	$M_r(\text{NH}_3) = 17$ $m = \frac{500 \times 17}{1000} = 8.5 \text{ (kg)}$	Award M1 for 17 anywhere, except if result of incorrect calculation ECF from incorrect M_r Moles CQ on ii Accept answer in g if unit is given If no conversion from kg to g in i, and no conversion from g to kg in iii, penalise these errors once only Award 2 marks for 8.5 (kg)	1 1
10 e	phosphorus AND potassium	Accept in either order Ignore symbols Reject phosphate/phosphine	1
		Total	13

Question number	Answer	Notes	Marks
11 b	<p>NITROGEN (simple) molecular structure</p> <p>intermolecular forces (of attraction) / forces between molecules</p> <p>weak / easily overcome /need little energy to overcome</p> <p>SILICA giant (structure)</p> <p>covalent (bonds)</p> <p>bonds strong / hard to break /need a lot of energy to break</p>	<p>Accept (nitrogen/N₂/small/discrete) molecules</p> <p>Accept intermolecular bonds Reference to intermolecular also scores M1</p> <p>If any reference to breaking covalent bonds or any other type of bonding, only M1 can be awarded</p> <p>Accept macromolecular Ignore lattice</p> <p>Covalent can be anywhere in answer</p> <p>If any reference to breaking intermolecular forces or any other type of bonding, only M4 can be awarded Any five for 1 mark each</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
		Total	9
		Total for paper	120

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UG037094 Summer 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
.....



Llywodraeth Cynulliad Cymru
Welsh Assembly Government



Rewarding Learning