

# Mark Scheme (Results)

June 2014

Pearson Edexcel International GCSE  
Chemistry (4CH0) Paper 1C  
Science Double Award (4SC0) Paper  
1C

Pearson Edexcel Level 1/Level 2  
Certificate  
Chemistry (KCH0) Paper 1C  
Science (Double Award) (KSC0) Paper  
1C

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

### **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at:

[www.pearson.com/uk](http://www.pearson.com/uk)

January 2014

Publications Code UG038368

All the material in this publication is copyright

© Pearson Education Ltd 2014

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Accept	Reject	Marks
1 (a)	<b>B</b> - (filter) funnel			1
	<b>D</b> - test tube/boiling tube			1
	<b>E</b> - pipette		teat pipette/dropping pipette	1
	<b>F</b> - beaker			1
(b)	<b>M1</b> - A			1
	<b>M2</b> - E			1

(Total marks for Question 1 = 6 marks)

Question number	Answer	Accept	Reject	Marks
2 (a) (i)	<b>D</b> - hydrocarbons			1
(b)	S U R V T First mark for S in box 1 <u>AND</u> R in box 3 Second mark for V in box 4 <u>AND</u> T in box 5			2

(Total marks for Question 2 = 3 marks)

Question number	Expected Answer	Accept	Reject	Marks
3 (a) (i)	12			1
(ii)	<b>M1</b> – 2  <b>M2</b> – two electrons in <u>outer/valence</u> shell Award M2 if M1 missing but not if incorrect Ignore references to magnesium and 2.8.2	roman numeral		1  1
(iii)	X <sup>2+</sup>	Mg <sup>2+</sup>		1
(b)	<b>M1</b> – (79 x 24) + (10 x 25) + (11 x 26)  <b>M2</b> – divide by <u>100</u>  <b>M3</b> – 24.3  Mark <b>M2</b> and <b>M3</b> csq on <b>M1</b> if one minor slip in numbers in M1 (eg 97 instead of 79 or 25 instead of 24)  <b>M3</b> dep on <b>M2</b>  Correct answer with no working scores 3  IGNORE units	(0.79 x 24) + (0.10 x 25) + (0.11 x 26) for 2 marks  24.32 with no working scores 2		1  1  1

(Total marks for Question 3 = 7 marks)

Question number	Answer	Accept	Reject	Marks
4 (a)	to increase the rate/speed (of the reaction)  IGNORE to start the reaction/to provide energy/references to the copper(II) oxide will not react without heat / to make it dissolve faster / to give particles more energy	to overcome the activation energy/to provide <u>activation</u> energy (for the reaction)	Answers referring to copper instead of copper(II) oxide	1
(b)	it stops disappearing  OR there is a (black) suspension/solid /copper(II) oxide  OR the mixture/it turns cloudy/black IGNORE crystals	stops dissolving  precipitate/ppt	any colour other than black	1
(c)	to remove (unreacted/excess) copper(II) oxide IGNORE references to impurities/crystals	to remove (unreacted/excess) solid  to obtain a solution (of copper(II) sulfate)	to separate copper(II) oxide from sulfuric acid	1
(d)	copper(II) sulfate/the crystals are less soluble in cold water (than in hot water) OR solubility decreases with temperature IGNORE reference to water evaporating	reverse argument <u>ions</u> join together (to form a lattice) <u>ionic</u> lattice forms	references to freezing	1

(e)	blue IGNORE shades of colour		any colour other than blue	1
(f)	on filter paper/kitchen towel/tissue paper OR leave / in a warm place / in the sun / on a radiator / near a window / in a (warm/drying) oven	OWTTE desiccator	heat / hot oven	1

(Total marks for Question 4 = 6 marks)



Question number	Answer	Accept	Reject	Marks
5 (a)	(i) B	lower case letters		1
	(ii) D			1
	(iii) A			1
	(iv) C			1
(b)	<b>M1</b> - (a substance) containing (two or more) elements IGNORE atoms for <b>M1</b> only		mixture for M1 only  molecules/particles bonded, etc for M1 and M2	1
	<b>M2</b> - bonded (together) / <u>chemically</u> combined (in a fixed ratio)			<u>chemically</u> joined
(c) (i)	<b>M1</b> - Na loses electron(s)			1
	<b>M2</b> - Cl gains electron(s)			1
	<b>M3</b> - Na becomes 2.8 AND chlorine becomes 2.8.8  If incorrect number of electrons transferred, max 2  IGNORE references to full shells  max 1 for mention of covalent bonding  All 3 marks can be scored from correct dot and cross diagrams showing electron transfer			1

(ii)	<p><b>M1</b> - Na = 23 <u>AND</u> Cl = 35.5</p> <p><b>M2</b> - 58.5</p> <p><b>M2</b> dep on <b>M1</b></p> <p><b>IGNORE units</b></p> <p><b>Correct answer with no working scores 2</b></p>			<p>1</p> <p>1</p>
------	--	--	--	-------------------


(Total marks for Question 5 = 11 marks)

Question number	Answer	Accept	Reject	Marks
6 (a)	(i) 13(.0)			1
	(ii) 1.4			1
	(iii) 25(.0)			1
(b)	indigo			1
	red			1
(c)	NaOH + HCl → NaCl + H <sub>2</sub> O IGNORE state symbols even if incorrect	H <sup>+</sup> + OH <sup>-</sup> → H <sub>2</sub> O		1

(Total marks for Question 6 = 6 marks)


Question number	Answer	Accept	Reject	Marks
7 (a)	magnesium chloride/MgCl <sub>2</sub> oxygen/O <sub>2</sub> sulfuric (acid)/H <sub>2</sub> SO <sub>4</sub> IGNORE hydrogen sulfate If name and formula given, both must be correct	carbon dioxide/CO <sub>2</sub>		1 1 1
(b)	$\text{Mg} + \text{H}_2\text{O} \rightarrow \text{MgO} + \text{H}_2$ IGNORE state symbols even if incorrect Penalise incorrect symbols and failure to use subscripts			1

(Total marks for Question 7 = 4 marks)

Question number	Answer	Accept	Reject	Marks
8 (a)	 <p><b>M1</b> – for both electron diagrams correct IGNORE inner electrons of N even if incorrect</p> <p><b>M2</b> – for both charges correct</p> <p><b>M3</b> – for correct ratio of ions</p>	any combination of dots and crosses		3
(b)	$6\text{Li} + \text{N}_2 \rightarrow 2\text{Li}_3\text{N}$ <p><b>M1</b> – all formulae correct</p> <p><b>M2</b> – balanced</p> <p><b>M2</b> dep on <b>M1</b></p> <p>IGNORE state symbols even if incorrect</p>	multiples and fractions		2
(c) (i)	<p>(i) l aq g</p> <p>(ii) <b>M1</b> – any number from 8 to 14</p> <p><b>M2</b> – LiOH/lithium hydroxide is a base/alkali <b>OR</b> hydroxide <u>ions</u>/OH<sup>-</sup> formed/present</p>	ammonia / <u>metal</u> hydroxides / <u>Group 1</u> hydroxides are bases/alkalis		1  1  1

(d)	ions cannot move OR ionic compounds <u>only</u> conduct when molten/in solution  IGNORE references to electrons	ionic compounds do not normally conduct when solid		1
-----	---	--	--	---

(Total marks for Question 8 = 9 marks)

Question number	Answer	Accept	Reject	Marks
9 (a) (i)	A	Methane		1
(ii)	C	Ethene		1
(iii)	C	Ethene		1
(b)	<b>M1</b> - (molecular) C <sub>4</sub> H <sub>10</sub> <b>M2</b> - (empirical) C <sub>2</sub> H <sub>5</sub> ECF from molecular formula	H <sub>10</sub> C <sub>4</sub> H <sub>5</sub> C <sub>2</sub>	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	1 1
(c) (i)	<b>M1</b> - (name) alkane(s) <b>M2</b> - (general formula) C <sub>n</sub> H <sub>2n+2</sub>			1 1
(ii)	 <p>IGNORE bond angles</p>		missing Hs and bonds	1

(d)	<b>M1</b> – incomplete combustion/insufficient oxygen	lack of oxygen /less oxygen / <u>only</u> 1½ oxygen (in equation)		1
	<b>M2</b> – toxic/poisonous/causes death IGNORE dangerous/harmful			
	<b>M3</b> – reduces the capacity of the blood to carry oxygen IGNORE references to suffocation/cannot breathe IGNORE blood carries no oxygen	correct references to haemoglobin /blood carries less oxygen/blood does not release oxygen as easily		1

(Total marks for Question 9 = 11 marks)




Question number	Answer	Accept	Reject	Marks
10 (a) (i)	Any two from: <ul style="list-style-type: none"> <li>• good conductor <u>of heat</u></li> <li>• high melting point</li> <li>• malleable</li> </ul> Apply list principle			2
(ii)	<b>M1</b> – ductile  <b>M2</b> – good conductor <u>of electricity</u> Apply list principle Answers can be given in any order			1  1
(b) (i)	strong(er) IGNORE references to density and rusting	other correct descriptions		1
(ii)	lower density / resists corrosion IGNORE lighter	does not rust greater strength to weight ratio		1
(c) (i)	heat / thermal energy / heat energy is given out OR transferred/lost to the surroundings IGNORE references to bond formation and breaking	produced  produces an increase in temperature  it gets hot		1
(ii)	<b>M1</b> - (aluminium/it is) more reactive  <b>M2</b> – (aluminium/it) displaces iron (from its oxide)  M2 DEP on M1	iron is less reactive  replaces it/aluminium takes oxygen away from iron (oxide)		1  1

(iii)	<b>M1</b> – aluminium	loses (three) electrons /oxidation number increases		1
	<b>M2</b> – gains oxygen <b>M2</b> DEP on <b>M1</b> IGNORE references to magnesium	combines with oxygen / forms aluminium oxide		1
(d)	temperature reached $\geq$ m.pt of iron IGNORE exothermic / heat produced / lots of energy produced	<u>high</u> temperature reached / gets <u>very</u> hot		1

(Total marks for Question 10 = 12 marks)

Question number	Answer	Accept	Reject	Marks
11 (a)	large hydrocarbons/alkanes/molecules become small ones IGNORE references to forming alkenes/ethene/more useful molecules	(large) hydrocarbons or alkanes or molecules become smaller ones long chains become short chains	references to polymers	1
(b)	<b>M1</b> – (add to) bromine (water)/Br <sub>2</sub> IGNORE Br  <b>M2</b> – (bromine) decolourised/turns colourless IGNORE starting colour and clear  <b>M2</b> dep on <b>M1</b> , but can be scored for a near miss in <b>M1</b> , eg Br or bromide (water)	(acidified) potassium manganate(VII)  decolourised/turns colourless		1  1
(c)	<b>M1</b> – (catalyst) silica / silicon dioxide / alumina / aluminium oxide  <b>N.B.</b> if both name and formula given, mark the name only  <b>M2</b> – 600-700 °C	correct formula aluminosilicate / zeolite  any value or range within this range equivalent temperatures in Kelvin		1  1

(Total marks for Question 11 = 5 marks)

Question number	Answer	Accept	Reject	Marks
12 (a) (i)	<p><b>M1</b> – divide all the masses by respective <math>A_r</math></p> <p><b>M2</b> – to give 0.02 : 0.02 : 0.04</p> <p><b>M3</b> – (mole) ratio is 1 : 1 : 2 Correct ratio or empirical formula with no working scores 0/3</p>		division by atomic number/division upside down for all marks	1
(ii)	<p><b>M1</b> – <math>204 \div 102 = 2</math> OR <math>102 \times 2 = 204</math></p>	$(2 \times 12) + (2 \times 19) + (4 \times 35.5) = 204$		1
	<p><b>M2</b> – <math>C_2F_2Cl_4</math> Correct answer with no working scores 2 marks</p>	symbols in any order	FI for F	1
(b)	 <p><b>M1</b> – all four bonding pairs correct</p> <p><b>M2</b> – rest of diagram correct</p> <p><b>M2</b> dep on <b>M1</b></p>	<p>FI for F</p> <p>any combination of dots and crosses</p>		2

	IGNORE inner shell electrons even if incorrect Award 1 mark for similar molecules, eg CCl <sub>4</sub> and CF <sub>4</sub>			
--	---	--	--	--

(Total marks for Question 12 = 7 marks)

Question number	Answer	Accept	Reject	Marks
13 (a)	covalent			1
(b) (i)	<p><b>M1</b> – giant covalent / giant structure/lattice/network</p> <p><b>M2</b> – strong (covalent) bonds/many (covalent) bonds</p> <p><b>M3</b> – lot of (thermal/heat) energy required</p> <p><b>M4</b> – to <u>break</u> bonds</p>	macromolecular giant molecular	Max 1 if bonding stated to be intermolecular/ionic/metallic	1 1 1 1
(ii)	<p><b>M1</b> –intermolecular forces(of attraction) / forces (of attraction) between molecules</p> <p><b>M2</b> – are weak / little (thermal/heat) energy required (to overcome the forces)</p> <p>M2 DEP on M1 Weak bonds on its own = 0</p>	intermolecular bonds in place of intermolecular forces	any indication that covalent/ionic/metallic bonds are broken scores 0	1 1
(c)	<p><u>theory B</u> AND since there are no/fewer gas molecules in space</p> <p>OR</p> <p>there is no/less gas in space</p> <p>OR</p> <p>space is a vacuum</p>	<p>fewer gas molecules at high altitude/less gas at high altitude</p> <p>air/specified gas in place of gas</p> <p>ORA</p>		1

(d)	high temperature AND since (forward) reaction is endothermic/absorbs heat  IGNORE references to le Chatelier's principle			1
-----	--	--	--	---

(Total marks for Question 13 = 9 marks)

Question number	Answer	Accept	Reject	Marks
14 (a)	<p><b>M1</b> –</p> <p><math>\left( \begin{array}{cc} \text{H} &amp; \text{H} \\   &amp;   \\ \text{---} \text{C} &amp; \text{---} \text{C} \\   &amp;   \\ \text{H} &amp; \text{H} \end{array} \right)_n</math></p> <p><b>M2</b> – any suitable use, eg:</p> <ul style="list-style-type: none"><li>• plastic bags</li><li>• buckets/bowls</li><li>• storage bottles (for food, drinks, chemicals)</li><li>• garden furniture</li><li>• gas pipes</li><li>• rubbish bins</li><li>• storage tanks for fuel</li><li>• cling film</li><li>• packaging</li><li>• clothing</li><li>• insulation (for electric cables)</li></ul> <p>Please research any unfamiliar use</p> <p><b>M3</b> – poly(propene)</p> <p><b>M4</b> –</p> <p><math>\begin{array}{c} \text{H} &amp; &amp; \text{CH}_3 \\   &amp; &amp;   \\ \text{C} &amp; = &amp; \text{C} \\   &amp; &amp;   \\ \text{H} &amp; &amp; \text{H} \end{array}</math></p> <p>IGNORE bond angles</p>	<p>continuation bonds not going through brackets</p> <p>polypropene polypropylene</p> <p>methyl group attached to any carbon methyl group displayed</p>	<p>just plastic</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>



(b)	Any two from <b>M1</b> – (many) small molecules/monomers join up <b>M2</b> - double bond becomes single bond/ it becomes saturated <b>M3</b> – increase in mass/chain length/size	OWTTE  double bond breaks and single bond forms		2
(c) (i)	inert(ness) IGNORE strong bonds / long chains	unreactive/non-polar		1
(c) (ii)	<b>M1</b> – produces greenhouse gases/toxic gases/poisonous gases <b>M2</b> – (landfill) uses up land / takes up space OR new sites hard to find	carbon dioxide		1

(Total marks for Question 14 = 9 marks)

Question number	Answer	Accept	Reject	Marks	
15 (a) (i)	<b>M1</b> – $M_r(\text{NaOH}) = 40$			1	
	<b>M2</b> – $10(.0) \div M1$			1	
	<b>M3</b> – 0.25 (mol) Correct answer with no working scores 3			1	
	(ii) <b>M1</b> – $0.25 \times 1000 \div 250$			1	
	<b>M2</b> – $1(.0) \text{ (mol/dm}^3\text{)}$ Correct answer with no working scores 2			M3 from (a)(i) $\div 250 / 0.001$ for 1 mark	1
	Mark csq throughout				

(b)	(i)	<p><b>M1</b> – (reading at end) 25.20</p> <p><b>M2</b> – (reading at start) 1.65</p> <p><b>M3</b> – (volume added) 23.55</p> <p>Award 1 mark for correct end and start readings in reverse order</p> <p>Mark <b>M3</b> csq on <b>M1</b> and <b>M2</b></p> <p>Penalise lack of two decimal places once only in a correct answer</p>			1
	(ii)	<p><b>M1</b> – (colour at start) yellow</p> <p><b>M2</b> – orange/pink</p>	red		1
	(iii)	<p>different volumes can be measured /continuously graduated</p> <p>/ addition (of acid) can be controlled</p> <p>/ volume required is not known</p> <p>IGNORE references to precision or accuracy</p>	pipette measures one volume only		1
(c)	(i)	<p><b>M1</b> – <math>2(.00) \times 200 \div 1000</math></p> <p><b>M2</b> – 0.4(0) (mol)</p> <p>Correct final answer with no working scores 2 marks</p>	400 for 1 mark		1
	(ii)	<p><b>M1</b> – <math>n(\text{CO}_2) = 0.2(0) / \frac{1}{2}</math> of <b>M2</b> from (c)(i) (mol)</p> <p><b>M2</b> – <math>\text{mass}(\text{CO}_2) = 8.8(0) \text{ (g)} / \text{M1} \times 44</math></p> <p>Correct final answer with no working scores 2 marks</p>			1

(Total marks for Question 15 = 15 marks)

