



GCSE (9–1) Chemistry A (Gateway Science)

Н

J248/04 Paper 4 (Higher Tier)

Sample Question Paper

Date - Morning/Afternoon

Time allowed: 1 hour 45 minutes

You must have:

· the Data Sheet

You may use:

- · a scientific or graphical calculator
- a rule



First name	
Last name	
Centre number	Candidate number

INSTRUCTIONS

- Use black ink. You may use an HB pencil for graphs and diagrams.
- Complete the boxes above with your name, centre number and candidate number.
- Answer all the questions.
- · Write your answer to each question in the space provided.
- Additional paper may be used if required but you must clearly show your candidate number, centre number and question number(s).
- Do not write in the bar codes.

INFORMATION

- The total mark for this paper is 90.
- The marks for each question are shown in brackets [].
- Quality of extended response will be assessed in questions marked with an asterisk (*).
- This document consists of 32 pages. Any blank pages are indicated.



SECTION A


Answer **all** the questions.

You should spend a maximum of 30 minutes on this section.

1	Which	h statement is correct for a Group 1 element?	
	Α	It dissolves in water to form a bleach.	
	В	It is a non-metal.	
	С	It is an inert gas.	
	D	It reacts with water to form hydrogen.	
	Your	answer	ני

2 The bar chart shows the amount of some of the fractions made from 100 tonnes of crude oil by fractional distillation.

It also shows the amount of each fraction needed for everyday uses.



Cracking converts large molecules into smaller more useful molecules to make the supply match the demand.

Which fractions are most likely to be cracked to make the supply match the demand?

Α	gas	OII	and	tuel	OI
---	-----	-----	-----	------	----

B gas oil and petrol

c naphtha, paraffin and fuel oil

D petrol and gases

Your answer

3	Urea	is a fertiliser.	
	The f	formula for urea is:	
		(NH ₂) ₂ CO	
	A stu	dent makes 1 mole of urea from 2 moles of ammonia.	
	What	t is the mass of urea that the student makes?	
	A	43.0 g	
	В	44.0 g	
	С	58.0 g	
	D	60.0 g	
	Your	answer	[1]
4	A st	udent is testing sodium carbonate solution.	
	She	adds barium chloride solution followed by excess dilute hydrochloric acid.	
	Whi	ch of these observations would not be seen?	
	Α	colourless solution at the end	
	В	gas bubbles when the dilute acid is added	
	С	white precipitate formed when the dilute acid is added	
	D	white precipitate formed when the barium chloride solution is added	
	You	r answer	[1]

5 A student is making a fertiliser called potassium nitrate, KNO₃.

Look at the equation for the reaction she uses.

$$KOH + HNO_3 \rightarrow KNO_3 + H_2O$$

The relative formula masses, $M_{\rm r}$, of each compound are shown in the table.

Compound	Formula	Relative formula mass
potassium hydroxide	KOH	56.1
nitric acid	HNO ₃	63.0
potassium nitrate	KNO ₃	101.1
water	H ₂ O	18.0

What is the atom economy for the reaction to make potassium nitrate?


Assume that water is a waste product.

- **A** 15.1%
- **B** 47.1%
- **C** 52.9%
- **D** 84.9%

Your answ	ver	

[1]

6 Which displayed formula includes the functional group of an alcohol?



Your answer	

[1]

7 Zinc nitrate thermally decomposes to give two gases.

$$2Zn(NO_3)_2(s) \rightarrow 2ZnO(s) + 4NO_2(g) + O_2(g)$$

A student heats 1.89 g of zinc nitrate until there is no further reaction.

What is the total volume of gas, measured at room temperature and pressure, made in this reaction?

Assume that one mole of gas occupies a volume of 24 dm³ at room temperature and pressure.

The molar mass of zinc nitrate is 189 g/mol.

- **A** $0.12 \, \text{dm}^3$
- **B** 0.48 dm^3
- \mathbf{C} 0.60 dm³
- **D** 1.20 dm^3


Your answer

[1]

8 A student investigates the reaction between calcium carbonate and hydrochloric acid.

He measures the volume of gas made every minute.

Look at the graph. It shows his results for the experiment.



What is the rate of reaction between 0 and 2 minutes in cm³/minute?

- **A** 7.5 cm³/min
- B 15 cm³/min
- C 30 cm³/min
- **D** 60 cm³/min

Your answer	

[1]

		ū	
9		ident investigates the reaction between 1.0 g of calcium carbonate and 20 cm ³ of nol/dm ³ hydrochloric acid at 25°C.	
		student does two experiments. He uses different sized pieces of calcium carbonate for experiment.	or
	The	rate of reaction is greater in the first experiment.	
	Whic	ch is the best explanation for this?	
	A	Small pieces of calcium carbonate have a larger surface area resulting in less frequency collisions.	ient
	В	Large pieces of calcium carbonate have a larger surface area resulting in less frequency collisions.	ıent
	С	Large pieces of calcium carbonate have a smaller surface area resulting in more frequent collisions.	
	D	Small pieces of calcium carbonate have a larger surface area resulting in more frequent collisions.	
	Your	answer	[1]
10	Thes	se statements explain how scientists think our modern-day atmosphere was formed.	
	1	Plants evolved and used carbon dioxide during photosynthesis to make oxygen.	
	2	As the Earth cooled down water fell as rain resulting in the formation of the ocean	S.
	3	The atmosphere today consists of nitrogen, oxygen and a small amount of carbon dioxide.	1
	4	Volcanoes gave out ammonia and carbon dioxide as well as methane and water vapour.	
	5	Ammonia was changed by bacteria in the soil into nitrogen gas.	
	Wha	t is the correct order that these events happened?	
	Α	4, 2, 5, 1, 3	
	В	2, 4, 5, 3, 1	
	С	4, 1, 5, 2, 3	
	D	1, 4, 2, 5, 3	
	Your	answer	[1]

11 A student bubbles ethene gas into bromine water.

Which displayed formula shows the product of this reaction?



$$\mathbf{c} \quad \bigvee_{\mathsf{C} = \mathsf{C}}^{\mathsf{H}} \quad \mathop{\mathsf{Br}}$$

[1]

Which of the following procedures is the most suitable for preparing a 0.100 mol/dm³ solution of sodium carbonate?

The relative formula mass, $M_{\rm r}$, of sodium carbonate is 106.

- **A** Dissolving 10.6 g of sodium carbonate in water to make 1.0 dm³ of solution.
- **B** Dissolving 10.6 g of sodium carbonate in 0.10 dm³ of water.
- C Dissolving 10.6 g of sodium carbonate in 1.0 dm³ of water.
- **D** Dissolving 106 g of sodium carbonate in water to make 1.0 dm³ of solution.

Your answer	
rour answer	

[1]

13 A student reacts some metals with different salt solutions and records her results.

She places a tick (\checkmark) in her results table if she sees a chemical change and a cross (x) if there is no reaction.

Some of the boxes are blanked out.

	Magnesium chloride	Silver nitrate	Copper(II) sulfate	Iron(II) sulfate
Magnesium		✓	✓	✓
Silver	х		х	х
Copper	х	✓		х
Iron	х	✓	✓	

Which metal has the least tendency to form a positive ion?

- A copper
- **B** iron
- **C** magnesium
- **D** silver

Your answer	

[1]

14 A student heats compound **X** with acidified potassium manganate(VII) solution.

The product of the reaction is compound Y.

What is the colour change seen during this reaction?

- A colourless to orange
- B colourless to purple
- **C** orange to colourless
- **D** purple to colourless

Your answer	

[1]

15 A condensation polymer is made from two monomers.

One of the monomers has two –OH groups in its molecules.

The other monomer has two –COOH groups in its molecule.

Which one of the following is the polymer?

- **A** polyamide
- **B** poly(chloroethene)
- **C** polyester
- **D** DNA

Your answer

[1]

SECTION B

Answer all the questions.

16 Zinc and dilute sulfuric acid react to make hydrogen.

$$Zn(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq) + H_2(g)$$

Inga measures the rate of this reaction by measuring the **loss in mass** of the reaction mixture.

She finds that the change in mass is very small and difficult to measure.

(a) Draw a labelled diagram to show a **better way** of measuring the rate of this reaction.

[3]

(b) The reaction between zinc and dilute sulfuric acid is slow.

Inga decides to try and find a catalyst for this reaction.

She tests four possible substances.

Each time she adds 0.5 g of the substance to 1.0 g of zinc and 25 cm³ of dilute sulfuric acid.

Look at her table of results.

Substance	Colour of substance at start	Colour of substance at end	Relative rate of reaction
no substance			1
calcium sulfate powder	white	white	1
copper powder	pink	pink	10
copper(II) sulfate powder	blue	pink	30
manganese(IV) oxide powder	black	black	1

(i)	It is important to do the reaction with only zinc and dilute sulfuric acid.	
	Explain why.	
		[1]
(ii)	It is important to do all of the reactions with the same concentration of acid.	
	Explain why.	
		[1]
(iii)	Which of the substances could be a catalyst for the reaction between zinc and dilute sulfuric acid?	
	Explain your answer.	
		[2]
(iv)	There is not enough evidence to confirm which substance is a catalyst. Suggest an extra piece of experimental evidence that could be collected to	
	confirm which substance is a catalyst.	
		[1]

		[O]
	Predict, with reasons, the relative rate of reaction.	
	This time she uses a lump of copper rather than copper powder.	
(v)	Inga does the experiment with copper, zinc and dilute sulfuric acid again.	



BLANK PAGE

PLEASE TURN OVER FOR THE NEXT QUESTION



17 The Group 7 elements are known as the halogens.

The halogens have similar chemical properties.

Their physical properties vary with increasing atomic number.

(a) Look at the table of information about the halogens.

Halogen	Atomic symbol	Atomic number	Molecular formula	Atomic radius in pm	Reaction of halogen with sodium iodide solution
fluorine	F	9	F ₂	64	Makes iodine and sodium fluoride
chlorine	Cl	17	Cl ₂	99	Makes iodine and sodium chloride
bromine	Br	35	Br ₂	114	
iodine	I	53	12	133	No reaction
astatine	At	85		3	No reaction

(iii)	Explain your answer to (ii) in terms of the reactivity of the halogens.	
	Put your answer in the table.	[1]
(ii)	Predict the reaction of bromine with sodium iodide solution.	
	Put your answers in the table.	[2]
(i)	Predict the molecular formula and atomic radius of astatine.	


	(b)	All h	alogens react with alkali metals to make a salt.	
		(i)	All halogens have similar chemical reactions.	
			Explain why in terms of electronic structure.	
			[[1]
		(ii)	Sodium reacts with bromine to make sodium bromide, NaBr.	
			Construct the balanced symbol equation for this reaction.	
			[[2]
				-
		(iii)	What is the formula of the product of the reaction between astatine and potassium?	
				[1]
18	Cha	mical	tests are used to identify gases, anions and cations.	
10				
			an unknown solution.	
	She	thinks	s that the solution contains copper(II) ions and bromide ions.	
	Des	cribe t	the chemical tests she does to confirm the presence of these two ions in the solution.	
				• • •
			[4]

19 Sarah does three titrations with dilute hydrochloric acid and potassium hydroxide solution.


Hydrochloric acid neutralises the alkali potassium hydroxide.

$$HCl(aq) + KOH(aq) \rightarrow KCl(aq) + H_2O(l)$$

Look at the apparatus she uses.



Look at the diagrams. They show parts of the burette during the first titration.



initial reading

final

reading

First titration


Here is Sarah's results table:

Titration number	1	2	3
final reading (cm ³)		37.5	32.1
initial reading (cm ³)		20.4	15.0
titre (volume of acid added) (cm ³)		17.1	17.1

(a)	Use the diagrams and table to help you calculate the mean titre.	
	Explain your answer.	
	Mean titre = cm ³	[2]
(b)	Sarah uses 25.0 cm³ of potassium hydroxide solution, KOH.	
	She also uses hydrochloric acid with a concentration of 0.100 mol/dm ³ .	
	Calculate the concentration, in mol/dm³, of the KOH(aq).	
	Concentration of KOH(aq) = mol/dm ³	[2]
(c)	Use your answer to (b) to calculate the concentration of the KOH(aq) in g/dm ³ .	

20 Poly(propenenitrile) is an addition polymer.

Look at the flow chart. It shows how poly(propenenitrile) is made from crude oil.




(a) Crude oil is a complex mixture of hydrocarbons.

Fractional distillation separates this mixture.

┏	- 1 - :	•	1	- 4	! 1	1 1	£	f., 1: 1	-1! - 1!11 - 1!
-v	าเวเก	ın	Tarme	\sim t	Intarma	IDCIIIOR	TOTODO	tractional	aletillation
ᆫᄭ	viaii i.	111	rem i	OI.	шисппо	ICCUIAL	TULCES.	пачиона	distillation.

[2]

(b) Look at the displayed formula for propenenitrile.



Propenenitrile is an unsaturated compound.

How you can tell from the displayed formula?

.....

21

The	reversible reaction between carbon dioxide and hydrogen makes methane and water.	
	carbon dioxide + hydrogen ⇌ methane + water	
(a)	In a sealed container this reversible reaction forms a dynamic equilibrium.	
	What is meant by the term dynamic equilibrium?	
	Refer to both concentration and rate of reaction in your answer.	
		[2]
(b)	Kayvan investigates this reaction.	
	He predicts that 11.0 g of carbon dioxide should make 4.0 g of methane.	
	In an experiment, he finds that 11.0 g of carbon dioxide makes 2.2 g of methane.	
	Calculate the percentage yield of methane.	
	Percentage yield =%	[2]

(c)*	Kayvan investigates the effect of changing the pressure and changing the temperature on
	this reaction.

carbon dioxide + hydrogen
$$\rightleftharpoons$$
 methane + water $CO_2(g)$ + $4H_2(g)$ \rightleftharpoons $CH_4(g)$ + $2H_2O(I)$

The table shows the percentage yield of methane in the equilibrium mixture under different conditions.

		Pressure in atmospheres			
	100	200	300	400	
	300	35%	52%	65%	80%
Temperature	600	30%	46%	58%	74%
in °C	900	23%	37%	47%	62%
	1200	14%	25%	36%	48%

Kayvan predicts that the reaction between carbon dioxide and hydrogen is endothermic and involves a reduction in the volume of gases.

Describe and explain whether Kayvan's predictions are supported by the reaction and

results in the table.
TE1

Ammonium sulfate can be manufactured from ammonia and sulfuric acid.

(a)		Haber Process is used to manufacture ammonia. Scientists think that the Haber cess is one of the most important chemical reactions.	
	Expl	ain the importance of the Haber Process in agriculture.	
			[2]
(b)	The	Contact Process is used to manufacture sulfuric acid.	
	The	Contact Process involves the reaction between sulfur dioxide and oxygen.	
	The	conditions used are 450°C and about 10 atmospheres pressure.	
	(i)	If the temperature is increased to 500°C the rate of reaction changes.	
		Describe and explain this change in rate of reaction.	
			[2]
	(ii)	If the pressure is reduced to 5 atmospheres the rate of reaction changes.	
		Describe and explain this change in rate of reaction.	
			[2]

It is made using the reaction between the alkali ammonia and sulfuric acid.

$$2NH_3 + H_2SO_4 \rightarrow (NH_4)_2SO_4$$

	214113 1 112004 7 (14114/2004	
(i)	Describe how a sample of solid ammonium sulfate is prepared in a laboratory.	
	Explain why this method is not suitable to be used industrially.	
	[4	i]
(ii)	Predict the maximum mass of ammonium sulfate that can be made from 51 tonnes of ammonia.	
	Maximum mass = tonnes [2	2]

23 Carbon dioxide is one of several greenhouse gases.

It is made by the combustion of fossil fuels such as coal, gas and oil.

Look at the table. It shows the amount of carbon dioxide produced in a large city in the years 2010 and 2016.

Between the years 2010 and 2016 the percentage increase of atmospheric carbon dioxide has been about 2.5%. During the same time, the increase in mean global temperature has been only 0.05°C.

Source of carbon	Carbon dioxide p	Percentage increase (%)	
dioxide	in 2010	in 2016	
Homes	500 000	600 000	20
Factories and industry	500 000	750 000	50
Transport	1 000 000	1 000 000	0
Electricity generation	750 000	900 000	

	(a)	Look at	the row	for elec	tricity gei	neration.
--	-----	---------	---------	----------	-------------	-----------

Calculate the percentage increase of carbon dioxide produced.

	Percentage increase = %	[4]
(b)	Some scientists think there is a link between the amount of fossil fuels burnt and climate change.	
	The data in the table does not support this view.	
	Suggest reasons why.	

[2]

24 Kasia investigates the corrosion of different metals.

She places a small strip of each metal in different samples of air.

She leaves the metals for one week before collecting her results.

Look at her table of results.

	Original	Appearance of metal after one week in			
Metal	appearance of metal	moist acidic air	moist alkaline air	dry air	moist air
aluminium	shiny silver	dull silver	dull silver	shiny silver	shiny silver
copper	shiny red- orange	dull red- orange	green red- orange	shiny red- orange	dull red- orange
iron	shiny silver	brown coating	brown coating	shiny silver	brown coating
magnesium	shiny silver	whitish coating	dull silver	shiny silver	dull silver
zinc	shiny silver	dark coating	dark coating	shiny silver	dull silver


(a)	Suggest, with a reason, one change to the experimental procedure that would improve the quality of the results.	ne
		[1]
(b)	Explain the conclusions that can be made from Kasia's results.	
		[3]

25 Aluminium is extracted from its ore using electrolysis.

Сор	per is	extracted from its ore by heating with carbon.				
(a)	Explain why different methods are used to extract aluminium and copper.					
			[2]			
(b)	Molten aluminium oxide contains Al^{3+} and O^{2-} ions.					
	The electrolysis of molten aluminium oxide makes aluminium and oxygen.					
	(i)	Write the balanced symbol equation for the electrode reaction that happens at the cathode.				
		Use the symbol e ⁻ to represent an electron.				
			[1]			
	(ii)	Solid aluminium oxide cannot be electrolysed.				
		Explain why.				
			[1]			

(c) Copper is also made by electrolysis of copper sulfate solution.

Look at the diagram of the apparatus used in this electrolysis.



Describe what you would **see** at each of the electrodes.

At the anode:	
A	
At the cathode:	

[2]

26	Iron	rusts	when	it	gets	wet.
----	------	-------	------	----	------	------

(a) The word equation for rusting is

iron + water + oxygen
$$\rightarrow$$
 rust (hydrated iron(III) oxide)

Balance the symbol equation for the formation of rust.

.....Fe(s) +
$$H_2O(I)$$
 + $O_2(g)$ \rightarrow $Fe_2O_3.3H_2O(s)$ [2]

(b) A 1.0 kg iron bar is left outside in the rain.

All of the iron turns to rust.

The rust forms at a rate of 60 g per day.

Calculate how long it will take for the iron bar to turn completely to rust.

Give your answer to the nearest day.

END OF QUESTION PAPER